

Edited by:

WIEBKE KIRLEIS, MARTA DAL CORSO, DRAGANA FILIPOVIĆ

MILLET AND WHAT ELSE?

The wider context of the adoption of millet cultivation
in Europe



MILLET AND WHAT ELSE?

MILLET AND WHAT ELSE?

The wider context of the adoption of millet cultivation
in Europe

Edited by:

WIEBKE KIRLEIS, MARTA DAL CORSO, DRAGANA FILIPOVIĆ

© 2022 Individual authors

Sidestone Press Academics

Published by Sidestone Press, Leiden
www.sidestone.com

Layout & cover design: CRC 1266/Anna Carina Lange and Sidestone Press

Cover image: Wiebke Kirleis and Anna Carina Lange

ISBN 978-94-6427-015-0 (softcover)

ISBN 978-94-6427-016-7 (hardcover)

ISBN 978-94-6427-017-4 (PDF e-book)

The STPAS publications originate from or are involved with the Collaborative Research Centre 1266, which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation; Projektnummer 2901391021 – SFB 1266).

Contents

Preface by the series editors	7
INTRODUCTION: Millet and what else? The wider context of the adoption of millet cultivation in Europe Wiebke Kirleis, Dragana Filipović, Marta Dal Corso	9
SECTION 1: INNOVATIONS IN THE BRONZE AGE SUBSISTENCE ECONOMY: REGIONAL PERSPECTIVES FROM UKRAINE TO FRANCE	25
The (pre)history of common millet (<i>Panicum miliaceum</i>) in Ukraine and its place in the traditional cuisine Galyna Pashkevych	27
Bronze Age plant spectra in Hungary before and after the introduction of millet cultivation Sofia Filatova	39
Bronze Age novelties in animal exploitation in the Carpathian Basin in a European context László Bartosiewicz	69
Unearthing millet in Bronze and Iron Age Croatia Kelly Reed, Jacqueline Balen, Ivan Drnić, Sara Essert, Hrvoje Kalafatić, Marija Mihaljević, Emily Zavodny	95
The earliest finds of millet and possible associated changes in material culture in Slovenia Tjaša Tolar and Primož Pavlin	107
On the 'ancient' evidence for <i>Panicum miliaceum</i> and <i>Vicia faba</i> in central Germany (primarily Saxony-Anhalt) Monika Hellmund	129
Millet in Bronze Age agriculture and food consumption in northeastern France Françoise Toulemonde, Julian Wiethold, Emmanuelle Bonnaire, Geneviève Daoulas, Marie Derreumaux, Frédérique Durand, Bénédicte Pradat, Oriane Rousselet, Caroline Schaal, Véronique Zech-Matterne	155

On-site to off-site: A multidisciplinary and multiscale consideration of the 13th to 11th century BCE transformation in northern Germany	185
Ingo Feeser, Stefanie Schaefer-Di Maida, Stefan Dreibrodt, Jutta Kneisel, Dragana Filipović	
SECTION 2: APPROACHING MILLET CULTIVATION AND CONSUMPTION THROUGH HIGH-END MICROSCOPY, CHEMISTRY AND ETHNOGRAPHY	217
Putting millet into a culinary context: Organic residue analysis and the identification of <i>Panicum miliaceum</i> in pottery vessels	219
Edward A. Standall, Oliver E. Craig, Carl Heron	
Tracing millet through biomarker analysis in archaeological sites in alluvial plains: The first miliacin data from the northern Italian Bronze Age	231
Marta Dal Corso, Marco Zanon, Carl Heron, Mauro Rottoli, Michele Cupitò, Elisa Dalla Longa, Wiebke Kirleis	
Exploring seed impressions within the fabric of pottery: Using a silicone cast method for reliable identification	253
Eiko Endo	
Traditional millet cultivation in the Iberian Peninsula: Ethnoarchaeological reflections through the lens of social relations and economic concerns	263
Andrés Teira-Brión	
Versatile usage of millet: Brooms and animal fodder from <i>Sorghum technicum</i>	279
Wiebke Kirleis and Marta Dal Corso	
CONCLUSION: Early cultivation of millet in Europe: What else and where next? Concluding the workshop proceedings	293
Dragana Filipović, Marta Dal Corso, Wiebke Kirleis	
List of authors	321

Preface by the series editors

With this book series, the Collaborative Research Centre 1266 *Scales of Transformation: Human-Environmental Interaction in Prehistoric and Archaic Societies* (CRC 1266), at Kiel University, enables collective and timely presentation of its research outcomes relating to the multiple aspects of socio-environmental transformations in ancient societies. As the editors of this publication platform, we are pleased to be able to publish monographs containing detailed primary data and comprehensive interpretations from different case studies and landscapes, as well as the output from scientific meetings and international workshops. The book series is dedicated to the fundamental research questions of the CRC 1266 related to transformations on different temporal, spatial and social scales, here defined as processes leading to a substantial and enduring reorganisation of socio-environmental interactions. Some of the general questions are: What are the transformations that describe human development from 15,000 years ago to the beginning of the Common Era? How did interactions between the natural environment and human populations change over time? What role did humans play as cognitive actors in the changing social and natural environments? Which factors triggered transformations that led to marked societal and economic inequality?

Understanding human practices within the often intertwined social and environmental contexts is one of the most fundamental aspects of archaeological research. Moreover, in current debates, the dynamics and feedback involved in human-environmental relationships have become a major issue, particularly when we look at the detectable and sometimes devastating consequences of human interference with nature. Archaeology, with its long-term perspective on human societies and landscapes, is in the unique position to trace and link comparable phenomena in the past, to study human involvement with the natural environment, to investigate the impact of humans on nature, and to outline the consequences of environmental change on human societies. Modern interdisciplinary research enables us to reach beyond simplistic, monocausal lines of explanation and overcome evolutionary perspectives. Looking at the period from 15,000 to 1 BCE, the CRC 1266 takes a diachronic view in order to investigate transformations in the development of Late Pleistocene hunter-gatherers, horticulturalists, early agriculturalists, and early metallurgists, as well as early state societies, thus covering a wide array of societal formations and environmental conditions.

The volume *Millet and what else? The wider context of the adoption of millet cultivation in Europe* brings into focus the contribution of dietary innovations to socio-environmental transformation processes, here in the context of the Bronze Age, which is characterised by increasing social complexity and intensified global interconnectedness. Broomcorn/common/proso millet (*Panicum miliaceum*) has recently become a major topic of interest in archaeobotanical as well as

archaeological research in Europe and Asia. Its origin in the far-from-Europe East Asia, its 'late' arrival in Europe (compared to *e.g.* wheat and barley), its distinctive biological traits, and the multiple available methods for tracing this species in the archaeological record make millet an exciting and attractive object for research. This edited volume sets an important benchmark for the current state of the art in millet-related research. It is the outcome of the CRC 1266 workshop 'Millet and what else? The wider context of the adoption of millet cultivation in Europe', which was held in November 2019 in Kiel and was aimed to inspire a deeper discussion on the circumstances and consequences of the introduction of broomcorn millet cultivation in Europe in the Bronze Age. The workshop also marked the closing of the CRC 1266's 'Millet Dating Programme' – the large, collaborative effort of the European archaeobotanical community, which was initiated by CRC 1266's archaeobotanists and led by Dragana Filipović. This volume presents and elaborates the topics and insights presented and discussed at the workshop, and raises new questions. It thus serves as a roadmap for future research and collaboration in the studies of millet, and agricultural or dietary innovations in general.

Both this volume and the workshop from which it proceeds cover the areas and topics investigated by the CRC 1266's subprojects *F3: Dynamics of Plant Economies in Ancient Societies*, and *D1: Population Agglomerations at Tripolye-Cucuteni Mega-sites*. We are grateful in particular to Dragana Filipović for the editorial work, carried out jointly with Marta Dal Corso and Wiebke Kirleis. We cordially acknowledge the support of the external reviewers. We are very thankful to our graphic illustrators, Anna Carina Lange and Carsten Reckweg, for their valuable help in the preparation of the publication. We also thank Karsten Wentink, Corné van Woerdekom and Eric van den Bandt, of Sidestone Press, for their unfailing support in completing this volume, and to Nicole Taylor, scientific coordinator of the CRC 1266, for overseeing the publication process.

Wiebke Kirleis and Johannes Müller

Millet and what else? The wider context of the adoption of millet cultivation in Europe

Wiebke Kirleis, Dragana Filipović, Marta Dal Corso

Introduction

This small-grained cereal – broomcorn/common or proso millet (*Panicum miliaceum*) – has in the past 10-15 years gained massive attention within the field of archaeology, as well as in discussions on modern global food security and sustainable agriculture. It can be considered a special crop in Europe, different from the Neolithic ‘founder crops’ in that it was domesticated in the Far East, namely China, and arrived in Europe ‘late’ – not before the full establishment of metallurgy, in the Bronze Age, when trade and exchange and production levels also peaked. Its distinctive biological, nutritional and ecological traits made this species particularly attractive to farmers in recent and modern communities and, together with its ‘special’ pathway of spread from its region of origin, likely contributed to its widespread and relatively quick adoption by communities in the past. As both previous and new research shows, including that presented in the book in front of us, broomcorn millet is an excellent example of a foodstuff that acted as a social agent in the (trans)formation of society, economy, culture and identity.

Broomcorn millet cultivation and its biological, ecological and nutritive traits

Taxonomically, broomcorn millet (Fig.1) is part of the Poaceae family and the Panicoideae subfamily. Broomcorn millet (henceforth referred to as ‘millet’) emerged as a cultivar in the Far East and has since been commonly used as a crop grown for food, beverage and forage. The primary cultivation areas of this small-grained cereal adapted to short-day cycles are found in the dry tropics or the subtropics. Millet is warmth loving, drought resistant and sensitive to frost. It cannot cope with temperatures below –2°C, and the germination requires ca. 8-10°C. The most important growth factor is the availability of sunlight and water during



Figure 1. Broomcorn millet (*Panicum miliaceum*) before harvest, in the Archäologisch-Ökologisches Zentrum Albersdorf, Schleswig-Holstein, end of August 2018 (photo: Wiebke Kirleis).

the germination time. Millet can grow on poor soils that warm up easily. However, due to the small size of the grains, fine seedbeds are beneficial (Becker-Dillingen 1927, 563-575; Lieberei and Reisdorff 2007; Miedaner and Longin 2012, 80-82). Millet has a rather wide ecological tolerance and can grow in higher altitudes as well as in floodplains (Chen *et al.* 2014; Moreno-Larrazabal *et al.* 2015; Miller *et al.* 2016; d'Alpoim Guedes 2018). However, millet cannot grow beyond 60° northern latitude (Motuzaite-Matuzeviciute pers. comm.). The northernmost archaeobotanical evidence originates from Slavonic sites in Novgorod and Staraja Ladoga, in Russia, and dates to medieval times (Alsleben 2012). The short growing cycle of millet, lasting only 60-90(100) days from seeding to harvesting, makes this species attractive in multiple ways: 1) In northerly regions, it allows millet to be used as a buffer crop to mitigate failure of the primary crops destroyed by late frosts, thus securing a summer crop yield. 2) In Mediterranean climates, it allows millet to be used as second (summer) crop in order to enhance the yearly crop yield (Taylor and Doudu 2020). 3) At higher altitudes which offer a very short vegetation period, it may render millet

superior to other crops (*e.g.* nowadays, broomcorn millet is grown as a summer crop at 3500 m asl. in the Himalayas, as noted by workshop participant Stefania Grando, International Crops Research Institute for the Semi-Arid Tropics, ICRISAT (pers. comm.)). 4) Among semi-sedentary or nomadic groups, such as mobile pastoralists, it also makes millet appealing since it fits with short-term camp occupation (Hermes *et al.* 2019; Dal Corso *et al.* 2022).

The **labour investment** for growing millet varies over the growing period (Fig. 2). Since a fine seedbed is beneficial for plant growth, the soil has to be tilled (ploughed and harrowed or raked) before sowing. Further maintenance is necessary when the plants begin to sprout and push out 3-5 leaves. Because the early leaf production is slow and the faster-growing weeds may outcompete the young millet plants simply by casting shade over them, intensive weeding is necessary (Miedaner and Longin 2012, 80). The weeding has to be carried out with utmost care because the predisposition for the yield is established in the early, 3-5-leaf stage of the plant's growth. Later in the growing period, hardly any labour input is needed.

However, millets are beloved fodder for birds prior to (and post-) harvesting. The plant thus requires effective protection measures that, again, may be labour-intensive. Another issue with millet is the uneven ripening of the grains in the panicle, whereby the ripening starts at the top of the panicle. It takes an experienced farmer to determine the optimal harvesting time and thus avoid grain loss. Despite being high maintenance, millet is an attractive crop because of its high propagation rate compared with *e.g.* rye, with high ratios of planted to harvested grain, even though the grain size is half of that of large-seeded cereals, such as rye or wheat. Further, the silica-rich glumes protect the grains from mildew and allow for long-term storage (Becker-Dillingen 1927, 572-575; Anderson and Greb 1987; Miedaner and Longin 2012, 80-82; Taylor and Duodu 2020).

Millets are adapted to hot and dry environments by following a **C4 pathway** of photosynthesis, which gives them high assimilatory power to produce biomass by reduction of CO₂. Plants normally close their stomata when the ambient temperature is high in order to limit water loss through evaporation. However, this makes the absorption of CO₂ for photosynthesis more difficult. C4 plants have developed a mechanism that enables them to use even the smallest amounts of CO₂ to produce biomass. This comes with increased energy costs, but these costs are buffered by the C4 plants' unique leaf anatomy, which allows CO₂ to concentrate in the so-called bundle sheath cells.

In C4 plants, the fixation of CO₂ is a two-step process. CO₂ is pre-fixed in the mesophyll cells close to the open stomata, where the enzyme PEP is transferred into a four-carbon molecule (*e.g.* malate). The latter is then actively transported with extra energy supply into the so-called bundle sheath cells, where CO₂ is released again and – also when stomata are closed – incorporated into the Calvin cycle to be fixed in sugar molecules (glucose). C4 plants are thus able to continue CO₂ fixation when the highest light intensity of the day is reached (around noon) and stomata are closed, while C3 plants fall back at a light intensity of 50 klx. In full sunlight, C4 plants have higher CO₂ assimilation rates above 30°C than C3 species and are thus better adapted to non-shaded (sub-)tropical habitats. Further, C4 plants have a high water and nitrogen use efficiency (Sitte *et al.* 1991, 277-288; Brown 1998, 479-485). These traits are particularly relevant when millet is grown in subtropical and tropical environments.

But is there a benefit to growing the C4 plant millet in the temperate climatic zone? Does millet allow for higher biomass production and higher yields if compared with C3 plants in central and northern Europe, with its temperate climate? The limiting factor for plant growth in higher latitudes is the availability of light. However, the uneven distribution of biologically effective global irradiance has only minor influence on biomass production (Geisler 1988, 83).



Figure 2. 2a-d – Broomcorn millet (*Panicum miliaceum*) planted on 6 May 2018, in the Archäologisch-Ökologisches Zentrum Albersdorf, Schleswig-Holstein: 2a-2b. Early growth, 30 May 2018; 2c-d. One week later, 6 June 2018; 2e-f. Late harvesting, 29 August 2018. 2g-2h – Broomcorn millet (*Panicum miliaceum*) planted on 27 April 2019, in the Archäologisch-Ökologisches Zentrum Albersdorf, Schleswig-Holstein: 2g. Early growth and intensive weeding, 24 May 2019; 2h. Weeds outcompeted, 21 June 2019 (photos: 2a-d Marta Dal Corso; 2e-f Angelika Hoffmann; 2g-h Wiebke Kirleis).

The observation that C4 plants have a higher net-photosynthesis efficiency than C3 plants and are more productive with respect to biomass production is valid only if C4 and C3 plants are compared under their respective optimal growing conditions. Direct comparison under temperate climate conditions does not allow us to infer variability in biomass production as being stimulated by the different pathways of photosynthesis (Brown 1998). This argument is further supported by the fact that there are hardly any C4 plants among the native wild plant species in temperate central Europe (Brandes 1995; Pyankov *et al.* 2010). What can we conclude from these observations on C4 plant metabolism? And what is the relevance of this for the northward spread of millet in the 2nd millennium BCE?



Figure 3. Illustration from the fairy tale 'Der Süße Brei', by the brothers Grimm (Maximumkarte, DDR 1985, Verlag Bild und Heimat).

In temperate regions, the rate of evapotranspiration through the leaf surface is low, and plants rarely suffer from drought because humidity in the central and northern parts of Europe is relatively high compared with the Mediterranean climate zone in southern Europe. The C4 metabolism can hardly count as a trigger for the northward spread of early millets. Whether weather extremes, such as drought events, fostered the spread of millet is still a desideratum for future research. In modern times, it has been observed that non-plant factors, such as soil type and fertility, length of the growing season, and disease and insect pressures, exert a strong influence on the selection and production of crops and that the selection of crop species by farmers is based more on economic concerns and (culinary) tradition than on biological traits (Brown 1998, 486).

From a **dietary perspective**, millet is rich in the three main macro-nutrients. The carbohydrate content (measured in % of the dry weight) is, as is typical for cereals, high, with the reported values ranging between 60% and 75% (e.g. Saha

et al. 2016; Das *et al.* 2019). The protein content varies between 8% and 13% and can be improved through plant breeding to be as high as 18%. The fat content is about 4-6%. Millet is a gluten-free cereal, easy to digest, and high in dietary fibre and antioxidants, the latter reducing the risk of cardio-vascular diseases (Stefania Grando pers. comm.). The iron content of millet is three times higher than that of wheat, meaning that daily iron requirements of humans can be met with only 50g of millet grains. The optimal nutritional uptake of iron, however, depends on the parallel consumption of vitamin C. Millet contains vitamin E and provitamin A and is particularly rich in vitamin B and minerals. It is in many ways beneficial for human metabolism and health.

Millet is today often promoted as a health-improving product, with benefits for hair, nails and connective tissue because the plant has a high silica content. However, the silica is embedded in the glumes and thus not part of human diet, because it is the dehusked (*i.e.* glume-free) grains that are consumed (Miedaner and Longin 2012, 83; Taylor and Duodu 2020). The hard glumes are essential when it comes to the storage of millet, however, as they perfectly protect the grains from mildew.

During cooking, millet grains gain in size considerably when soaked in hot water or boiled. Thus, although it is a small-seeded cereal, with far lower grain weight than large-seeded cereals (the mass of 1000 grains of millet is 4-9 g and of wheat, 40-50 g; Miedaner and Longin 2012, 80), millet is perceived as a highly filling food. This perception has even found its way into cultural memory through fairy tales. The remarkable swelling of millet grain in food preparation is, for instance, described in 'Der Süße Brei' ('Sweet Porridge') tale by the brothers Grimm (Fig. 3). The main foods that can be made from millet are porridge, soup, flatbread and beer.

Sociocultural dimensions of millet consumption

The cultural dimensions associated with millet and the meanings ascribed to this staple crop are discussed in ethnohistorical studies on the recent past in central Europe (Wiegelmann 1967, 112-156), as well as in ethnological studies on modern-day India (*e.g.* Erler *et al.* 2020; Hardenberg 2021). These studies show that the sociocultural dynamics of foodstuffs in general, and millet in particular, are multidimensional. They are related to and affect societal reality and are evident in the ritual and religious spheres as well as the economic sphere. Some crops have individual agency and play a role in ideology and other cultural behaviours (*e.g.* Hastorf 2017); they are highly dynamic and transformative.

In the 19th century, the cultivation of millet in central Europe rapidly declined. The reasons for this are manifold, and to understand the context, it should be emphasised that, at the time, the production of millet served two highly divergent purposes. It is astonishing that one and same cereal, millet, was attributed two entirely opposite meanings – as a highly valued grain and a symbol of fortune and as food for the poor. On one hand, it was an everyday food for the poor and was widely used in communal kitchens and military hospitals, while it also served as famine food. On the other hand, it was a component of festive meals and was prepared for Christmas, New Year's Eve and weddings, because of the belief in inherent happiness related to the property of millet to 'grow' when made into gruel, which was perceived as a signal that it provides endlessly increasing wealth.

The traditional inclusion of millet in festive meals continued through the 19th century, but as an everyday food it was replaced by potato, especially in areas where millet was less important. One reason for this was a change in consumption habits. At the time, millet was consumed as gruel only, while potato offered a wide range of applications, and was prepared as a bake, gruel, soup or salad, and turned

into flour for making dumplings and pancakes. The change in culinary choice from millet to potato as the main staple occurred alongside the increasing influence and spread of urban eating habits. These entailed new ways of cooking and eating, such as combining different foods into complex meals instead of preparing simple stews, and the increasing use of individual plates with fork and knife, which supplanted earlier, collective eating from one bowl. Further, increasing meat consumption superseded millet gruel. With respect to cultivation, the wider ecological amplitude of potato, which is able to cope with late frosts, for instance, was another of its advantages that accelerated the decline of millet cultivation and consumption in the 19th century (Wiegelmann 1967, 86-156).

Hardenberg (2021) studied cultural dimensions of millet consumption among the Dongria Kond, an Adivasi tribal group living in the Eastern Ghats mountain range, in India. The Dongria Kond traditionally grow millet and ascribe to it a number of different values and meanings. Millet is considered extremely nutritious; it is given economic value because it is used as a currency in exchange for labour, as a kind of wages; it is also an appropriate gift that strengthens social relationships; finally, it serves as a suitable offering in religious contexts. Many of these cultural ascriptions were transferred to rice, which became a cash crop after it had been introduced in the region as a substitute for millet as part of the programme run by development aid organisations. Only recently have millets been promoted as the 'Forgotten Food for the Future' by the 'Smart Food Initiative' of the ICRISAT, as explained by Stefania Grando at our workshop. For people living in Africa and India, the re-introduction of millet growing is part of a programme aimed at furthering sustainable agriculture.

This promotion of a 'forgotten' crop leads to a politically driven transformation of the role of the hill tribes in northern India, formerly perceived as poor people who had to produce rice to survive, but now becoming custodians of endangered millet species thanks to their traditional knowledge of millet cultivation. This strengthens the cultural identity of the Dongria Kond, and they are now re-instating their traditional lifeways. If the practice of exchange of millets traditionally served to consolidate social bonds, it now also has the role of preserving and enhancing the biodiversity of millets.

This tale shows similarities with the study by Erler *et al.* (2020), who explore the fact that, in today's city of Bengaluru, in India, millets are marketed as a superfood made in India, whereby their positive effects on health and their contribution to optimising body weight are emphasised. As a result of this new approach, millets are now shopped for by middle-class people, in organic food shops. This is a noteworthy transformation of attitude since, traditionally, millets had been perceived as inferior crops compared with rice and wheat, representing food of the poor. The new image as superfoods turned millets into a symbol of social distinction. Buying and consuming millet has become a status determinant for the middle class. Although impoverished farmers in India often perceive the consumption of (finger) millet as a consequence of their food scarcity, the members of Bengaluru's middle class consume (a variety of) millets in order to secure their social standing.

These and similar ethnological studies provide examples of changes in culinary practices being deeply intertwined with social transformations – a perspective difficult to explore for prehistory. Although the superseding of millet by potato in the 19th century was stimulated by the wider ecological plasticity of the latter, it was underpinned by the adaptation of rural populations to bourgeois customs and morals and lifestyles, which found its expression in new culinary practices. The examples from India show how the consumption of millet creates, fosters or transforms individual and/or communal identity depending on the habitus and the socio-political context. These changes are in this case almost completely independent from any environmental affordances or constraints. This observation becomes highly relevant when we try to understand causes and consequences of the fast acceptance of broomcorn millet as a new staple crop in Bronze Age Europe.

Millet in the frame of research of the CRC 1266 'Scales of Transformation'

The spread of millet cultivation is one of the central themes of the subproject 'Dynamics of Plant Economies in Ancient Societies' in the research programme CRC 1266 'Scales of Transformation – Human-Environmental Interaction in Prehistoric and Archaic Societies' at Kiel University, which investigates transformational processes in a wide array of societal formations and environmental settings (Müller and Kirleis 2019). The project is investigating the role of millet in the substantial and enduring reorganisation of socio-environmental interaction in prehistoric Europe. It searches for patterns and indicators in crop cultivation and food consumption that reveal the importance of this new resource in the wider processes of cultural and environmental transformation.

In order to place the research and discussions on a firm spatiotemporal footing, the project executed a large, continental-scale radiocarbon dating programme targeting the 'earliest' millet grains in Europe. The dating results proved an earlier assumption (e.g. Hunt *et al.* 2008) that broomcorn millet arrived in Europe in the Bronze Age (Filipović *et al.* 2020). The workshop 'Millet and what else? The wider context of the adoption of millet cultivation in Europe', held in Kiel in November 2019, marked the closing of the 'Millet Dating Programme' and celebrated this major collaborative work of the European archaeobotanical community (Fig. 4). It also provided a roadmap for future research and collaborations, and this book is a step in that direction.

More than 20 invited workshop participants – archaeobotanists, zooarchaeologists, isotope and biomarker specialists, and ethnographers – presented and discussed dietary, economic and cultural aspects of the period in which the new crop was introduced in Europe, i.e. the Middle Bronze Age and the Late Bronze Age, the methods used in archaeology to track and reconstruct the growing and consumption of millet, and the growing and consumption of millet today or in the recent past. The workshop was organised in three sessions.

In the first session, titled 'Innovations in the Bronze Age subsistence economy – regional perspectives from Ukraine to France', the presenters addressed a series of questions suggested by the workshop organisers:

- Why did broomcorn millet spread in the 2nd millennium BCE, and why then? What were the pre-conditions (social, environmental, technological)? Did millet arrive as both human and animal (e.g. chicken) food?
- What was food production (plant and animal) in Europe like before and after the arrival of millet? What was the role of gathered resources?
- Once adopted, how, when and where was millet cultivated, processed, stored, consumed?
- What else changed at the time? What was the role of millet in these changes?

In the second session, titled 'Beyond seeds: Isotopic and biomolecular evidence of millet cultivation and consumption', participants presented new, state-of-the-art methods, such as stable isotope and biomolecular analyses, which detect millet cultivation, cooking and consumption directly in the soil, ceramic vessels and consumers' tissue, while also pointing to a specific yield-securing/improving cultivation practice, i.e. manuring.

In session 3, titled 'Growing and eating millet: Insights from ethnography, agronomy and experimentation', participants shared observations of recent or present-day millet cultivation, processing, storing and uses (as food or fodder); the agricultural routine and amount of labour required in millet cultivation; and the economic and nutritional status of millet in relation to other crops.

The workshop programme and the abstracts are available at <https://www.sfb1266.uni-kiel.de/en/events/sessions-workshops/millet-and-what-else-the-wider-context-of-the-adoption-of-millet-cultivation-in-europe>

WEDNESDAY, 27th NOVEMBER 2019		
08:00 – 08:30	Arrival of participants to the venue	
08:30 – 09:00	Welcome & Opening talk Wiebke Kirleis & Dragana Filipović (CAU Kiel)	
	SESSION 1: Innovations in Bronze Age subsistence economy – regional perspectives from Ukraine to France Session chairs: Wiebke Kirleis & Marta Dal Corso (CAU Kiel), Almuth Alsleben (Akademie der Wissenschaften und der Literatur, Mainz), Susanne Jahns (Brandenburgisches Landesamt für Denkmalpflege und Archäologisches Landesmuseum, Zossen)	
09:00 – 09:20	Millet finds in the territory of Ukraine Galyna Pashkevych (NASU, Kiev), Marta Dal Corso & Wiebke Kirleis (CAU Kiel)	
09:20 – 09:40	Chasing Chinese millets in Ukraine using seed impressions in pottery Eiko Endo (Meiji University, Tokyo)	
09:40 – 10:00	Millet cultivation in the Belarus and the Baltic States Giedrė Motuzaite Matuzevičiūtė & Mindaugas Grikpėdis (Vilnius University)	
10:00 – 10:20	The Bronze Age in Poland – archaeological and environmental traces of subsistence strategies Aldona Mueller-Bieniek (Polish Academy of Sciences, Warsaw) et al.	
10:20 – 10:40	Timing the introduction of <i>Panicum miliaceum</i> to the Middle Danube Region – dual evidence Mária Hajnalová (Constantine the Philosopher University, Nitra) & Peter Barta (Comenius University, Bratislava)	
10:40 – 11:10	COFFEE BREAK	
11:10 – 11:30	Bronze Age agricultural changes in the Czech Republic Adéla Pokorná, Petr Kočár & Tereza Šálková (Czech Academy of Sciences, Prague)	
11:30 – 11:50	Millets in Bulgaria – diachronic overview of their role in the subsistence and critical review of the archaeological finds Elena Marinova (State Office for Cultural Heritage Baden-Württemberg, Hemmenhofen), Ivanka Hristova (Umeå University) & Delphine Fremontdeau (University of Leuven)	
11:50 – 12:10	Archaeobotany in Romania – investigations in the Late Bronze Age fortification Cornesti-Jarculi Astrid Stobbe, Christoph Herbig & Maren Gummior (Johann Wolfgang Goethe-Universität, Frankfurt am Main)	
12:10 – 12:30	Before and after millet in Slovenia, south of the Alps to the Balkans Tjaša Tolar (ZRC SAZU, Ljubljana)	
12:30 – 14:00	LUNCH BREAK The successful spread of millets in Northern Italy Mauro Rottoli & Barbara Proserpio (Musei Civici, Como)	
14:00 – 14:20	On the „early“ evidence of broomcorn millet in central Germany, primarily Saxony-Anhalt Monika Hellmund (Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Halle)	
14:20 – 14:40	Millets in Bronze Age agriculture and food consumption in Northeastern France Françoise Toulemonde (AASPE National Museum for Natural History/CNRS Paris) & Julian Wiethold (INRAP, Metz) et al.	
14:40 – 15:00	The start of millet cultivation in Iberia Hans-Peter Stika (University of Hohenheim, Stuttgart)	
15:00 – 15:20	Weedy millets and millet weeds Helmut Kroll (Kiel)	
15:20 – 15:40	COFFEE BREAK	
15:40 – 16:00	Ukrainian dishes from millet Galyna Pashkevych (NASU, Kiev)	
16:00 – 16:20	Novelties in animal herding and consumption in Bronze Age Europe László Bartosiewicz (Stockholm University)	
16:20 – 16:50	DISCUSSION KEYNOTE LECTURE Change in the later second millennium BC: plants, pots, and people Anthony Harding (University of Exeter)	
16:50 – 17:20	WORKSHOP DINNER	
17:30 – 19:00	THURSDAY, 28th NOVEMBER 2019 Arrival of participants to the venue	
08:30 – 09:00	SESSION 2: Beyond seeds: Isotopic and biomolecular evidence of millet cultivation and consumption Session chairs: Lorenz Schwark & John Meadows (CAU Kiel)	
09:00 – 09:20	Miliacin in palaeosols and sediments, a powerful biomarker for telling stories about broomcorn millet Blandine Courel (Université de Strasbourg & The British Museum), Pierre Adam (Université de Strasbourg), Philippe Schaeffer (Université de Strasbourg), Jérémy Jacob (Université d'Orléan & LSCE Gif-sur-Yvette)	
09:20 – 09:40	Molecular and isotopic identification of millet in prehistoric pottery. New results from Bruszczewo, Poland Edward Standal (The British Museum & University of York), Carl Heron (The British Museum), Oliver Craig (University of York), Jutta Kneisel, Wiebke Kirleis & Johannes Müller (CAU Kiel), Janusz Czebreszuk (Adam Mickiewicz University, Poznań)	
09:40 – 10:00	Towards the quantification of millet in ancient diets using stable isotopes Oliver Craig (University of York)	
10:00 – 10:20	Detecting the manuring of millet in the past Amy Syring (University of Oxford)	
10:20 – 10:40	Early integration of pastoralism and millet cultivation in Bronze Age Eurasia Taylor Hermes (CAU Kiel)	
10:40 – 11:10	COFFEE BREAK	
11:10 – 11:30	Chicken feed: tracking the introduction and incorporation of new plants and animals Naomi Sykes (University of Exeter) & Holly Miller (University of Nottingham)	
11:30 – 11:50	SESSION 3: Growing and eating millet: Insights from ethnography, agronomy and experimentation Session chair: Anna Maria Mercuri (University of Modena and Reggio Emilia)	
11:50 – 12:10	Traditional millet growing in NW Iberia: from the ethnographic insights to the archaeobotanical implications Andrés Teira-Brion (University of Santiago de Compostela)	
12:10 – 12:30	Looking Away from Europe: A Wider Perspective on Millets in Dry Lands Marco Madella & Carla Lancelotti (Universitat Pompeu Fabra, Barcelona)	
12:30 – 13:00	Proso millet: cultivation, agronomic practices, and uses Stefania Grando (International Crops Research Institute for the Semi-Arid Tropics)	
13:00 – 14:00	DISCUSSION & CLOSING REMARKS LUNCH	

Figure 4. Programme of the millet-themed workshop in Kiel, November 2019.

This volume

Some of the knowledge on millet presented at the workshop is compiled in these proceedings. One of the aims of this volume is to illuminate the wider context of the spread and acceptance of millet in parts of Europe: agricultural, social, technological, environmental, architectural, culinary, ritual and symbolic. The papers do so by re-iterating the importance of looking beyond the list of crops when studying the history of food and food production. The broader-based approach advocated here extends to the methodologies of millet-related research, whereby sophisticated techniques and non-conventional proxies are crucial, but the information they offer must be checked against the results produced by other approaches, as a two-way control. To this point, we note that radiocarbon dating of millet grains has clearly demonstrated the importance of applying direct and adequate methods, which go as far as dismantling old narratives, but that the dates alone cannot tell the story of the agricultural change, its causes and ramifications.

The papers here ‘put flesh on the bones’. They examine the cultural and environmental circumstances surrounding the start of millet cultivation in Europe. They present novel methods to investigate the production and use of millet on the past. They showcase rare examples of recent and modern-day growing of millets in Europe.

The book is organised in two sections: the first, ‘Innovations in the Bronze Age subsistence economy: regional perspectives from Ukraine to France’, includes nine contributions that describe the social, technological, economic, dietary and natural environment in different regions in Europe at the time of the establishment of millet cultivation. The second section, ‘Approaching millet cultivation and consumption through high-end microscopy, chemistry and ethnography’, presents some new, non-carpological methods of tracing the cultivation and consumption of millets in the past – using high-resolution microscopic analysis of millet grain impressions and casts, as well as biomolecular and isotopic studies. It also emphasises the ethnographic route to understanding millet agriculture, where the living agronomic knowledge illuminates past practices. Below we provide sneak-peeks into the contributions.

Innovations in the Bronze Age subsistence economy: Regional perspectives from Ukraine to France

The paper by **Galina Pashkevich** critically approaches the research history on finds of broomcorn millet in the Ukraine, the region with the earliest dating of millet on its westwards shift to Europe in the 17th century BCE. Prominent finds of millet from the main periods of prehistory and history are listed here, some previously mentioned only in the literature written in Russian or Ukrainian. The paper further describes ancient millet cuisine based on written sources and discusses the sociocultural relevance of millet dishes in Ukraine. It is accompanied by two recipes for millet dishes that allow the reader to improve their own cooking practices and widen their culinary horizon.

Sofia Filatova explores the environmental, cultural and archaeobotanical context of the introduction of broomcorn millet in Bronze Age Hungary. She summarises the settlement history of the 3rd and 2nd millennium BCE, changes in which seem to reflect the apparently fluctuating population density and possible diachronic shifts in food economy between more and less labour-intensive productions. The author then conducts an evaluation of the existing quantitative archaeobotanical data for Bronze Age sites in Hungary, acknowledging that the field and analytical methodologies have varied among sites and among specialists. Using standardised values for the numerous crops recorded at Bronze Age sites, the author shows how the range and quantities of crops varied among cultures and changed over time. As

is clear from this overview, broomcorn millet arrived in the Carpathian Basin at a time of cultural transformation – from Middle to Late Bronze Age and the spread of the Tumulus culture – and a time of changing climatic conditions and vegetation cover, from more to less humid and from less to more open, respectively.

Laszlo Bartosiewicz describes the dynamics of animal herding and use during the Bronze Age, primarily in the Carpathian Basin, but also in neighbouring regions, especially those to the east of the Basin, pointing out regional variations in the uses of domesticates. He offers a crisp summary of the general chronological and environmental background to the Bronze Age, noting in particular the predominance in central Europe of a cooler and increasingly humid climate by the middle of the Subboreal, the period encompassing the Bronze Age. For the Carpathian Basin, he gives an overview of the representation of the major animal species in the Bronze Age, emphasising the high variability in this within and between settlement types and the general importance of cattle as a source of both meat and secondary products. The author also documents the sharply rising rate of use/importance through time of caprines vs. pigs in the region and reviews the evidence suggesting the production of wool as a factor driving this trend. The contribution benefits from the bonus commentaries on the (early) finds of chicken, specialised antler working in the Bronze Age, and the curious finds of perforated hare and dog metapodia at a few Bronze Age sites in Hungary. A prominent place is given to the discussion of the appearance and role of horse, and how the animal's mobility enabled a greater degree of movement on the part of humans. The author illustrates the 'special status' of horses with examples of horse burials in the West Asian steppe.

Kelly Reed, Jacqueline Balen, Ivan Drnić, Sara Essert, Hrvoje Kalafatić, Marija Mihaljević and Emily Zavodny provide an overview of the archaeobotanical records of millet and the stable isotopic evidence of millet consumption by humans and/or animals in Croatia. They offer more detail about the archaeological sites and contexts yielding the grains of broomcorn and foxtail millet that were dated within the 'Millet Dating Programme'. They observe that the data on the Bronze Age plant economy and diet, and changes in these, in Croatia are still very sparse but that the available evidence clearly points to the Middle to Late Bronze Age as the period when crop cultivation began to rely on broomcorn millet and, later on, a few other crops not known in previous periods. As one possible reason for the diversification of crop growing, they suggest risk buffering, which is particularly relevant in agriculturally marginal environments, where a wider range of crops ensured a regular food supply.

The paper by **Tjaša Tolar** and **Primož Pavlin** builds a picture of prehistoric crop cultivation and wild plant gathering in Slovenia, from the thus far earliest crop traces in the region, registered at Eneolithic sites, through to the Early Iron Age. Prior to presenting the evidence of plant production and consumption, they introduce us to the region and its later prehistory, emphasising key developments in technologies of material production (e.g. copper and bronze metallurgy). They also summarise the characteristics of burial ritual, settlements and houses, emphasising the fortified nature of the habitations in the Late Bronze Age and Early Iron Age, including the sites from which the dated broomcorn millet grains derived. This comprehensive contribution also includes an overview of finds of harvesting tools, which takes us back in time to the early days of wild grass cutting in the Natufian culture of prehistoric southwest Asia. In Slovenia, flint pieces used as implements in sickles are known from the Late Neolithic, whereas the Bronze Age was the time of metal sickles, commonly found in hoards of metal objects, suggesting that they were highly valued.

In her paper, **Monika Hellmund** revisits a number of reportedly Neolithic broomcorn millet finds in central Germany and, using the radiocarbon dates, in many cases obtained directly on the millet grains, demonstrates how all of them are later intrusions, many from the Late Bronze and Early Iron Age. Along the way,

she takes us through the exceptionally rich archive of prehistoric millet finds in the German state of Saxony-Anhalt, including many cases of lumps of fused grains (millet ‘porridge’) or otherwise large deposits. She also looks beyond the borders of Saxony-Anhalt and discusses some of the ‘early’ millet finds in the neighbouring German states, emphasising again the insecure temporal attribution of these finds in the absence of direct radiocarbon dates. This critical and detailed overview emphasises the importance of accurate chronological determination of archaeobotanical finds, since broader narratives on crop introductions and agricultural innovations directly depend on accurately establishing their timing. In addition to scrutinising the finds of broomcorn millet in the study region and adjacent areas, Hellmund puts another late addition to the crop spectrum in focus – faba bean (*Vicia faba*). She observes the coinciding temporal and contextual occurrence of the two crops in the Late Bronze Age and the Early Iron Age in central Germany and argues for their importance in both the mundane and the ritual spheres.

Françoise Toulemonde, Julian Wiethold, Emmanuelle Bonnaire, Geneviève Daoulas, Marie Derreumaux, Frédérique Durand, Bénédicte Pradat, Oriane Rousselet, Caroline Schaal and Véronique Zech-Matterne provide an in-depth overview of the spread of early millet in northeastern France in the Late Bronze Age, from the 14th century BCE onwards, that is based on an impressive compilation of new and already published archaeobotanical data and new radiocarbon dates. It is interesting to learn that the establishment of millet cultivation in this region coincides with an increase in human population size and with human settlements expanding into previously unexploited environments. The widening of the crop spectrum in northeastern France in the Bronze Age extends beyond the integration of millet and includes the newcomers faba bean and gold-of-pleasure (*Camelina sativa*). Further, new farming strategies emerge involving a dynamic agropastoral economy. In addition to the socio-economic relevance of millet, the authors stress the sociocultural dimensions of millet consumption in the Bronze Age, since it occurs in collective feasting sites as well as in funerary contexts.

In an ambitious integration of off-site and on-site palaeo-environmental, archaeobotanical and archaeological evidence, **Ingo Feeser, Stefanie Schaefer-Di Maida, Stefan Dreibrodt, Jutta Kneisel and Dragana Filipović** chart the process of cultural and environmental development before and during the last centuries of the 2nd millennium BCE in northern Germany. Their focus is on a long-lived burial site in Schleswig-Holstein, Mang de Barga, where they found evidence of changing landscape and intensifying agricultural and ritual use of the land, all culminating around the 12th century BCE, which is when broomcorn millet was introduced in north-central Europe and when a new cultural phenomenon, the Urnfield culture, spread across the region. The authors conclude that the shifts they see in the local vegetation composition resulted from abrupt, concurrent and considerable changes in local funerary, settlement and agricultural activities during the 13th to 11th centuries BCE. They then look at the evidence from the wider region around the site and beyond it, from Schleswig-Holstein; they find that the environmental and cultural transformation took place at all three scales observed – from local to regional to supra-regional.

Approaching millet cultivation and consumption through high-end microscopy, chemistry and ethnography

The paper by **Edward Standall, Oliver Craig and Carl Heron** shifts our focus from the growing of millet to its processing for food or drink. The authors describe novel methods – biomarker and carbon and nitrogen stable isotope analyses – applied in the identification of millet in organic residues. As an illustration, they summarise examples of the studies that detected the diagnostic biomolecule miliacin in ceramic

vessels: in charred surface deposits (so-called foodcrusts) and/or absorbed into the ceramic fabric. This molecule is retained after the processing of millet or millet-containing foods, in the pots. Besides highlighting the advantages, the authors, importantly, outline the limitations of these new approaches and provide guidance for their future development and application.

In the contribution by **Marta Dal Corso, Marco Zanon, Carl Heron, Mauro Rottoli, Michele Cupitò, Elisa Dalla Longa** and **Wiebke Kirleis** broomcorn millet has been attested by the biomarker miliacin in the archaeological sediments at the Terramare culture site of Fondo Paviani (ca. 1350-1100 BCE), in the Po River valley, northern Italy. This molecule is typical of *Panicum miliaceum* and it was absent in the other modern panicoid grasses analysed for reference. The study presents the results of investigation of on-site sediments and, together with the available archaeobotanical and isotopic records, inserts them into the picture of millet consumption and cultivation during the Late Bronze Age in the Po Valley.

Eiko Endo describes in detail the method she applies in the analysis of plant impressions within the fabric of pottery, which has proved highly efficient and reliable in her work in Japan, but also in Ukraine. In her paper, she gives details on the analytical procedure, which allows her to identify the impressions down to species as well as to plant part (e.g. grain vs. chaff). As an illustration of the application of this method, she summarises the study she conducted on a pottery assemblage from a large number of prehistoric and historic sites in the Japanese archipelago. She also lists the types of questions one can ask based on the record of impressions, such as those related to formation processes and the use of crop-processing by-products. The paper highlights the potential benefit of this method for the study of crop cultivation and use in general, and of millets in particular. Particularly promising is the application at sites or in regions and periods where plant impressions are the major or the only recovered source of information on plant use.

Andrés Teira-Brión's ethnographic study presents the living tradition of broomcorn and foxtail millet cultivation in northwestern Spain and northern Portugal, both regions of high rainfall. Earlier ethnographic work in this area, in which the author participated, documented millet-growing, -processing and -preparation procedures. In this paper, the author emphasises variations in these practices between neighbouring regions. For instance, in the relatively low-lying Galicia and Minho regions, the farmers' focus is on broomcorn millet, which they use as both food and fodder. In mountainous Asturias, foxtail millet is grown as green fodder, whereas the occasional broomcorn millet in foxtail millet fields is a tolerated weed. But the author does not just describe these general observations. He looks at a much broader socio-economic context – the place of millet cultivation in agricultural systems, which, as he emphasises, are built around, and shaped by, much more than a presence/absence of a crop. Land ownership and inheritance rules, the organisation of production, the structure of community, the balance between arable and pastoral farming or a focus on one of the two, production for own needs and/or market, flexibility of the agrarian cycle – all these and other aspects play a role in agricultural decision making, change, adaptation and adoption. The author urges us to consider the broader economic and social contexts of agriculture in general and millet growing in particular before coming to conclusions about millet as a highly desired 'enabler crop'. This contribution reminds us of the possibilities and limitations of the use of ethnological analogies to inform archaeological interpretations and the danger of directly superimposing present-day perceptions and knowledge onto the past.

Wiebke Kirleis and **Marta Dal Corso** present an interesting example of highly specialised handicraft – broom production – reliant on millet cultivation, in this case of the species *Sorghum technicum*. Their ethnographic observations emphasise

how rare this handicraft is in Europe, as it is nowadays practised in Moldova and northern Serbia, but hardly anywhere else. The richly illustrated paper presents the *chaîne opératoire* of the almost-extinct practice, covering the production steps from harvesting the plants to tying the brooms. This contribution shifts the attention to the non-food purposes of millet cultivation, highlighting the place of millet and millet-related activities in the production of objects for everyday use and the craft associated with the use of millet beyond solely foodstuff.

Acknowledgements

The work on broomcorn millet conducted within the CRC 1266 is one more ‘panicle’ in the global field of millet research. Besides the three of us, many collaborators and colleagues have helped this panicle to grow and the grains to ripen. Some of our associates were directly involved, throughout the research process or at some of its stages, and others contributed with their insights, opinions or practical help. A heartfelt thank-you to all, those named below and those who remain unnamed.

A core group at Kiel included our colleagues within the CRC 1266, above all our key collaborator, John Meadows. Another group, variously linked with the CRC 1266, working both at Kiel and elsewhere, included all of our co-authors on the millet papers that have been published or are forthcoming. Many of them joined us at the workshop in Kiel, and we are delighted that we could host them. We are also lucky that a number of other colleagues accepted our invitation to the workshop and made the event all the more diversified; we are grateful for their interest and enthusiasm.

Both the Millet Dating Programme and the workshop relied immensely on the organisational and technical support of the staff and students at the Institute for Prehistoric and Protohistoric Archaeology at Kiel University, which hosts the CRC 1266. Key help was provided by Tanja Reiser, Yasmin Dannath, Anna Carina Lange, Ralf Opitz, Carsten Reckweg and Janina Stampe. Tomasz Goslar, at the Poznan Radiocarbon Laboratory, facilitated radiocarbon dating of most of the samples included in the programme and provided all the necessary information and details. Christian Hamann, at Kiel, oversaw millet dating done at the Leibniz-Laboratory and provided some crucial advice.

The preparation of the proceedings was possible thanks to the eagerness, time and effort of the authors of the papers included here. We are thrilled and grateful that they responded to our call for contributions. A group of colleagues gave advice or acted as reviewers for this book, and we are very much indebted to them for their readiness to join our team.

Funding for the Millet Dating Programme, the workshop, and this book was secured through the German Research Foundation (DFG – project number 2901391021 – SFB 1266).

References

- Alsleben, A., 2012. The plant economy of northern medieval Russia. In: M. Brisbane, N. Makarov, E. Nosov, eds., *The Archaeology of Medieval Novgorod in Context: Studies in Centre/Periphery Relations*. Oxford: Oxbow, 321-350.
- Anderson, R.L. and Greb, B.W., 1987. Residual herbicides for weed control in proso millet (*Panicum miliaceum* L.). *Crop Protection*, 6 (1), 61-63. [https://doi.org/10.1016/0261-2194\(87\)90030-5](https://doi.org/10.1016/0261-2194(87)90030-5)
- Becker-Dillingen, J., 1927. *Handbuch des Getreidebaues einschließlich Mais, Hirse und Buchweizen*. Berlin: Verlag Paul Parey.
- Brandes, D., 1995. Breiten sich die C4-Pflanzen in Mitteleuropa aus? In: Bundesanstalt für Vegetationskunde, Naturschutz und Landschaftspflege in Bonn-Bad Godesberg, Bundesforschungsanstalt für Naturschutz und Landschaftsökologie, Deutschland Bundesamt für Naturschutz, eds., *Dynamik und Konstanz: Festschrift für Herbert Sukopp*. Schriftenreihe für Vegetationskunde 27. Münster: Landwirtschaftsverlag, 365-372.
- Brown, R.H., 1998. Agronomic implications of C4 photosynthesis. In: R.F. Sage and R.K. Monson, eds., *C4 Plant Biology*. San Diego: Academic Press, 473-507.
- Chen, F.H., Dong, G.H., Zhang, D.J., Liu, X.Y., Jia, X., An, C.B., Ma, M.M., Xie, Y.W., Barton, L., Ren, X.Y., Zhao, Z.J., Wu, X.H., Jones, M.K., 2015. Agriculture facilitated permanent human occupation of the Tibetan Plateau after 3600 BP. *Science*, 347 (6219), 248-250. <https://doi.org/10.1126/science.1259172>
- Dal Corso, M., Pashkevych, G., Filipović, D., Liu, X., Motuzaite-Matuzeviciute, G., Stobbe, A., Shatilo, L., Videiko, M., Kirleis W., 2022. Between cereal agriculture and animal husbandry – Millet in the early economy of the North Pontic region. *Journal of World Prehistory*.
- d'Alpoim Guedes, J., 2018. Did foragers adopt farming? A perspective from the margins of the Tibetan Plateau. *Quaternary International*, 489, 91-100. <https://doi.org/10.1016/j.quaint.2016.12.010>
- Das, S., Khound, R., Santra, M., Santra D.K., 2019. Beyond bird feed: proso millet for human health and environment. *Agriculture*, 9 (3): 64. <https://doi.org/10.3390/agriculture9030064>
- Erler, M., Keck, M., Dittrich, C., 2020. The changing meaning of millets: organic shops and distinctive consumption practices in Bengaluru, India. *Journal of Consumer Culture*. <https://doi.org/10.1177%2F1469540520902508>
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Kloof, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A.M., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H.-P., Stobbe, A., Tolar, T., Wasylkowa, K., Wiethold, J., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://www.nature.com/articles/s41598-020-70495-z>
- Geisler, G., 1988. *Pflanzenbau: Ein Lehrbuch: Biologische Grundlagen und Technik der Pflanzenproduktion*. 2nd ed. Berlin: Verlag Paul Parey.
- Hardenberg, R., 2021. The rise of a new millet assemblage in India. In: T. Schade, B. Schweizer, S. Teuber, R. Da Vela, W. Frauen, M. Karami, D. Kumar Ojha, K. Schmidt, R. Sieler, M. Toplak, eds., *Exploring Resources: on Cultural, Spatial and Temporal Dimensions of Resource Cultures*. RessourcenKulturen 13. Tübingen: Tübingen University Press, 141-156.
- Hastorf, C.A., 2017. *The Social Archaeology of Food*. Cambridge: Cambridge University Press.

- Hermes, T.R., Frachetti, M.D., Doumani Dupuy, P.N., Mar'yashev, A., Nebel, A., Makarewicz, C.A., 2019. Early integration of pastoralism and millet cultivation in Bronze Age Eurasia. *Proceedings of the Royal Society B*, 286, 20191273. <https://doi.org/10.1098/rspb.2019.1273>
- Hunt, H.V., Van der Linden, M., Liu, X., Motuzaite-Matuzeviciute, G., Colledge, S., Jones, M.K., 2008. Millet across Eurasia: chronology and context of early records of the genera *Panicum* and *Setaria* from archaeological sites in the Old World. *Vegetation History and Archaeobotany*, 17, S5-S18. <https://doi.org/10.1007/s00334-008-0187-1>
- Lieberei, R. and Reisdorff, C., 2007. *Nutzpflanzenkunde*. 7th ed. Stuttgart: Thieme.
- Miedaner, T. and Longin, F., 2012. *Unterschätzte Getreidearten: Einkorn, Emmer, Dinkel & Co*. Clenze: Agrimedia-Verlag.
- Miller, N.F., Spengler, R.N., Frachetti, M., 2016. Millet cultivation across Eurasia: origins, spread, and the influence of seasonal climate. *The Holocene*, 26, 1566-1575. <https://doi.org/10.1177%2F0959683616641742>
- Moreno-Larrazabal, A., Teira-Brión, A., Sopelana-Salcedo, I., Arranz-Otaegui, A., Zapata, L., 2015. Ethnobotany of millet cultivation in the north of the Iberian Peninsula. *Vegetation History and Archaeobotany*, 24 (4), 541-554. <https://doi.org/10.1007/s00334-015-0518-y>
- Müller, J. and Kirleis, W., 2019. The concept of socio-environmental transformations in prehistoric and archaic societies in the Holocene: an introduction to the special issue. *The Holocene*, 29 (10), 1517-1530. <https://doi.org/10.1177%2F0959683619857236>
- Pyankov, V.I., Ziegler, H., Akhane, H., Deigle, C., Luetge, U., 2010. European plants with C4 photosynthesis: geographical and taxonomic distribution and relations to climate parameters. *Botanical Journal of the Linnean Society*, 163 (3), 283-304.
- Saha, D., Channabyre Gowda, M.V., Arya, L., Verma, M., Bansal, K.C., 2016. Genetic and genomic resources of small millets. *Critical Reviews in Plant Sciences*, 35, 56-79. <https://doi.org/10.1080/07352689.2016.1147907>
- Sitte, P., Ziegler, H., Ehrendorfer, F., Bresinsky, A., 1991. *Strasburger Lehrbuch der Botanik für Hochschulen* (33rd edition). Stuttgart: Gustav Fischer Verlag.
- Taylor, J. and Duodu, K.G., eds., 2018. *Sorghum and Millets: Chemistry, Technology, and Nutritional Attributes*. Cambridge: Elsevier.
- Wiegmann, G., 1967. *Alltags- und Festspeisen: Wandel und gegenwärtige Stellung*. Münster: Elwert Verlag.

**SECTION 1: INNOVATIONS IN THE
BRONZE AGE SUBSISTENCE ECONOMY:
REGIONAL PERSPECTIVES FROM UKRAINE
TO FRANCE**

The (pre)history of common millet (*Panicum miliaceum*) in Ukraine and its place in the traditional cuisine

Galyna Pashkevych

Abstract

The generally high scientific interest in the Tripolye prehistoric culture of Ukraine peaked several years ago. Conferences were held and many relevant publications appeared at the time. There were journalists at these conferences and, wishing to entertain the conference participants, they prepared a *kulish* meal. In their opinion, *kulish* was a typical Tripolye dish. However, the archaeobotanical evidence shows that this is not the case, because the main ingredient in this traditional dish is millet (*Panicum miliaceum*) and the association of millet with the Tripolye period is wrong. Millet appeared in Ukraine much later than this culture. The history of the emergence of millet on the territory of Ukraine has been unravelled thanks to archaeobotanical and other research, the results of which are summarised here.

Introduction

The wide array of cultivated plants and the traditional methods of their cultivation, processing and dietary use are a part of human culture and reflect the talent, intellect and diligence of many generations. There has, therefore, been growing awareness of the need to study and preserve the heritage of culinary knowledge and practices. This has recently started to manifest itself in the increased attention being paid to ancient crop varieties and other plants as sources of products for healthy nutrition and wellbeing.

Archaeobotanical materials are an important source of information on the history of the origin, distribution and use of plants. One of the plants that has attracted the interest of both archaeobotanists and laypersons is common/broomcorn/proso millet (*Panicum miliaceum*). The cultivation and consumption of this cereal has a long tradition in Ukraine. This paper revisits the ‘early’ finds of millet in Ukraine in light of the recent determinations of the precise age of these finds. It then moves through time and presents some of the records of millet from the classical antiquity and medieval periods, after which it turns to the recent and modern-day role of millet in the region.

The ‘early’ finds of common millet in Ukraine

The Tripolye culture extended over great parts of the modern forest-steppe zone in Ukraine, in the area between the Prut and Dnieper rivers. Calibrated ^{14}C dates indicate that the Tripolye culture covered the period from the 5th to the 3rd millennium BCE (e.g. Müller *et al.* 2016). The major cereals grown by these groups were the hulled wheats einkorn (*Triticum monococcum*) and emmer (*T. dicoccum*); barley (*Hordeum vulgare*) was a minor crop. The list of cultivated plants from this period further includes pea (*Pisum sativum*), bitter vetch (*Vicia ervilia*), lentil (*Lens culinaris*) and flax (*Linum usitatissimum*) (Янушевич 1976, 1980; Пашкевич 1980, 1989, 1990a, 1993, 2000a, 2011; Pashkevich/Pashkevych 2003, 2005, 2012; 2016; Пашкевич and Відейко 2006; Kirleis and Dal Corso 2016).

For the territory of present-day Ukraine and Moldova, the presence of millet on Neolithic and Eneolithic sites was previously reported as involving rare, single impressions of grain in pottery or daub (Янушевич 1976; Pashkevych 2012). Yanushevich notes that some impressions said to be millet grains in pottery and daub from the settlements of the Bugo-Dniester culture and the early phase of the Tripolye culture are not very clear and that some of them probably are of the weedy species *Setaria glauca*, *Setaria viridis* or *Echinochloa crus-galli* (Янушевич 1976, 153). Impressions of what seems to be millet grain were also detected in the clay materials of the middle and late phases of the Tripolye culture, where it was reported that they preserved morphological features of the grains (Pashkevych 2012).

Neolithic and Eneolithic records of millet are known not only from Ukraine, but also from central and western Europe. Detailed lists of these ‘oldest’ finds of millet grains in Europe are given by Zohary and Hopf (2000, 83-85) and Hunt *et al.* (2008), the latter already critically asking when millet actually arrived in Europe. This issue was pursued by Motuzaite-Matuzeviciute *et al.* (2013) who, through radiocarbon dating some of the ‘Neolithic’ millet grains from Europe, demonstrated that they date much later. This raised suspicion about the age of other such ‘early’ millet finds on the continent and emphasised the importance of obtaining reliable dates in order to answer the question of when millet arrived. With the completion of a much larger scale Millet Dating Programme at Kiel University, it is now clear that millet became a staple crop in Europe only in the 2nd millennium BCE and that the millet finds from earlier archaeological contexts represent later intrusions (Filipović *et al.* 2020).

Before the start of archaeobotanical analysis in Ukraine, the determination of archaeological plant remains was based on the knowledge of archaeologists, which was insufficient, and on some general assumptions. For example, *Setaria* was referred to as ‘millet’, but without specifying whether it was foxtail (*italica*) or some other *Setaria* species. For instance, at the Tripolye settlement site of Grebeni (excavated from 1961-1963 by S.N. Bibikov) yielded a vessel filled with grains, which was stored in the site’s archive under the label ‘millet’. Analysis under a microscope revealed that these were instead grains of the weedy species *Setaria glauca* (Пашкевич 1991). One plant of this species produces about 6000 grains and, when they are ripe, shatters them rapidly and infests the soil. *S. glauca* seeds do not lag much behind common millet in terms of their nutritional value and can be used as food (Рева М.І. and Рева Н.Н. 1976). The grains found in the vessel may have been collected and kept in the pot for some kind of use.

In the 20th century, archaeobotanists rarely used scanning electron microscopy, and the radiocarbon dating technique was much less precise than nowadays. Results obtained with conventional microscopy and non-AMS radiocarbon dating can, therefore, benefit from a revision. An excellent example is the recent re-examination of the plant impressions on figurines of the Usatovo culture of prehistoric Ukraine. This study was carried out as part of the research on the timing and pace of the movement of millets from China to Europe. Archaeobotanical studies of Usatovo

culture materials were carried out by N. Kuzminova (Кузьминова and Петренко 1989), who identified impressions of millet grains on 70 anthropomorphic clay figurines. Re-analysis of these impressions was recently conducted using a scanning electron microscope, and this re-analysis showed that none of these impressions bear morphological features of common millet (An *et al.* 2018).

Another example is the radiocarbon dating of charred grains of common millet from the territory of Ukraine, performed jointly with archaeobotanists from Kiel University (Filipović *et al.* 2020; Dal Corso *et al.* 2022). The results revealed that common millet first appeared in Ukraine in the last quarter of the 17th century BCE; the dated grains from the Bronze Age settlement at the site of Vinogradniy Sad, in southeastern Ukraine are, for now, the earliest finds of common millet in Europe. Perhaps Ukraine was the gateway through which millet entered Europe on its westward journey from the areas of origin in China.

Finds of common millet from the Bronze Age to early modern times in Ukraine

For most of its duration, the Bronze Age in Europe (ca. 3000-500 BCE) coincided with the Subboreal climatic period. Under the impact of climatic deterioration, with frequent droughts and cold spells, significant changes took place in the economy, social organisation and material culture of the populations of central and eastern Europe. For instance, the Tripolye culture disappeared at the time of a dry Subboreal phase.

According to the radiocarbon AMS-dates, millet grains from Vinogradniy Sad, a settlement of the Bronze Age Sabatinovka culture, date from the period 1631-1455 BCE (Filipović *et al.* 2020; Dal Corso *et al.* 2022). This settlement was excavated at the end of the 1980s; soil samples were taken from the cultural layers and wet sieved, and charred plant remains were extracted and archived. The crop assemblage from Vinogradniy Sad is composed mainly of hulled wheat, naked barley and common millet (Пашкевич and Костилюв 1992). The data from this and other sites of the Sabatinovka culture show that barley, common millet and emmer were the principal crops of the Bronze Age (Pashkevich 2003). The Bronze Age economy in the steppe and forest-steppe zone of Ukraine was based on stock breeding and agriculture. Archaeobotanical records show that agriculture gained in economic importance during the Middle Bronze Age and Late Bronze Age (Пашкевич and Шовкопляс 2012).

As shown by the radiocarbon dating, by the beginning of the 12th century BCE, common millet was being grown throughout Europe, including in its northern part. In Ukraine, millet grains from the Belozerska culture settlement of Dikiy Sad, located in the town of Nikolaev, gave the following radiocarbon dates: 1214-1001 BCE and 1209-979 BCE (Filipović *et al.* 2020; Dal Corso *et al.* 2022). The settlement itself dates to the end of the Bronze Age. Archaeobotanical analysis of the pit fills and layers within structures documented the presence of barley, common millet, free-threshing wheats and hulled wheats, as well as seeds of crop weeds (Горбенко and Пашкевич 2010).

From the Bronze Age onwards, common millet frequently occurs in large quantities throughout Ukraine, in both the steppe and the forest-steppe zones, and even in the forested areas. In particular, it seems to have had great importance in the Iron Age in the forest-steppe zone. In *The Histories*, Book IV, Ch. 18 and 75-76, Herodotus writes about the use of common millet and hemp by the Scythians (Godley 1920). Hulled wheats (principally emmer) and hulled barley, along with common millet, dominated the food production of the Scythians in the forest-steppe zone (Janushevich 1981; Янушевич 1986; Пашкевич 1991, 2001; Пашкевич and Горбаненко 2019). Charred grains of common millet were recovered at the sites of Ivane-Puste and Zalissyia, in

the western part of this zone (Пашкевич 1991). Crop cultivation became prominent in the predominantly pastoral food economy of the Scythians in the steppe zone in the period 500-400 BCE, when these groups occupied the lower Dnieper region. The crops they cultivated were barley and millet and, to a lesser extent, emmer (Pashkevich 1984; Гаврилюк and Пашкевич 1991; Пашкевич 2000b). The composition of plant assemblages, the types of tools and some ethnographic data offer a basis for the reconstruction of the nature of crop cultivation. It seems that, at this time, parts of the population shifted their economic base more to farming. The need for winter fodder for the herds led to a focus on certain crops, primarily common millet and barley that have traditionally been known as forage plants. Apparently, crop cultivation emerged as a key subsistence strategy of the Scythians in the steppe zone due to the need for animal fodder (Dal Corso *et al.* 2022).

There were Early Iron Age settlements in the northern Black Sea region that were located in similar environments, but their economies were different. Most obviously, the great importance of bread wheat is a distinctive feature of the ancient Greek towns in the North Pontic; besides bread wheat, hulled barley was also very common here. Free-threshing wheat arrived in the northern Black Sea region with the Greek colonists from Miletus, in Asia Minor, and, in the course of time, was integrated into agricultural production over the entire territory of modern Ukraine. It is evident that the Greek settlers brought new food production strategies and used the crops known from their homeland. Hulled wheats were of less significance, whereas a range of pulses were grown, including bitter vetch, pea, lentil and chickpea (Янушевич 1976, 1986; Janushevich and Nikolaenko 1979; Janushevich 1989; Пашкевич 1990b, 2002, 2004, 2005, 2006, 2016).

The significance of hulled wheats among the plants constituting food for the residents of the ancient Greek city of Chersonesos, on the Crimean Peninsula, may have been connected with an economic downturn. Grain reserves stored in individual homesteads were intended only for consumption and not for sale. The importance of crafts increased and the inhabitants of Chersonesos recovered the lost trade income by selling their handicrafts to the groups living in the hilly regions (Сорочан *et al.* 2001), where a different set of grain crops was cultivated to that in the *chora* of Chersonesos; the dominant ones where hulled wheats, rye and common millet.

The importance of millet in the Greek colonies of the northern Black Sea region increased in the first centuries CE, when contacts and mixing with the local population became reflected in the crop spectrum of the Greek communities, which now also included hulled wheats and barley. These crops, which had long been locally grown, were well adapted to the climatic and soil conditions of the region. The last centuries of the 1st millennium BCE and the first centuries CE were the time when millet was widespread in eastern and central Europe. Charred grains and their impressions from this period were found at sites in Slovakia, Germany and Poland.

Over a long period, millet was the main grain crop of the Slavic tribes in Ukraine, starting from the 3rd century BCE and continuing up to 7th century CE. At some of the sites, it makes up almost 60% of the plant assemblage (Пашкевич 1988a, b; Горбаненко and Пашкевич 2010; Пашкевич and Горбаненко 2010).

The Rus' people of early medieval eastern Europe grew bread wheat, rye and common millet. The associated archaeobotanical samples date from the 9th to the 14th centuries CE and were collected from such places as old Rus' towns, fortified villages and military camps; they vary in volume from a few grams to several kilograms (Пашкевич 2010; Козловська *et al.* 2012). The significant finds come from the period spanning the 11th to 13th centuries CE, from the territory of ancient Kyiv, where archaeological work has been going on for more than 100 years. During the excavations of 1907-1908, charred grains of rye, millet and free-threshing wheat were most commonly found in the Rus' structures. In some areas wheat prevailed, in others millet (Пашкевич 2010).

In the 18th and 19th centuries CE, Russia occupied first place in the world for the cultivation of common millet, as highlighted by the words of the famous botanists who travelled through the Russian Federation at the time. The well-known scientist Piotr Zhukovsky stated that millet arrived in Europe from East Asia together with nomadic tribes.

‘Of the cereals, millet is probably the most ancient food plant in China. All of the most ancient traces of Chinese history from around 2700 BCE testify to this. Even younger evidence, dating back to the Yin era (XVIII-XII BCE), speaks of the leading significance of millet’¹ (Жуковский 1964, 229-230).

According to N.I. Vavilov, common millet was the favourite cereal of the nomads in Eurasia, and it was with these tribes that he associated the appearance of millet in Europe.

‘Millet can be sown very late, at different periods, and does not bind the nomads. Sowing [...] requires very few grains; millet is extremely transportable, unusually drought resistant, it grows even on sandy soils [...] and therefore is still an indispensable attribute of the nomadic economy of the semi-desert regions of Asia and the southeast of the European part of the Russian Federation’² (Вавилов 1987, 65).

In the selected works by N.I. Vavilov, various forms of economy among the Afghan tribes are described: *‘Even nomads were engaged in agriculture, especially during long migrations from the south of the country to the north’* (Вавилов 1959, 105-106). The advantage of sowing millet is also emphasised:

‘A small amount of grains per hectare is needed for sowing and, at the same time, [it is] undemanding in terms of conditions. Therefore, millet is a good companion of nomadic and semi-nomadic tribes’³ (Вавилов 1959, 262-264).

Some insights on the role of millet in ancient and traditional cuisines

Millet is one of the most important grain crops. The grains removed from the glumes (groats) are called *psheno* in Ukrainian. It has a high protein content (12-14%) and a low percentage of fat (1-3.5%), e.g. lower than that of oat groats. Millet boils quickly and absorbs water easily (Вавилов *et al.* 1986, 143-152). Unlike buckwheat and oat groats, millet has little fibre; vitamin PP; and microelements such as copper, nickel, manganese and zinc. The proteins it contains are quickly oxidised leading the groats to become bitter. Other parts of plant, i.e. straw and chaff, are used as hay or green fodder.

The first written evidence for the use of common millet is found in the works of the ancient Roman author Columella:

*‘Bread is made from millet, which can be eaten with pleasure while it is still warm. From the mogar [or foxtail millet, *Setaria italica*], when crushed and freed from the brans, as well as from the millet, porridge is obtained in any quantity; it is quite tasty, especially with milk’⁴ (after Сергеев 1970, 14).*

1 Translated by the author.

2 Translated by the author.

3 Translated by the author.

4 Translated by the author.

Pliny the Elder reported in his *Natural History*, Book XVIII:

'Millet flourishes particularly well in Campania, where it is used for making a white porridge; it also makes extremely sweet bread. Moreover, the Sarmatian tribes live chiefly on millet porridge, and even on raw meal, mixed with mare's milk or with blood taken from the veins in a horse's leg. Millet and barley are the only grains known to the Ethiopians' (Rackham 1938, 253).

Common millet and foxtail millet (or *mogar*) are mentioned by other classical sources too. According to the ancient Greek authors, in the 1st century CE, millet and mogar were the main crops. Strabo named millet as one of the main crops in the typical Campanian crop rotation. At the same time, he pointed to its reliability: *'... the most reliable protection against hunger, because millet is stable in any weather and there is never a shortage of it'*⁵ (cited in Кузищин 1966, 111-113). Herodotus (485-425 BCE), in his Book IV, Ch. 17, mentioned consumption of lentil and millet among the Scythians who inhabited the Dnieper and Bug basins (Godley 1920). Xenophon (434-359 BCE), in *Anabasis*, Book 1, Ch. 22, describes how, in a campaign through Cilicia, he encountered fields sown with barley, wheat and millet (Brownson *et al.* 1922).

From an agronomic point of view, this cultivar is more thermophilic than other cereals and survives higher temperatures than do winter rye, wheat and oat. Unlike other cereals, millet is almost indifferent to the sowing time. It gives an equally good yield whether sown in early or late spring, or in the middle of summer. This has an advantage because, in the event of bad winter or spring weather affecting the wheat harvest, millet comes to the rescue. According to Columella's calculations for millet, on average, the number of grains needed for sowing is 20 times lower than for wheat and 40 times lower than for emmer (after Сергеев 1970, 149).

Today, millet is grown mostly in the Middle East, central Asia and India. In the past few decades, millet production has also been started in some American and European countries. In Ukraine, in contrast, the cultivation of millet has been drastically reduced in recent years, despite millet being a well-known and widespread crop here until the 21st century. Various dishes were made from it – porridge, soup and even a drink called *buza*, which contains 4-6% alcohol.

In the recent history of Ukraine, a range of millet-based dishes are found in the cuisine of the Zaporozhye Cossacks. This was a military organisation in the steppe zone of Ukraine, formed in the 16th century CE from the local Slavonic tribes and nomads. Their life was on the road, and so they prepared food on an open fire, in big copper or iron pots. Porridges were widespread, prepared from millet or buckwheat. The millet porridge had different names and ingredients: *teterya* was a combination of millet and sour rye dough, fat, milk or fish; *bratko* was a simple porridge made purely of millet. A more liquid millet dish was called *kandiour*. The Ukrainians also eat millet porridge with pieces of pumpkin. Sometimes they wrap millet porridge in cabbage leaves; cover the rolls with sour cream, spices and tomato sauce; and then boil for several minutes. This last dish is called *golubcy*.

Kulish is the most famous millet-based traditional Ukrainian dish, still popular today. Interestingly, the word 'kulish' is of Hungarian origin (*köles*=millet). Kulish is normally eaten for lunch in the field or for dinner at home. It is prepared on an open fire. Fat (*salo*), millet, onions, carrot, salt and pepper are the basic ingredients (Fig. 1). Potato was added in the 20th century CE. Kulish with meat is especially tasty, and the dish is also sometimes cooked with fish or mushrooms (Хименко 1960, 110; Шалімов and Шадюра 1981, 39). In ancient times, when the ingredients were deficient, kulish was also made with underground plant parts (bulbs or tubers, *e.g.* of *Typha*), which are rich in starch.

5 Translated by the author.




Figure 1. Two common recipes for Ukrainian kulish (suggested by Galyna Pashkevych; figure by Anna Carina Lange).

Conclusions

For a long time, it was thought that the earliest finds of ancient millet grains on the territory of modern Ukraine dated to the Neolithic. However, modern research methods have shown that the earliest millet grains known from Ukraine are no older than the Bronze Age. These small grains often move within and between archaeological layers, as they find their way through cracks, root channels, and rodent burrows or are carried by insects and rodents, ending up as younger intrusions in older layers. This is particularly evident at multi-period sites.

From the Bronze Age to the present day, common millet has been one of the main cultivated plants in the territory of Ukraine, associated both with agriculturalists and mobile pastoralists, such as the Scythians, who lived in villages and towns, and with nomads, such as the Cossacks. Pliny, Columella and other classical authors indicated the varied uses of common millet in ancient cuisines. This diversity was preserved over centuries and millennia, and the Cossacks of the Zaporizhzhya cooked various dishes from millet. The most popular dish was a porridge called kulish. Kulish recipes have survived to this day and are commonly used in Ukrainian cuisine.

Acknowledgements

I would like to thank my fellow archaeobotanists from the Institute for Pre- and Protohistory, Kiel University, Germany, for inviting me to take part in the workshop in November 2019 in Kiel. I am very pleased that, thanks to the workshop and this book, the results and insights from Ukraine can be presented, offering some new perspectives and helping answer some questions. I am grateful to Dragana Filipović, Marta Dal Corso and Wiebke Kirleis for their interest in my work, collegial attitude and support.

References

In Russian or Ukrainian

- Вавилов, Н.И., 1959. Избранные труды, том 1: Земледельческий Афганистан [Selected works, vol. 1: agricultural Afghanistan]. Москва-Ленинград: Академия наук СССР.
- Вавилов, Н.И., 1987. Происхождение и география культурных растений [The origin and geography of cultivated plants]. Ленинград: Наука.
- Вавилов, П.П., Гриценко, В.В., Кузнецов, В.С., 1986. Растениеводство [Growing plants]. Москва: Агропромиздат.
- Гаврилюк, Н.А. and Пашкевич Г.А., 1991. Земледельческий компонент в экономике степной Скифии. [The agricultural component of the economy of the steppe Scythians] *Советская археология*, 2, 51-63.
- Горбаненко, С.А. and Пашкевич, Г.О., 2010. Землеробство давніх слов'ян (кінець 1 тис. до н.е. – 1 тис. н.е.) [Farming of the ancient worlds (end of 1st century BCE-1st century CE)]. Київ: Академперіодика.
- Горбенко, К.В. and Пашкевич Г.О., 2010. Палеоетноботанічні дослідження на території городища Дикий Сад [Palaeoethnobotanical research on the territory of the Dikiy Sad hillfort]. Емінак: Науковий цоквартальник, 1-4 (5), 5-19.
- Жуковский, П.М., 1964. Культурные растения и их сородичи. Систематика, география, цитогенетика, иммунитет, экология, происхождение, использование. [Cultivated plants and their wild relatives: systematics, geography, cytogenetics, immunity, ecology, origin, use]. Москва: Колос.
- Козловська, Н.А., Пашкевич, Г.О., Хамайко, Н.В., 2012. Палеоботанічні дослідження матеріалів з розкопок по вул. Спаській, 35 у м. Києві [Palaeobotanical materials from the archaeological excavations at 35 Spas'ka str. in Kyiv]. *Археологічні дослідження в Україні 2011*, 257-258.
- Кузищин, И.М., 1966. Очерки по истории земледелия Италии II в. до н.э. – 1 в. н.э. [Essays on the history of agriculture in Italy, 2nd century BCE-1st century CE]. Москва: Издательство Московского Государственного Университета.
- Кузьминова, Н.Н. and Петренко, В.Г., 1989. Культурные растения на Западе Северного Причерноморья в середине 3 – 2-ом тыс. до н.э. (по данным

- палеоботаники) [Cultivated plants in the western northern Black Sea region from the middle of the 3rd through the 2nd millennium BCE (according to the palaeobotanical data)]. *Проблемы древней истории и археологии Украинской ССР*. Киев: Наукова думка, 119-120.
- Пашкевич, Г.А., 1980. Культурные растения трипольских поселений Поднепровья [Cultivated plants of the Trypolye settlements of the Dnieper region]. *Первобытная археология. Поиски и находки*. Киев: Наукова думка, 234-242.
- Пашкевич, Г.А., 1988а. Палеоботанический анализ злаков [The palaeobotanical analysis of cereals]. In: Е.В. Максимов, В.А. Петрашенко, eds., *Славянские памятники у с. Монастырек на Среднем Днепре*. Киев: Наукова думка, 131-134.
- Пашкевич, Г.А., 1988b. Палеоботанические исследования в области славянской археологии [Palaeobotanical research in the field of Slavic archaeology]. In: В.Д. Баран, ed., *Труды V Международного Конгресса археологов-славистов*. Т. 4 [Vol. 4] *Древние славяне*. Киев: Наукова думка, 169-174.
- Пашкевич, Г.А., 1989. Палеоботанические исследования трипольских материалов междуречья Днепра и Южного Буга [Palaeobotanical studies of the Trypillian materials in the Dnieper-Southern Bug interfluvium]. *Первобытная археология*. Киев: Наукова думка, 132-142.
- Пашкевич, Г.А., 1990а. Результаты палеоботанического исследования крупных трипольских поселений [Results of the palaeobotanical study of the large Trypillian settlements]. In: В.Г. Збенович, ed., *Тезисы докладов 1-го полевого семинара 'Раннеземледельческие поселения – гиганты Трипольской культуры на Украине'*, Тальянки – [Веселый Кут – Майданецкое]. Киев: Институт археологии национальной АН Украины, 131-134.
- Пашкевич, Г.А., 1990b. Состав культурных и сорных растений из раскопок поселений сельской округи Ольвии [Composition of the cultivated plants and weeds from the excavations of settlements in the rural area of Olbia]. *Античные поселения Нижнего Побужья (археологическая карта)*. Киев: Наукова думка, 114-119.
- Пашкевич, Г.А., 1991. Палеоэтноботанические находки на территории Украины: Памятники 1 тыс. до н.э. – II тыс. н.э. [Palaeoethnobotanical finds on the territory of Ukraine: sites from the 1st millennium BCE-2nd millennium CE]. Киев: Институт археологии национальной АН Украины.
- Пашкевич, Г.А., 1993. Особенности палеоэтноботанических комплексов энеолита – бронзы территории Украины [Characteristics of the palaeoethnobotanical complexes of the Eneolithic-Bronze Age in the Ukrainian territory]. In: P. Georgieva, ed., *The Fourth Millennium B.C.: Proceedings of the international symposium, Nessebar, 28-30 August 1992*. Sofia: New Bulgarian University, 99-108.
- Пашкевич, Г.А., 2000а. Земледелие в Степи и Лесостепи Восточной Европы в неолите – бронзовом веке (палеоэтноботанические свидетельства) [Agriculture in the steppe and forest-steppe of eastern Europe in the Neolithic through Bronze Age (palaeoethnobotanical evidence)]. *Stratum plus*. Археология и культурная антропология, 2, 404-418.
- Пашкевич, Г.А., 2000b. Палеоэтноботанические исследования скифских памятников степной зоны Северного Причерноморья [Palaeoethnobotanical investigations of the Scythian settlements in the steppe zone of the northern Black Sea region]. In: В.И. Гуляев, В.С. Ольховский, eds., *Скифы и сарматы в VII-III вв. до н.э. Палеоэкология, антропология и археология*. Москва: Институт археологии РАН, 101-109.
- Пашкевич, Г.А., 2001. Палеоботанические исследования материалов Мотронинского городища [Palaeobotanical analysis of the materials from the settlement of Motroninsky]. In: С.С. Бессонова, С.А. Скорый, eds., *Мотронинское городище скифской эпохи (по материалам раскопок 1988-1996 гг.)*. Киев, Краков: Институт археологии национальной АН Украины, 133-137.

- Пашкевич, Г.А., 2002. К проблеме изучения Северного Причерноморья в античную эпоху [On the problem of studying the northern Black Sea region in ancient times]. *Боспорский феномен. Погребальные памятники и святилища*. Санкт-Петербург: Издательство Государственного Эрмитажа, 302-311.
- Пашкевич, Г.О., 2004. Про склад рослин, вирощуваних на початку грецької колонізації Північного Причорномор'я [On the composition of plants cultivated at the beginning of the Greek colonisation of the northern Black Sea]. In: С.Д. Крыжицкий, ed., *Borysthenika-2004: Материалы международной научной конференции к 100-летию исследований острова Березань Э.Р. фон Штерном*. Николаев: Национальная Академия наук Украины, 131-138.
- Пашкевич, Г.А., 2005. Палеоэтноботанические исследования Ольвии и ее округи [Palaeoethnobotanical investigations of Olbia and its surroundings]. *Stratum plus*, Археология и культурная антропология, 3, 13-76.
- Пашкевич, Г.А., 2006. Современное состояние палеоэтноботанических исследований Херсонеса [The current state of research of the palaeoethnobotanical investigations at Chersonesos]. *Херсонесский сборник*, 15, 165-179.
- Пашкевич, Г.О., 2010. Палеоетноботанічні дослідження давньоруського часу та середньовіччя на території України [Palaeoethnobotanical research on the Old Russian time and the Middle Ages on the territory of Ukraine]. Проблеми давньоруської та середньовічної археології. Археологія і давня історія України 1. Київ: Інститут археології національної АН України, 477-483.
- Пашкевич, Г.О., 2011. Значення Бернашівки-1 для палеоетноботанічної характеристики раннього трипілля [The importance of Bernashivka-1 for the palaeoethnobotanical characterisation of the early Trypillia]. In: О.Г. Корвін-Піотровський, ed., *Прадавні землероби Південно-Східної Європи. Тези доповідей міжнародної наукової конференції до 30-річчя дослідження поселення-гіганта Тальянки та 10-річчя співпраці з 'Search Foundation' USA*. Київ, Тальянки: Інститут археології національної АН України, 65-68.
- Пашкевич, Г.А., 2016. Археоботанические исследования Боспора [Archaeobotanical studies in the Bosphorus]. *Боспорские исследования*, XXXII, 205-303.
- Пашкевич, Г.О. and Відейко, М.Ю., 2006. *Рільництво племен трипільської культури* [Agriculture of the Trypillia culture tribes]. Київ: Електронне видання.
- Пашкевич, Г.О. and Горбаненко, С.А., 2010. Каталог палеоетноботанічних матеріалів пам'яток слов'янських культур [Catalogue of the palaeoethnobotanical finds from the sites of the Slavic cultures]. Колекції наукових фондів Інституту археології НАН України. Матеріали та дослідження. (Vol. 3 in the series *Археологія та давня історія України*). Київ: Інститут археології національної АН України, 105-122.
- Пашкевич, Г.О. and Горбаненко, С.А., 2019. Палеоетноботанічні дослідження [Palaeoethnobotanical research]. In: С.А. Скорий, В.П. Білозор, О.Б. Супруненко, І.М. Кулатова, eds., *Селища скіфського часу в системі Великого укріплення Більського городища*. Київ: Майдан, 320-328.
- Пашкевич, Г.О. and Костильов, О.М., 1992. Синтаксономічний аналіз палеоботанічних даних на прикладі матеріалів епохи бронзи [Syntaxonomic analysis of the palaeobotanical data using the Bronze Age materials]. *Ойкумена. Український екологічний вісник*, 3, 72-77.
- Пашкевич, Г.О. and Шовкопляс, В.М., 2012. Особливості суббореального періоду голоцену [Characteristics of the Holocene Subboreal period]. *Тектоніка та стратиграфія*, 39, 145-151.
- Рева, М.Л. and Рева, Н.Н., 1976. *Дикі їстівні рослини України* [Wild edible plants of Ukraine]. Київ: Наукова думка.
- Сергеенко, М.Е., 1970. *Учение земледельцы Древней Италии* [Deliberations on agriculture in ancient Italy]. Перевод с латинского, примечания и введение. Ленинград: Наука.

- Сорочан, С.Б., Зубарь, В.М., Марченко, Л.В., 2001. *Жизнь и гибель Херсонеса* [The life and fall of Chersonesus]. Харьков: Майдан.
- Хименко, И.С., 1960. *Українські страви* [Ukrainian dishes]. Київ: Державне видавництво технічної літератури УРСР.
- Шалімов, С.А. and Шадура, О.А., 1981. *Сучасна українська кухня* [Modern Ukrainian cuisine]. Київ: Видавництво техніка.
- Янушевич, З.В., 1976. *Культурные растения юго-запада СССР по палеоботаническим исследованиям* [Cultivated plants of the southwestern USSR according to palaeobotanical studies]. Кишинев: Штиинца.
- Янушевич, З.В., 1980. Земледелие в раннем Триполье [Agriculture in the early Tripolye]. In: И.И. Артеменко, ed., *Первобытная археология – поиски и находки: сборник научных трудов*. Киев: Наукова думка, 225-234.
- Янушевич, З.В., 1986. *Культурные растения Северного Причерноморья. Палеоэтноботанические исследования* [Cultivated plants of the northern Black Sea region: palaeoethnobotanical investigations]. Кишинев: Штиинца.

In English

- An, T., Pashkevich, G., Jones, M., 2018. Re-examining millet impressions in Usatovo clay materials from NW Black Sea region, Ukraine. *Archaeological and Anthropological Sciences*, 11, 3201-3211. <https://doi.org/10.1007/s12520-018-0718-3>
- Brownson, C.L., Marchant, E.C., Todd, O.J., Miller, W., 1922. *Xenophon in Seven Volumes*. Cambridge, MA: Harvard University Press.
- Dal Corso, M., Pashkevych, G., Filipović, D., Liu, X., Motuzaite-Matuzeviciute, G., Stobbe, A., Shatilo, L., Videiko, M., Kirleis, W., 2022. Between cereal agriculture and animal husbandry: millet in the early economy of the North Pontic region. *Journal of World Prehistory*.
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Kloof, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A.M., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H.-P., Stobbe, A., Tolar, T., Wasylkowa, K., Wiethold, J., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Godley, A.D., trans. 1920. *Herodotus. The Histories*. Cambridge, MA: Harvard University Press. In: G.R. Crane (ed.) *Perseus Digital Library*. Web project at Tufts University [online]. <http://www.perseus.tufts.edu/hopper/text?doc=Perseus%3A-text%3A1999.01.0126%3Abook%3D4%3Achapter%3D17%3Asection%3D1>
- Hunt, H.V., Vander Linden, M., Liu, X., Motuzaite-Matuzeviciute, G., Colledge, S., Jones, M.K., 2008. Millet across Eurasia: chronology and context of early records of the genera *Panicum* and *Setaria* from archaeological sites in the Old World. *Vegetation History and Archaeobotany* 17, S5-S18. <https://doi.org/10.1007/s00334-008-0187-1>
- Janushevich, Z.V., 1981. Die Kulturpflanzen Skythiens. *Zeitschrift für Archäologie*, 15 (1), 87-96.
- Janushevich, Z.V., 1989. Agricultural evolution north of Black Sea from the Neolithic to the Iron Age. In: D.R. Harris and G.C. Hillman, eds., *Foraging and Farming: the Evolution of Plant Exploitation*. London: Unwin Hyman, 607-619.
- Janushevich, Z.V. and Nikolaenko, G.M., 1979. Fossil remains of cultivated plants in the ancient Tauric Chersonesos. In: U. Körber-Grohne, ed., *Festschrift Maria Hopf: zum 65. Geburtstag am 14. September 1979*. Archaeo Physika 8. Köln: Rheinland Verlag, 115-134.

- Kirleis, W. and Dal Corso, M., 2016. Trypillian subsistence economy: animal and plant exploitation. In: J. Müller, K. Rassmann, M. Videiko, eds., *Trypillia Mega-sites and European Prehistory 4100-3400 BCE*. Themes in Contemporary Archaeology 2. London and New York: Routledge, 195-205.
- Motuzaitė-Matuzevičiūtė, G., Staff, R.A., Hunt, H.V., Liu, X., Jones, M.K., 2013. The early chronology of broomcorn millet (*Panicum miliaceum*) in Europe. *Antiquity*, 87 (338), 1073-1085. <https://doi.org/10.1017/S0003598X00049875>
- Müller, J., Hofmann, R., Brandtstätter, L., Ohlrau, R., Videiko, M., 2016. Chronology and demography: how many people lived in a mega-site? In: J. Müller, K. Rassmann, M. Videiko, eds., *Trypillia Mega-sites and European Prehistory 4100-3400 BCE*. Themes in Contemporary Archaeology 2. London and New York: Routledge, 133-170.
- Pashkevich, G., 1984. Palaeoethnobotanical examination of archaeological sites in the lower Dnieper region, dated to the last centuries BC and first centuries AD. In: W. Van Zeist and W.A. Casparie, eds., *Plants and Ancient Man: Studies in Palaeoethnobotany*. Boston: A.A. Balkema, 277-284.
- Pashkevich, G., 2003. Palaeoethnobotanical evidence of agriculture in the steppe and the forest-steppe of East Europe in the Late Neolithic and Bronze Age. In: M. Levine, C. Renfrew, K. Boyle, eds., *Prehistoric steppe adaptation and the horse*. Cambridge: McDonald Institute for Archaeological Research, 287-297.
- Pashkevich, G., 2005. Palaeoethnobotanical evidence of the Tripolye culture. In: G. Dumitroaia, J. Chapman, O. Weller, C. Preoteasa, R. Monteanu, D. Nicola, D. Monah, eds., *Cucuteni: 120 ans de recherches: le temps du bilan*. Bibliotheca memoriae antiquitatis 16. Piatra Neamț: Editura Constantin Matasă, 231-245.
- Pashkevych, G., 2012. Environment and economic activities of Neolithic and Bronze Age populations of the Northern Pontic area. *Quaternary International*, 261, 176-182. <https://doi.org/10.1016/j.quaint.2011.01.024>
- Pashkevych, G., 2016. Data of the Trypillian agriculture: new palaeobotanical evidence. In: C. Preoteasa and C.D. Nicola, eds., *Cucuteni Culture within the European Neo-Eneolithic Context*. Proceedings of the International Colloquium 'Cucuteni 130', 15-17 October 2014, Pietra Neamț, Romania. Piatra Neamț: Editura Constantin Matasă, 24-27.
- Rackham, H., 1938. *Natural History by Pliny the Elder*. London: W. Heinemann. <https://archive.org/details/naturalhistory05plinuoft/page/252/mode/2up?view=theater>
- Zohary, D. and Hopf, M., 2000. *Domestication of Plants in the Old World*. (3rd edition). Oxford: Oxford University Press.

Bronze Age plant spectra in Hungary before and after the introduction of millet cultivation

Sofia Filatova

Abstract

The current paper provides an overview of the plant spectra from archaeological sites dating to the Bronze Age in Hungary, and seeks to identify changes and continuities therein before and after the establishment of broomcorn millet as a crop. The overview includes a discussion of the general trends that have been identified in the Bronze Age habitation and environment in the region, as well as a quantitative consideration of macro-botanical data from 52 Bronze Age sites in Hungary. The results confirm the introduction of millet cultivation in Hungary during the Late Bronze Age and further demonstrate the diverse spectra of plants within individual settlements of all three phases of the Bronze Age.

Introduction

Macro-botanical finds of broomcorn millet (*Panicum miliaceum*) in modern-day Hungary have been reported dating as early as the Neolithic (6th millennium BCE; Gyulai 2014, 32). Recent AMS-dating of charred broomcorn millet has, however, revealed that the earliest evidence of millet in Hungary dates to the 15th century cal BCE (the onset of the Late Bronze Age), indicating the redeposition of the small grains from younger to older archaeological layers (Filipović *et al.* 2020, 6, Table 2). Isotopic studies of human bone samples from the Great Hungarian Plain dating from the Neolithic to the Bronze Age have, furthermore, revealed an increase in C4 plant dietary input from the Late Bronze Age onwards, suggesting the inclusion of millet in the diet of humans and/or animals (Gamarra *et al.* 2018, 12-13). The Bronze Age has also been distinguished as the time when millet cultivation was introduced into other regions of Europe (Filipović *et al.* 2020, 13).

The current paper provides an overview of the spectra of plants from archaeological sites dating to the Bronze Age in Hungary, and seeks to identify changes and continuities therein before and after the establishment of broomcorn

millet as a crop. The paper starts with brief discussion on the general trends identified in Bronze Age habitation and environment, in order to contextualise the situation in which millet appeared in the region. This is followed by an introduction to the plant spectra based on macro-botanical studies from Bronze Age settlements throughout Hungary. The paper concludes with a consideration of the patterns that could be distinguished from the assemblages. Note that part of the information presented below applies to the Carpathian Basin as a whole, in which Hungary is situated.

Hungarian Bronze Age chronology and society

The Hungarian chronology of the Bronze Age was established by the 1980s (Fischl *et al.* 2015, 505) and an important part of it is based on Istvan Bóna's (1975) analysis of material from more than 1200 archaeological sites (Bóna 1975, 13). Although it has been further elaborated and refined since, its foundation has remained unaltered and it is still in use today (Fischl *et al.* 2015, 505). The Bronze Age in Hungary is divided into three main phases (Early, Middle and Late), and together these cover a period that starts in the second half of the 3rd millennium BCE and ends at the beginning of the 1st millennium BCE (Table 1).

A diverse array of communities, or archaeologically determined groups, inhabited the Carpathian Basin during the Bronze Age, and they are distinguished based on ceramic styles, burial customs and types of settlement. Traditionally, the areas occupied by different archaeological groups have been defined on the basis of uniformity in material culture, thereby forming 'cultural territories' throughout the Carpathian Basin (*e.g.* Bóna 1992, 16-17). Although this view has been criticised due to the underlying assumption that there exists a direct relationship between ethnicity and material culture, the traditional definition of Bronze Age groups and their related territories remain in use within Hungarian archaeology (Kienlin 2012, 2015, 36; Staniuk 2020, 38).

Starting with the transition to the Early Bronze Age (2800/2700 cal BCE), a group known as Makó-Kosihy-Čaka inhabited areas in northern, central and eastern Hungary, occupying small campsites as well as larger settlements (Kulcsár 2009, 14-16, 64). A few centuries later (from ca. 2600/2500 cal BCE), a variety of groups become visible in the archaeological record of the Carpathian Basin, including Bell Beaker groups inhabiting the area north of Budapest and Nagyrév groups first settling on the western bank of the river Danube (for a more elaborate description, see Fischl *et al.* 2015, 505-509). Some of these groups vanished in the course of the Early Bronze Age (*e.g.* the Bell Beaker groups), while others persisted and laid the foundations for subsequent Middle Bronze Age groups (Kienlin 2015, 33). Of special importance are Nagyrév groups, who in some cases constructed their houses one on top of the other in succession, creating superposed building horizons eventually leading to the formation of tells, which were occasionally fortified. This practice spread along the Danube and the Tisza, and to the northern and eastern parts of the Carpathian Basin (Fischl *et al.* 2015, 513). The occupation of tells is one of the defining features of the Early and Middle Bronze Age in central and eastern Hungary, although only small parts of the inhabited regions actually contained tells (Gogâltan 2008, 40; Kienlin 2015, 34). Tell-like, mound-like and flat settlements were inhabited alongside tells, while in Transdanubia (western Hungary) tells never appeared and settlements were occupied for shorter periods (Fischl *et al.* 2015, 514).

During the Middle Bronze Age (2000/1900-1500/1450 cal BCE), a number of tells that had been established in the Early Bronze Age were continuously inhabited by the succeeding Middle Bronze Age groups (*e.g.* Vátya groups on the Nagyrév tells). This continuity in occupation is attributed to population concentration and/or growth, which is also reflected in the increase in the number and size of settlements

Bronze Age phase	Date (cal BCE)	Important characteristics
Late Bronze Age II	1250/1150-800/750	Entire Carpathian Basin occupied by local groups of the Urnfield culture; settlements include campsites, hamlets and fortified hilltop settlements; bronze metallurgy becomes widespread.
Late Bronze Age I	1500/1450-1250/1150	Entire Carpathian Basin occupied by the Tumulus culture; settlements are more dispersed and sizes decrease compared with previous phase.
Middle Bronze Age	2000/1900-1500/1450	Continuity of tell occupation in the central and eastern Carpathian Basin by a variety of communities; settlements increase in size and number.
Early Bronze Age	2600/2500-2000/1950	Appearance of a variety of communities; onset of development of tells in the central and eastern Carpathian Basin.
Copper Age/Early Bronze Age	2800/2700-2600/2500	Transitional phase: Early Bronze Age settlements ranging from larger, more permanent sites to small, temporary campsites.

Table 1. Chronology of the Hungarian Bronze Age and summarised characteristics based on the information provided in the text. Dates after Fischl *et al.* (2015, 504-505, Table 1a, b).

(Szeverényi and Kulcsár 2012, 292; Fischl *et al.* 2015, 513). In some cases, (fortified) tells are surrounded by smaller settlements and a hierarchical relationship has been proposed, whereby tells, occupied by an elite, acted as primary centres and controlled movement and communication in the surroundings (Gogâltan 2008, 53; Earle and Kolb 2010, 72-73). This is, however, not the case in all regions of the Carpathian Basin (e.g. Szeverényi and Kulcsár 2012, 336) and because very few 'satellite settlements' have been systematically excavated, this hierarchical relationship remains speculative (Kienlin 2012, 282; Jaeger 2016, 69; Staniuk 2020, 43).

The end of the Middle Bronze Age is marked by the gradual abandonment of tells and by the burial of bronze hoards, the latter practice happening during what is known as the Koszider horizon (Poroszlai 2003, 155; Jaeger 2016, 99-100). Although this practice was initially interpreted as a rapid event stimulated by a violent invasion by the Tumulus people from the west, the absence of evidence for settlement destruction and the increasing evidence for gradual settlement abandonment, both prior to and after the Koszider horizon, have altered this view (Fischl *et al.* 2013, 358; Vicze 2013, 15; Jaeger 2016, 99-100; Duffy *et al.* 2019).

During the first part of the Late Bronze Age (1500/1450-1200/1150 cal BCE), the open, more dispersed and generally smaller settlements of the Tumulus groups came to dominate the landscape (Sánta 2010, 257). These groups occupied the entire region of the Carpathian Basin and are especially known for their distinctive burials, in kurgans or tumuli (Csányi 2003, 162). There is no consensus on what may have caused the sudden decline in number and size of settlements, but explanations that have been put forward include a shift from intensive to extensive agricultural practices, with a more pronounced reliance on animal husbandry (Sánta 2010, 522), and a reduction in agricultural yields due to (a) climatic deterioration, (b) scarcity in available arable land, and/or (c) exhaustion of the cultivated soil (Gyulai 1993, 19; Reményi 2012, 281; Fischl *et al.* 2013, 366).

The final phase of the Late Bronze Age (1250/1150-800/750 cal BCE) witnessed another major change in occupation. The Carpathian Basin was now inhabited by a range of local Urnfield groups, such as the Gáva in the east and the Piliny and Kyjatice in the west (Marková and Ilon 2013, 827). These groups occupied a variety of (marginal) landscapes, ranging from the plains to the mountains, and their settlements included large, fortified hilltop settlements and small hamlets and campsites (Szábo 2003, 162-164; Marková and Ilon 2013, 827-828). The Urnfield groups are especially known for their burial rite, which took the form of cremations in urns, and for their extensive development of bronze metallurgy and trade. The bronze metallurgy was characterised by a variety of tools that were no longer solely prestige items, but became objects of common use (Szábo 2003, 164; Marková and Ilon 2013, 827).

Bronze Age environment

The Bronze Age in Hungary coincides with the Subboreal biozone, which spans the Late Copper Age to the start of the Iron Age (ca. 3600/3500-900/800 cal BCE; Somogyi 1987, 30). In contrast to the preceding Atlantic phase, which witnessed the Holocene climatic optimum, the Subboreal is distinguished by cooler summer temperatures and high levels of humidity (Kordos 1987, 13-14, Figs. 2, 3). Stable hydrogen, oxygen and carbon isotope data obtained from cave stalagmites in southern Hungary and from bivalve shells from Lake Balaton in western Hungary reflect a short-term humid phase during the first part of the Middle Bronze Age, with a subsequent increase in humidity during the second part of the Middle Bronze Age. After the humidity peak, precipitation decreased and gave way to drier and warmer summer temperatures during the second phase of the Late Bronze Age. These conditions have been interpreted to reflect a period of fluctuating humidity but with a relatively mild climate, suitable for arable farming (Demény *et al.* 2019).

Despite the mosaic-like character of the vegetation that prevailed in Hungary during the Bronze Age, a few general commonalities can be observed. Although the exact combination of taxa varied locally, the arboreal pollen in the records in northeastern, central and western Hungary show a strong presence of species indicative of mixed oak forests (*e.g.* species of oak (*Quercus* sp.), elm (*Ulmus* sp.) and lime (*Tilia* sp.)) throughout the entire period and an increase in hornbeam (*Carpinus* sp.) and beech (*Betula* sp.) towards the Late Bronze Age (Sümegi and Bodor 2000, 89; Juhász 2007a, 34, 2007b, 47-49; Juhász and Szegvári 2007, 324; Magyar *et al.* 2010, 926-927; Náfrádi *et al.* 2011, 363). The increasing presence of hornbeam and beech can be seen as an indicator for more arid and humid conditions, respectively (Magyar *et al.* 2010, 927), but their combined presence can more likely be interpreted as a reflection of woodland clearance. The exploitation of oak, elm and lime for the construction of houses and fortifications during the Bronze Age would have promoted the spread of hornbeam and beech (Sümegi and Bodor 2000, 89; Náfrádi *et al.* 2011, 361). At least part of the mixed oak forest that prevailed in Hungary during the Bronze Age was in the form of a forest-steppe vegetation, the so-called Pannonian steppe, which existed throughout the semi-arid regions of the Carpathian Basin during most of the Holocene (Nagy-Bodor *et al.* 2000, 121; Sümegi *et al.* 2012, 17). Over the long term, the exploitation of oak, elm and lime caused thinning of the forest, blurring the boundaries between closed woodlands, gallery (riverine) forests and forest-steppe, and an increase in grasslands, eventually resulting in an expansion of the forest-steppe (Sümegi *et al.* 2012, 18).

From the second half of the Middle Bronze Age onwards, anthropogenic indicators, such as plantain (*Plantago* sp.) and dock (*Rumex* sp.), are more pronounced in the pollen record than before and suggest disturbance through pastoral activities and arable agriculture (Sümegi and Bodor 2000, 87; Juhász 2007a, 34, 2007b, 47-49; Juhász and Szegvári 2007, 324; Magyar *et al.* 2010, 926-927; Náfrádi *et al.* 2011, 362, Fig. 3; Knipl and Sümegi 2012, 444). This is in line with the archaeological evidence presented above that indicates flourishing of communities and a population growth within the Carpathian Basin from the Middle Bronze Age onwards.

Macro-botanical finds of the Bronze Age in Hungary

The material

Macro-botanical material retrieved from Bronze Age settlements in Hungary has been intensively studied during two phases: in the 1960s-1970s and in the 1990s-2000s (Gyulai 2010, Appendix Bronze Age data). An extensive compilation of macro-botanical data in Hungary was published by Gyulai in 2010, and this publication forms the main source of information in the current contribution. Gyulai's data have been supplemented by recent macro-botanical studies of Tarhos Gyepesi Átkelő (Tarhos 26; Békés County, Hungary) by Duffy (2010, 275, Table 9.1); Százhalombatta-Földvár (Pest County, Hungary) by Stika and Heiss (2013, 79-80, Table 1); and Kakucs-Turján (Pest County, Hungary) by the present author (Filatova 2020).

The data compiled by Gyulai have a few drawbacks that need to be addressed. First, as is inevitable with macro-botanical studies that span several decades, the sampling strategies applied to retrieve the remains were not uniform. Whereas some sites have been sampled systematically, most have not, and there is therefore a discrepancy in the reliability of the data (e.g. Gyulai 2010, 100). These data are nevertheless of great value to our understanding of the diversity of plant spectra of the Bronze Age, especially when viewed on a regional scale.

Second, the methods of quantification in Gyulai's compilation vary. Most of the remains have been recorded in absolute quantities, but records also include semi-quantitative estimates (e.g. x, xx, xxx) and quantifications in weight and volume. In order to include the entire body of data, these quantifications were converted into absolute numbers according to the guidelines shown in Tables 2 and 3. The semi-quantitative ranges were converted into the median values of those ranges. The conversions of weight and volume into numbers were calculated based on 100 remains of the same species from the Early and Middle Bronze Age settlement of Kakucs-Turján in central Hungary. Exceptions are grass pea (*Lathyrus sativus*), club wheat (*Triticum aestivum* ssp. *compactum*) and millet (*Panicum miliaceum*), which either were not identified in the samples of Kakucs-Turján or were only represented by a few grains/seeds per sample. Grass pea was estimated as equal to the weight and volume of pea (*Pisum sativum*), and club wheat as falling between the weight and volume of barley (*Hordeum vulgare*) and einkorn (*Triticum monococcum* ssp. *monococcum*). The volume and weight of 100 grains of millet were measured based on the same finds from the Byzantine site of Caričin Grad (Lebane municipality, Serbia; Reuter in press), which were readily available for measurement in the archaeobotanical laboratory at Kiel University. Despite the fact that the converted values probably diverge from the actual values, they have the advantage that they

Table 2. Weights and volumes of grains and seeds from Kakucs-Turján and Caričin Grad used to convert weights and volumes into absolute quantities in the dataset compiled by Gyulai (2010). The weights and volumes are given per 100 grains or seeds. Due to the unavailability of the sufficient amount of material, the weight and volume of grass pea is estimated as equal to that of pea and the weight and volume of club wheat as falling between the weight and volume of einkorn and barley. The volume and weight of millet grains is based on the volume and weight 100 millet grains from Caričin Grad (Reuter in press).

Scientific name	Weight (g)	Volume (ml)	Common name
<i>Hordeum vulgare</i>	1.1	3.5	Barley
<i>Panicum miliaceum</i>	0.1	0.5	Broomcorn millet
<i>Triticum monococcum</i> ssp. <i>monococcum</i>	0.9	3	Einkorn
<i>Triticum turgidum</i> ssp. <i>dicoccon</i>	1	3.25	Emmer
<i>Triticum aestivum</i> ssp. <i>compactum</i>	1	3.25	Club wheat
<i>Lathyrus sativus</i>	2.6	5.5	Grass pea
<i>Lens culinaris</i>	0.7	1.5	Lentil
<i>Pisum sativum</i>	2.6	5.5	Pea

Table 3. Absolute quantities used as a basis for conversion of the semi-quantitative indications in the dataset compiled by Gyulai (2010).

Category	Original quantity	New quantity
x	1-10	5,5
xx	10-100	55
xxx	100-1000	550
xxxx	1000-10,000	5500

Site	Phase*	Taxon	Plant part	Preservation	Original quantification	Adapted quantification
Tiszaalpár-Várdomb	MBA	<i>Hordeum vulgare</i> subsp. <i>hexastichum</i>	Grain	Charred	10 g + 59	968
Tiszaalpár-Várdomb	MBA	<i>Hordeum vulgare</i> subsp. <i>tetrastichon</i>	Grain	Charred	14 g + 540	1813
Tiszaalpár-Várdomb	MBA	<i>Hordeum vulgare</i> var. <i>nudum</i>	Grain	Charred	5.5 g	500
Tiszaalpár-Várdomb	MBA	<i>Hordeum vulgare</i>	Grain	Charred	93.4 g + 3960	12,451
Tiszaalpár-Várdomb	MBA	<i>Lathyrus sativus</i>	Seed	Charred	101 g + 744	4629
Tiszaalpár-Várdomb	MBA	<i>Lens culinaris</i>	Seed	Charred	20.6 g + 3253	2943
Tiszaalpár-Várdomb	MBA	<i>Triticum aestivum</i> ssp. <i>compactum</i>	Grain	Charred	11 g + 35	1135
Tiszaalpár-Várdomb	MBA	<i>Triticum monococcum</i>	Grain	Charred	71.4 g + 3930	11,863
Tiszaalpár-Várdomb	MBA	<i>Triticum dicoccon</i>	Grain	Charred	23.9 g + 372	1107
Tiszaalpár-Várdomb	MBA	<i>Triticum spec.</i>	Grain	Charred	722 g + 1988	2710
Százhalmabatta-Téglagyár	MBA	<i>Triticum monococcum</i>	Grain	Charred	480 cm ³ + 592	16,592
Százhalmabatta-Téglagyár	MBA	<i>Triticum dicoccon</i>	Grain	Charred	500 cm ³ + 32	15,417
Pákozd-Várhegy	MBA	<i>Pisum sativum</i>	Seed	Charred	80 cm ³	1455
Pákozd-Várhegy	MBA	<i>Triticum monococcum</i>	Grain	Charred	40 cm ³ + 104	1437
Mende-Leányvár	MBA	<i>Hordeum vulgare</i>	Grain	Charred	20 cm ³ (=20 ml)	100
Mende-Leányvár	MBA	<i>Hordeum vulgare</i> subsp. <i>distichon</i>	Grain	Charred	20 cm ³ (=20 ml)	100
Mende-Leányvár	MBA	<i>Triticum monococcum</i>	Grain	Charred	520 cm ³	17,333
Mende-Leányvár	MBA	<i>Triticum dicoccon</i>	Grain	Charred	1000 cm ³	30,769
Jászdózsa-Kápolnahalom	MBA	<i>Triticum dicoccon</i>	Grain	Charred	4 cl	123
Balatonboglár-Szárszó	MBA	<i>Triticum monococcum</i>	Grain	Charred	20 cl	6666
Balatonboglár-Szárszó	MBA	<i>Triticum dicoccon</i>	Grain	Charred	50 cl	15,385
Poroszló-Aponhát	LBA	<i>Panicum miliaceum</i>	Grain	Charred	20 cl	4000

Table 4. List of sites and taxa from the dataset published by Gyulai (2010), originally recorded in weights or volumes and converted for the current overview based on the values in Table 2. The names of the sites and taxa represent the original names reported by Gyulai, although some have been adapted in the current study. * MBA=Middle Bronze Age; LBA=Late Bronze Age.

allow for the inclusion of all data known from the Bronze Age in Hungary and therefore the potential error created by the conversion is considered acceptable. The converted values per site included in the dataset are given in Tables 4 and 5.

Furthermore, the Bronze Age chronology used by Gyulai in his database does not correspond to the chronological framework suggested more recently for Hungary. The dates have therefore been adjusted according to the data provided by Fischl *et al.* (2015, 504-505, Tables 1a-b). In a few cases, the classification of archaeological groups likewise deviated from recent classifications; where necessary, these were adjusted according to information extracted from Figler *et al.* (1997), Horváth (2005), Kulcsár (2009), Fülöp (2017), Filatova (2020) and the Hungarian National Museum Archaeology Database (2020).

Lastly, Gyulai's data are presented based on summarised remains per site and there are no contextual data that indicate the number of remains per archaeological context per site. Although this does not form a major drawback in the current paper,

Site	Taxon	Plant part	Phase	Preservation	Original quantification	Adapted quantification
Poroszló-Aponhát	<i>Triticum aestivum</i> ssp. <i>spelta</i>	Grain	LBA	Charred	xxx	550
Poroszló-Aponhát	<i>Triticum aestivum</i> ssp. <i>spelta</i>	Spikelet fork	LBA	Charred	x	5.5
Poroszló-Aponhát	<i>Triticum monococcum</i>	Spikelet fork	LBA	Charred	x	5.5
Poroszló-Aponhát	<i>Triticum turgidum</i> ssp. <i>dicocon</i>	Spikelet fork	LBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Atriplex patula</i>	Seed	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Cicer arietinum</i>	Seed	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Equisetum arvense</i>	Stem	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Hordeum vulgare</i>	Grain	MBA	Charred	xxx	550
Tószeg-Laposhalom	<i>Hordeum vulgare</i> ssp. <i>distichum</i>	Grain	MBA	Charred	xxx	550
Tószeg-Laposhalom	<i>Lathyrus sativus</i>	Seed	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Lens culinaris</i>	Seed	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Pisum sativum</i>	Seed	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Pyrus</i> sp.	Seed	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Triticum aestivum</i> ssp. <i>vulgare</i>	Grain	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Triticum monococcum</i>	Grain	MBA	Charred	x	5.5
Tószeg-Laposhalom	<i>Vicia angustifolia</i>	Seed	MBA	Charred	x	5.5
Tiszafüred-Ásotthalom	<i>Panicum miliaceum</i>	Grain	MBA	Charred	xxx	550
Tiszaalpár-Várdomb	<i>Triticum monococcum</i>	Grain	MBA	Charred	xx	55
Síhalom-Földvár	<i>Cucumis</i> sp.	Seed	MBA	Charred	x	5.5
Síhalom-Földvár	<i>Panicum miliaceum</i>	Grain	MBA	Charred	x	5.5
Százhalombatta-Téglagyár	<i>Hordeum vulgare</i> ssp. <i>distichum</i>	Grain	MBA	Charred	xx	55
Százhalombatta-Téglagyár	<i>Triticum monococcum</i>	Spikelet fork	MBA	Charred	x	5.5
Solymár-Várhegy	<i>Hordeum vulgare</i>	Grain	MBA	Charred	x	5.5
Mende-Leányvár	<i>Triticum dicocon</i>	Spikelet fork	MBA	Charred	xx	55
Mende-Leányvár	<i>Triticum monococcum</i>	Spikelet fork	MBA	Charred	xx	55
Jászdózsa-Kápolnahalom	<i>Triticum turgidum</i> ssp. <i>dicocon</i>	Spikelet forks	MBA	Charred	x	5.5
Felsődobsza-Várdomb	<i>Hordeum vulgare</i> var. <i>nudum</i>	Grain	MBA	Charred	x	5.5
Felsődobsza-Várdomb	<i>Lens culinaris</i>	Seed	MBA	Charred	x	5.5
Felsődobsza-Várdomb	<i>Triticum aestivum</i> ssp. <i>vulgare</i>	Grain	MBA	Charred	x+9	14.5
Felsődobsza-Várdomb	<i>Triticum monococcum</i>	Grain	MBA	Charred	x+4	9
Felsődobsza-Várdomb	<i>Triticum vulgare antiquorum</i>	Grain	MBA	Charred	x	5.5
Dunaújváros-Kosziderpadlás	<i>Hordeum vulgare polystichum</i> (H. <i>tetrastichum</i> +H. <i>hexastichum</i>) var. <i>nudum</i>	Grain	MBA	Charred	x	5.5
Dömsöd-Apaj	<i>Bromus</i> sp.	Grain	MBA	Charred	x	5.5
Dömsöd-Apaj	<i>Hordeum vulgare</i>	Grain	MBA	Charred	x	5.5
Dömsöd-Apaj	<i>Triticum monococcum</i>	Spikelet fork	MBA	Charred	x	5.5
Dömsöd-Apaj	<i>Triticum turgidum</i> ssp. <i>dicocon</i>	Spikelet fork	MBA	Charred	x	5.5

Table 5. List of sites and taxa from the dataset published by Gyulai (2010) that were originally recorded as semi-quantitative values and were converted for the current overview based on the values in Table 3. The names of the sites and taxa represent the original names reported by Gyulai, although some of these have been adapted in the current study.

Site	Taxon	Plant part	Phase	Preservation	Original quantification	Adapted quantification
Békés-Várdomb	<i>Agrostemma githago</i>	Seed	MBA	Charred	x	5.5
Békés-Várdomb	<i>Calystegia</i> sp.	Seed	MBA	Charred	x	5.5
Békés-Várdomb	<i>Hordeum vulgare</i>	Grain	MBA	Charred	x	5.5
Békés-Várdomb	<i>Melampyrum</i> sp.	Seed	MBA	Charred	x	5.5
Békés-Várdomb	<i>Quercus</i> sp.	Glans	MBA	Charred	x	5.5
Békés-Várdomb	<i>Triticum monococcum</i>	Grain	MBA	Charred	x	5.5
Baracs-Bottyánsánc	<i>Hordeum vulgare polystichum</i> (<i>H. tetrastichum</i> + <i>H. hexastichum</i>)	Grain	MBA	Charred	x	5.5
Baracs-Bottyánsánc	<i>Triticum monococcum</i>	Grain	MBA	Charred	x	5.5
Ároktő-Dongóhalom	<i>Secale cereale</i>	Grain	MBA	Charred	xx	55
Ároktő-Dongóhalom	<i>Triticum aestivum</i> ssp. <i>vulgare</i>	Grain	MBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Agrimonia eupatoria</i>	Glans	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Agropyron repens</i>	Grain	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Bromus tectorum</i>	Grain	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Camelina sativa</i>	Seed	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Centaurea</i> sp.	Achenium	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Convolvulus arvensis</i>	Seed	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Datura stramonium</i>	Seed	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Hordeum vulgare</i>	Grain	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Lens culinaris</i>	Seed	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Lolium</i> sp.	Grain	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Malva</i> sp.	Seed	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Pisum sativum</i> ssp. <i>microspermum</i>	Seed	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Rumex acetosella</i>	Glans	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Rumex crispus</i>	Glans	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Rumex</i> sp.	Glans	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Triticum aestivum</i> ssp. <i>compactum</i>	Grain	EBA	Charred	x	5.5
Pécs-Nagyárpád	<i>Verbascum</i> sp.	Seed	EBA	Charred	x	5.5

Table 5. continued.

since it represents a regional overview, it prohibits the calculation of more nuanced values, such as relative frequencies of plant remains expressed as percentages of the total number of samples.

The resulting data comprise a total of 52 sites (Fig. 1, Table 6) dating to the Early (n=12), Middle (n=25) and Late (n=15) Bronze Age. The majority of the finds consist of charred macro-botanical remains, although a few sites also include waterlogged finds (designated with a (w)). For practical reasons, species of cereals and pulses (e.g. *Triticum* sp. and cf. *Vicia faba*) have been merged into the categories of ‘Cerealia indet.’ and ‘Leguminosae sativae indet.’.

Table 6 (right and overleaf). Overview of the archaeological sites of the Bronze Age in Hungary that have been included in the current contribution. The assignments of the sites to archaeological groups partially rely on the data in Figler et al. 1997; Horváth 2005; Kulcsár 2009; Fülöp 2017; Filatova 2020; Hungarian National Museum Archaeology Database 2020. Note that the database in Gyulai (2010) included Százhalombatta-Földvár and Százhalombatta-Téglagyár as separate entries. According to the Hungarian National Museum Archaeology Database (2020), the latter is a synonym of the former; the two sets of data have therefore been merged here under the name Százhalombatta-Földvár.

	Site	Archaeological group	Number of samples	Total number of remains
Late Bronze Age (n=15)	Mosonmagyaróvár-Németbánya	Unknown	1	21
	Lébény-Billedomb	Unknown	2	1875
	Győr-Szabadrétdomb	Unknown	4	13
	Százhalombatta-Földvár	Urnfield	2	851
	Sopron-Krautacker 1	Urnfield	1	37
	Gór-Kápolnadomb	Urnfield	16	20,420
	Budapest, Albertfalva-Kitérő út	Urnfield	4	780
	Polgár 31	Tumulus	6	393
	Mosonmagyaróvár-Németdűlő (w)	Tumulus	1	1232
	Isztimér-Csőszpuszta	Tumulus	Unknown	264
	Dunakeszi-Székesdűlő (Auchan) (w)	Tumulus	35	1692
	Börcs-Paphomlok	Tumulus	10	354
	Balatonmagyaród-Hídvégpuszta	Tumulus	1	2304
	Ludas, Varjú-dűlő	Kyjatice	231	4928
	Poroszló-Aponhát	Gáva	2	9214.5
Middle Bronze Age (n=25)	Balatonboglár-Szárszó	Unknown	Unknown	22,078
	Tiszaalpár-Várdomb (w)	Vatya	43	42,444
	Százhalombatta-Földvár	Vatya	257	55,031.5
	Solymár-Várhegy	Vatya	1	5.5
	Pákozd-Várhegy	Vatya	Unknown	2953
	Mende-Leányvár (w)	Vatya	4	46,501
	Kakucs-Turján	Vatya	369	81,447.5
	Dömsöd-Apaj	Vatya	Unknown	22
	Cegléd 4/1	Vatya	1	26
	Budapest, Bocskai-Fehérvári úti aluljáró	Vatya	6	408
	Ménfőcsanak-Szeles (w)	Transdanubian Encrusted Pottery	32	298
	Túrkeve-Terehalom	Otomani	4	75,315
	Tarhos Gyepesi Átkelő (Tarhos 26)	Otomani	10	150
	Békés-Várdomb	Otomani	Unknown	38.5
	Dunaújváros-Kosziderpadlás	Nagyrév/Vatya	1	479.5
	Bölcske-Vörösgyír (w)	Nagyrév/Vatya	Unknown	38,030.5
	Baracs-Bottyánsánc	Nagyrév/Vatya	1	921
	Budapest, Csepel-Szenyvíz telep (w)	Nagyrév/MBA	13	74
	Tószeg-Laposhalom	Nagyrév/Hatvan	Unknown	17,764
	Süttő-Hosszúvölgy	Magyarád	1	157
	Tiszafüred-Ásotthalom	Hatvan/Füzesabony	Unknown	1650
	Szihalom-Földvár	Hatvan/Füzesabony	1	16.5
	Jászdózsa-Kápolnahalom	Füzesabony	Unknown	168.5
	Felsődobsza-Várdomb	Füzesabony	Unknown	112.5
	Ároktő-Dongóhalom	Füzesabony	1	275.5

	Site	Archaeological group	Number of samples	Total number of remains
Early Bronze Age (n=12)	Budapest, Corvin tér	Unknown	2	66
	Pécs-Nagyárpád	Somogyvár-Vinkovci	9	422.5
	Kakucs-Turján	Nagyrev	28	3676
	Mosonszentmiklós-Pálmajor	Makó-Kosihy-Čaka	6	19
	Kiskundorozsma (M5 45, 26/59)	Makó-Kosihy-Čaka	27	1
	Endrőd 161	Makó-Kosihy-Čaka	6	87.5
	Biatorbágy-Hosszúrét	Makó-Kosihy-Čaka	4	34
	Szigetszentmiklós-Vízmű	Bell Beaker	10	88
	Dunakeszi-Székesdűlő (w)	Bell Beaker	16	372
	Budapest, Csepel-Hollandi u.	Bell Beaker	6	132
	Budapest, Albertfalva - Hunyadi J. U	Bell Beaker	37	2700
	Budakalász M0 motorway, 12	Bell Beaker	51	69




Figure 1. Distribution of the Bronze Age settlements in Hungary included in the current overview. The majority of the sites are plotted based on estimations of their locations because the exact coordinates of the sites are not available to the author. **Early Bronze Age (yellow):** 1. Budakalász M0 motorway, 12; 2. Budapest, Albertfalva-Hunyadi J. U; 3. Budapest, Csepel-Hollandi u.; 4. Dunakeszi-Székesdűlő; 5. Szigetszentmiklós-Vízmű; 6. Biatorbágy-Hosszúrét; 7. Endrőd 161; 8. Kiskundorozsma (M5 45, 26/59); 9. Mosonszentmiklós-Pálmajor; 11. Pécs-Nagyárpád; 12. Budapest, Corvin tér. **Early and Middle Bronze Age (green):** 10/31. Kakucs-Turján. **Middle Bronze Age (blue):** 13. Ároktő-Dongóhalom; 14. Felsődobsza-Várdomb; 15. Jászdózsza-Kápolnahalom; 16. Szihalom-Földvár; 17. Tiszafüred-Ásotthalom; 18. Süttő-Hosszúvölgy; 19. Budapest, Csepel-Szennyvíz telep; 20. Tószeg-Laposhalom; 21. Baracs-Bottyánsánc; 22. Bölske-Vörösgyír; 23. Dunaújváros-Kosziderpadlás; 24. Békés-Várdomb; 25. Tarhos Gyepesi Átkelő (Tarhos 26); 26. Túrkeve-Terehalom; 27. Ménfőcsanak-Szeles; 28. Budapest, Bocskai-Fehérvári úti aluljáró; 29. Cegléd 4/1; 30. Dömsöd-Apaj; 32. Mende-Leányvár; 33. Pákozd-Várhegy; 34. Solymár-Várhegy; 36. Tiszaalpár-Várdomb; 37. Balatonboglár-Szárszó. **Middle and Late Bronze Age (white):** 35/50. Százhalombatta-Földvár. **Late Bronze Age (red):** 38. Poroszló-Aponhát; 39. Ludas, Varjú-dűlő; 40. Balatonmagyaród-Hídvépuszta; 41. Börcs-Paphomlok; 42. Dunakeszi-Székesdűlő (Auchan); 43. Isztimér-Csőszpuszta; 44/45. Mosonmagyaróvár-Németbánya; 46. Polgár site 31; 47. Budapest, Albertfalva-Kitérő út; 48. Gőr-Kápolnadomb; 49. Sopron-Krautacker site 1; 51. Győr-Szabadrétadomb; 52. Lébény-Billedomb (Basemap: Stamen Design; Map data: Open Street Map 2018).

The plant spectra

For practical reasons, the crops and gathered plants have been summarised per phase and archaeological group in Table 7. As this does not provide an insight into the diversity of plants at individual settlements, the absolute quantities and relative frequencies of taxa have been presented according to plant groups per settlement in Figures 2 and 3. Cereals prevail among the macro-botanical finds from the Early Bronze Age, while other plant groups occur in modest amounts. The most common cereal taxa vary per archaeological group: barley was most prominent at Bell Beaker and Makó-Kosihy-Čaka group sites, while einkorn was of greater importance at Nagyrév and Somogyvár-Vinkovci group sites. The absolute quantities and relative frequencies of taxa¹ show a relatively uniform picture, with some expected variations in the frequencies of taxa per site. It is noteworthy that there is a great diversity of weeds and wild plants (see Table 7), which can be seen as an indication of the presence of products and by-products of harvested crops. Grains of millet have been identified in small amounts in two Bell Beaker settlements and one Makó-Kosihy-Čaka settlement.

Middle Bronze Age macro-botanical assemblages show a greater diversity compared with the previous phase. Cereals prevail among most of the groups distinguished archaeologically, followed by pulses, oil and fibre crops, and gathered plants. The most important taxa represented in different archaeological groups are either einkorn or barley among the cereals, lentil (*Lens culinaris*) and bitter vetch (*Vicia ervilia*) of the pulses, and safflower (*Carthamus tinctorius*) and flax/linseed (*Linum usitatissimum*) among oil and fibre crops. There are, however, several exceptions. At first glance, one of these seem to be sites of the Otomani communities, where gathered plants make up the largest proportion of finds. This can be attributed to the settlement site of Túrkeve-Terehalom, where an unusually large amount of cleaned white goosefoot (*Chenopodium album*) fruits has been interpreted as gathered and consumed by the inhabitants (Gyulai 2010, 104). Another exception are some Hatvan/Füzesabony settlements, where the only macro-botanical finds were millet grains. Lastly, several settlements include a larger proportion of wild than cultivated plants (Ároktő-Dongóhalom, Borsod-Abaúj-Zemplén County; Budapest, Csepel-Szennyvíztelep telep, Pest County; and Cegléd 4/1, Pest County), while others include a larger proportion of pulses than cereals (Kakucs-Turján and Pákozd-Várhegy, Fejér County).

Similar to the plant spectra of the Middle Bronze Age, those of the Late Bronze Age are also varied. Cereals are the most common finds at the majority of the settlements, and whereas during the previous phases the cereal finds include large proportions of barley and einkorn, during the Late Bronze Age the most important cereals were spelt (*Triticum aestivum* ssp. *spelta*), emmer (*Triticum turgidum* ssp. *dicoccon*) and millet. Millet prevails in the Gáva and the three Tumulus community settlements; the plant spectra of the Tumulus groups also include a variety of pulses and a large proportion of wild plants and weeds. The latter is especially a reflection of the large assemblages of waterlogged remains of wild plants and weeds from the Tumulus community settlements of Dunakeszi-Székesdűlő (Auchan; Pest County) and Mosonmagyaróvár-Németdűlő (Győr-Moson-Sopron County). Except for cultivated millet, wild species of Panicoideae (e.g. *Digitaria ischaemum* and *Echinochloa crus-galli*) are identified in larger quantities, adding to the variety of taxa in Late Bronze Age assemblages.

1 In this paper, frequency of taxa denotes the number of different taxa within each plant category (cereals, pulses, etc.) expressed as percentage of the total number of taxa per site. As context-based data are unavailable to the author, the calculation of taxa frequency expressed as percentage of the total number of samples was not possible.

(Transition to) Early Bronze Age (2800/2700-2000/1900 cal. BCE)											
Period	Archaeological group	Plant part	Scientific name	Bell Beaker		Makó-Kosihy-Čaka		Nagyrév	Somogyvár-Vinkovci	Unknown	
No. of sites				5	4	1	1	1			
No. of samples				120	43	28	9	2			
Preservation	Count	No. of occurring sites	Count	No. of occurring sites	Count	Common name					
Cereals											
		Grain		Charred				92.0		Barley	
		Grain		Charred	1256.0	5.0	68.0	3.0	35.0	5.5	Hulled barley
		Grain		Charred	17.0	4.0					Naked barley
		Rachis fragment		Charred	10.0	1.0		9.0			Barley
		Rachis fragment		Waterlogged							Barley
		Grain		Charred	1.0	1.0					Rye
		Grain		Charred	1.0	1.0				5.5	Club wheat
		Grain		Charred	3.0	2.0	5.0	2.0		10.0	Common wheat
		Rachis fragment		Charred						1.0	Common wheat
		Grain		Charred	1.0	1.0					Spelt
		Spikelet fork		Charred							Spelt
		Grain		Charred							Free-threshing wheat
		Spikelet fork		Charred							Free-threshing wheat
		Grain		Charred	84.0	3.0			848.5	238.0	Einkorn
		Spikelet fork		Charred	593.0	3.0			1992.0		Einkorn
		Grain		Charred							Hulled wheat
		Spikelet fork		Charred	19.0	2.0			2.5		Hulled wheat
		Grain		Charred	101.0	5.0	4.0	1.0	77.0	82.0	Emmer

Table 7 (a, b, c). Overview of the macro-botanical assemblages in Bronze Age Hungary per archaeological group. Compilation of macro-botanical data after Gylai (2010, Appendix Bronze Age data), Duffy (2010, 275, Table 9.1), Stika and Heiss (2013, 79-80, Table 1) and Filatova (2020).

Period	(Transition to) Early Bronze Age (2800/2700-2000/1900 cal. BCE)							
Archaeological group	Bell Beaker	Makó-Kosihy-Čaka		Nagyrev	Somogyvár-Vinkovci	Unknown		
No. of sites	5	4		1	1	1		
No. of samples	120	43		28	9	2		
Scientific name	Plant part	Preservation	Count	No. of occurring sites	Count	No. of occurring sites	Count	Common name
<i>Triticum turgidum</i> ssp. <i>dicocon</i>	Spikelet fork	Charred	126.0	4.0	6.0	2.0		Emmer
<i>Triticum turgidum</i> ssp. <i>dicocon</i>	Spikelet fork	Waterlogged						Emmer
<i>Triticum turgidum</i> ssp. <i>durum</i>	Rachis fragment	Charred						Macaroni wheat
<i>Triticum timopheevii</i>	Spikelet fork	Charred						New Type Glume (NTG) wheat
<i>Triticum</i> sp.	Grain	Charred						Wheat species
<i>Cerealia indet</i>	Grain	Charred	271.0	2.0	25.0	2.0	283.0	Indeterminate cereals
<i>Cerealia indet</i>	Spikelet fork	Charred						Indeterminate cereals
Millet								
<i>Panicum miliaceum</i>	Grain	Charred	9.0	2.0	1.0	1.0		Broomcorn millet
<i>Panicum miliaceum</i>	Grain	Waterlogged						Broomcorn millet
Pulses								
<i>Cicer arietinum</i>	Seed	Charred						Chick-pea
<i>Lathyrus sativus</i>	Seed	Charred						Grass pea
<i>Lens culinaris</i>	Seed	Charred	1.0	1.0			186.0	Lentil
<i>Pisum sativum</i>	Seed	Charred	1.0	1.0	26.5	2.0	5.5	Pea
<i>Vicia ervilia</i>	Seed	Charred					1.0	Bitter vetch
<i>Vicia faba</i>	Seed	Charred	1.0	1.0				Broad bean
<i>Vicia sativa</i>	Seed	Charred						Common vetch
<i>Leguminosae sativae indet</i>	Seed	Charred						Indeterminate pulses

Table 7a. continued.

(Transition to) Early Bronze Age (2800/2700-2000/1900 cal. BCE)									
Period	Bell Beaker		Makó-Kosihy-Čaka		Nagyrev		Somogyvár-Vinkovci		Unknown
Archaeological group									
No. of sites	5		4		1		1		1
No. of samples	120		43		28		9		2
Scientific name	Plant part	Preservation	Count	No. of occurring sites		Count	Count		Common name
Oil, fibre and dye crops									
<i>Camelina sativa</i>	Seed	Charred					5.5		Gold-of-pleasure
<i>Carthamus tinctorius</i>	Achenium	Charred							Safflower
<i>Linum usitatissimum</i>	Seed	Charred				4.0			Flax/linseed
<i>Linum usitatissimum</i>	Seed	Waterlogged							Flax/linseed
<i>Linum usitatissimum</i>	Fruit + perianth	Charred							Flax/linseed
<i>Papaver somniferum</i>	Seed	Charred							Opium poppy
Gathered plants									
<i>Chenopodium album</i>	Fruit	Charred							White goosefoot
<i>Cornus mas</i>	Fruit endocarp	Charred				0.5			Cornelian cherry
<i>Cornus mas</i>	Fruit endocarp	Waterlogged							Cornelian cherry
<i>Corylus avellana</i>	Fruit	Charred				4.0			Common hazel
<i>Crataegus monogyna</i>	Fruit	Waterlogged							Common hawthorn
<i>Fragaria vesca</i>	Fruitlet	Charred							Wild strawberry
<i>Malus silvestris</i>	Seed	Charred							Crab apple
<i>Malus silvestris</i>	Seed	Waterlogged							Crab apple
<i>Malus silvestris</i>	Seed	Mineralized							Crab apple
<i>Malus silvestris</i>	Fruit	Charred							Crab apple
<i>Malus silvestris</i>	Fruit	Waterlogged	8.0	1.0					Crab apple
<i>Portulaca oleracea</i>	Fruit	Charred							Common purslane
<i>Prunus spinosa</i>	Fruit endocarp	Charred							Blackthorn
<i>Prunus</i> sp.	Fruit endocarp	Charred							Prunus species

Table 7a. continued.

(Transition to) Early Bronze Age (2800/2700-2000/1900 cal. BCE)									
Period	Bell Beaker		Makó-Kösihy-Čaka		Nagyrév		Somogyvár-Vinkovci		Unknown
Archaeological group									
No. of sites	5		4		1		1		1
No. of samples	120		43		28		9		2
Scientific name	Plant part	Preservation	Count	No. of occurring sites	Count	No. of occurring sites	Count	Count	Common name
<i>Pyrus</i> sp.	Seed	Charred							Pear
<i>Quercus</i> sp.	Fruitlet	Charred							Oak
<i>Rubus caesius</i>	Fruitlet	Charred							Blackberry
<i>Rubus caesius</i>	Fruitlet	Waterlogged							Blackberry
<i>Rubus</i> sp.	Fruitlet	Charred							
<i>Sambucus ebulus</i>	Fruit endocarp	Charred	1.0	1.0					Danewort
<i>Sambucus ebulus</i>	Fruit endocarp	Waterlogged							Danewort
<i>Sambucus nigra</i>	Fruit endocarp	Charred							Elder
<i>Sambucus nigra</i>	Fruit endocarp	Waterlogged							Elder
<i>Sambucus</i> sp.	Fruit endocarp	Waterlogged							Sambucus species
<i>Valerianella dentata</i>	Fruit	Charred							Narrowfruit consalad
<i>Vitis vinifera</i>	Seed	Charred							Grape vine
<i>Vitis vinifera</i>	Seed	Waterlogged							Grape vine
Panicoideae									
<i>Digitaria ischaemum</i>	Fruit	Charred	4.0	2.0					Smooth crabgrass
<i>Digitaria sanguinalis</i>	Fruit	Charred							Hairy crabgrass
<i>Echinochloa crus-galli</i>	Fruit	Waterlogged							Cockspur
<i>Echinochloa crus-galli</i>	Fruit	Charred	1.0	1.0					Cockspur
<i>Setaria lutescens</i>	Fruit	Charred							Yellow foxtail
<i>Setaria viridis</i>	Fruit	Charred							Yellow foxtail
<i>Setaria viridis</i>	Fruit	Waterlogged							Green foxtail
<i>Setaria</i> sp.	Fruit	Charred							Foxtail grasses
Wild plants and weeds (combined)			843.0	5.0	5.0	3.0	141.5	75.0	1.0

Table 7a. continued.

Scientific name		Plant part	Middle Bronze Age (2000/1900-1500/1450 cal. BCE)										Middle Bronze Age (2000/1900-1500/1450 cal. BCE)										Common name		
			Flúzsabony					Hatvan / Flúzsabony		Magyarád		Nagyrév/ MBA		Nagyrév/ Válya		Otomani		Transdanubian Encrusted Pottery		Vatya		Unknown			
			Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites		Count	No. of occurring sites
				3		>1		2		1		1		1		4		3		1		9		1	
				>1		>1					1					>2		>14		32		424		unknown	
Cereals																									
<i>Hordeum vulgare</i> (undiff.)	Grain	Charred																							Barley
<i>Hordeum vulgare</i> (h)	Grain	Charred	51.0	2.0						1.0		3.0	11671.5	4.0	11.5	3.0	74.0					22521.0	8.0		Hulled barley
<i>Hordeum vulgare</i> (n)	Grain	Charred	5.5	1.0									14300.5	3.0			1.0					566.0	4.0		Naked barley
<i>Hordeum vulgare</i>	Rachis fragment	Charred											2.0	1.0								257.0	2.0		Barley
<i>Hordeum vulgare</i>	Rachis fragment	Waterlogged																							Barley
<i>Secale cereale</i>	Grain	Charred	110.0	1.0													2.0					8.0	1.0		Rye
<i>Triticum aestivum</i> ssp. <i>compactum</i>	Grain	Charred								15.0												1135.0	1.0		Club wheat
<i>Triticum aestivum</i>	Grain	Charred	25.0	2.0									11.0	2.0	49.0	1.0	3.0					186.0	2.0		Common wheat
<i>Triticum aestivum</i>	Rachis fragment	Charred																							Common wheat
<i>Triticum aestivum</i> ssp. <i>spelta</i>	Grain	Charred											33.0	1.0	12.0	1.0	7.0					77.0	1.0		Spelt
<i>Triticum aestivum</i> ssp. <i>spelta</i>	Spikelet fork	Charred																				5.5	2.0		Spelt
<i>Triticum aestivum</i> / <i>durum</i>	Grain	Charred	14.5	1.0																		58.0	2.0		Free-threshing wheat
<i>Triticum aestivum</i> / <i>durum</i>	Spikelet fork	Charred																							Free-threshing wheat
<i>Triticum monoccocum</i>	Grain	Charred	29.5	2.0						141.0			20834.0	3.0	196.5	2.0	55.0					64630.5	6.0	6666.0	Einkorn
<i>Triticum monoccocum</i>	Spikelet fork	Charred											103.0	3.0	20.0	1.0						2730.0	6.0		Einkorn
<i>Triticum monoccocum</i> / <i>dicocum</i>	Grain	Charred																				191.0	2.0		Hulled wheat
<i>Triticum monoccocum</i> / <i>dicocum</i>	Spikelet fork	Charred																				6510.0	2.0		Hulled wheat
<i>Triticum turgidum</i> ssp. <i>dicoccon</i>	Grain	Charred	135.0	2.0							2.0		1414.0	4.0	161.0	2.0	47.0					49899.0	6.0	15385.0	Emmer
<i>Triticum turgidum</i> ssp. <i>dicoccon</i>	Spikelet fork	Charred	5.5	1.0									43.0	1.0	40.0	1.0						210.5	6.0		Emmer
<i>Triticum turgidum</i> ssp. <i>dicoccon</i>	Spikelet fork	Waterlogged									8.0														Emmer
<i>Triticum turgidum</i> ssp. <i>durum</i>	Rachis fragment	Charred																				1.0	1.0		Macaroni wheat

Table 7b.

Period	Middle Bronze Age (2000/1900-1500/1450 cal. BCE)										Middle Bronze Age (2000/1900-1500/1450 cal. BCE)				
	Archaeological group	Fűzesabony	Hatvan / Fűzesabony	Magyarád	Nagyrév/ MBA	Nagyrév/ Válya	Otomani	Transdanubian Enchusted Pottery	Válya	Unknown					
No. of sites		3	2	1	1	4	3	1	9	1					
No. of samples		>1	>1	1	13	>2	>14	32	424	unknown					
Scientific name	Plant part	Count	No. of occurring sites	Count	Count	Count	No. of occurring sites	Count	Count	No. of occurring sites	Count	Count	Count	Count	Common name
<i>Triticum timopheevii</i>	Spikelet fork								249.0	1.0					New Type Glume (NTG) wheat
<i>Triticum</i> sp.	Grain														Wheat species
<i>Cerealia indet.</i>	Grain			550.0	8.0	2261.0	2.0	1.0	11792.0	4.0					Indeterminate cereals
<i>Cerealia indet.</i>	Spikelet fork						47.0	2.0	235.0	1.0					Indeterminate cereals
Millet															
<i>Panicum miliaceum</i>	Grain			555.5		8.0	2.0		60.0	2.0					Broomcorn millet
<i>Panicum miliaceum</i>	Grain														Broomcorn millet
Pulses															
<i>Cicer orientum</i>	Seed					5.5	1.0								Chick-pea
<i>Lathyrus sativus</i>	Seed					5.5	1.0		4677.5	2.0					Grass pea
<i>Lens culinaris</i>	Seed	5.5	1.0		1.0	4706.5	3.0	53.0	46607.5	5.0					Lentil
<i>Pisum sativum</i>	Seed					16.5	2.0	103.0	9404.5	5.0					Pea
<i>Vicia ervilla</i>	Seed					537.0	2.0	220.0	469.5	3.0					Bitter vetch
<i>Vicia faba</i>	Seed					19.0	1.0		123.0	3.0					Broad bean
<i>Vicia sativa</i>	Seed					4.0	1.0		5.5	1.0					Common vetch
<i>Leguminosae sativae indet.</i>	Seed								128.5	1.0					Indeterminate pulses
Oil, fibre and dye crops															
<i>Camelina sativa</i>	Seed								11.0	1.0					Gold-of-pleasure
<i>Carthamus tinctorius</i>	Achenium							1246.0		1.0					Safflower
<i>Linum usitatissimum</i>	Seed								31.0	3.0					Flax/linseed
<i>Linum usitatissimum</i>	Seed								3.0	1.0					Flax/linseed
<i>Linum usitatissimum</i>	Fruit + perianth								1.0	1.0					Flax/linseed
<i>Papaver somniferum</i>	Seed														Opium poppy
Gathered plants															
<i>Chenopodium album</i>	Fruit										36609.0	1.0			White goosefoot
<i>Cornus mas</i>	Fruit endocarp	12.0	1.0			2.0	1.0	2.0	46.0	2.0					Cornelian cherry
<i>Cornus mas</i>	Fruit endocarp														Cornelian cherry
<i>Corylus avellana</i>	Fruit								35.0	1.0					Common hazel
<i>Crataegus monogyna</i>	Fruit					1.0	1.0								Common hawthorn

Table 7b (continued).

Period	Middle Bronze Age (2000/1900-1500/1450 cal. BCE)										Middle Bronze Age (2000/1900-1500/1450 cal. BCE)									
	Archaeologi- cal group	Fűzesabony	Hatvan / Fűzesabony		Magyarád	Nagyrév/ MBA	Nagyrév/ Vátya		Otomani		Transdanubian Encrusted Pottery	Vátya	Unknown	Common name						
			No. of occurring sites	Count			No. of occurring sites	Count	No. of occurring sites	Count					No. of occurring sites	Count				
No. of sites		3	2	1	1	4	3	1	9	1	unknown									
No. of samples		>1	>1	1	13	>2	>14	32	424											
Scientific name	Plant part	Preservation	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites	Count							
<i>Fragaria vesca</i>	Fruitlet	Charred									4.0	1.0		Wild strawberry						
<i>Malus silvestris</i>	Seed	Charred									6.0	1.0		Crab apple						
<i>Malus silvestris</i>	Seed	Waterlogged												Crab apple						
<i>Malus silvestris</i>	Seed	Mineralized									1.0			Crab apple						
<i>Malus silvestris</i>	Fruit	Charred												Crab apple						
<i>Malus silvestris</i>	Fruit	Waterlogged									1.0			Crab apple						
<i>Portulaca oleracea</i>	Fruit	Charred										124.0	1.0	Common purslane						
<i>Prunus spinosa</i>	Fruit endocarp	Charred				5.0	1.0				1.0	7.0	2.0	1.0	Blackthorn					
<i>Prunus</i> sp.	Fruit endocarp	Charred										8.0	1.0		Prunus species					
<i>Pyrus</i> sp.	Seed	Charred				5.5	1.0								Pear					
<i>Quercus</i> sp.	Fruitlet	Charred				1.0	1.0	5.5	1.0	62.0	8.0	2.0			Oak					
<i>Rubus caesius</i>	Fruitlet	Charred										1.0	1.0		Blackberry					
<i>Rubus caesius</i>	Fruitlet	Waterlogged													Blackberry					
<i>Rubus</i> sp.	Fruitlet	Charred	3.0	1.0								17.0	2.0							
<i>Sambucus ebulus</i>	Fruit endocarp	Charred	1.0	1.0								11.0	2.0		Danewort					
<i>Sambucus ebulus</i>	Fruit endocarp	Waterlogged									24.0				Danewort					
<i>Sambucus nigra</i>	Fruit endocarp	Charred										4.0	2.0		Elder					
<i>Sambucus nigra</i>	Fruit endocarp	Waterlogged													Elder					
<i>Sambucus</i> sp.	Fruit endocarp	Waterlogged													Sambucus species					
<i>Valerianella dentata</i>	Fruit	Charred										25.0	1.0		Narrowfruit consalad					
<i>Vitis vinifera</i>	Seed	Charred										2.0	1.0		Grape vine					
<i>Vitis vinifera</i>	Seed	Waterlogged													Grape vine					
Panicoidae																				
<i>Digitaria ischaemum</i>	Fruit	Charred										1.0	1.0		Smooth crabgrass					
<i>Digitaria sanguinalis</i>	Fruit	Charred													Hairy crabgrass					
<i>Echinochloa crus-galli</i>	Fruit	Waterlogged													Cocksbur					
<i>Echinochloa crus-galli</i>	Fruit	Charred													Cocksbur					
<i>Setaria kteszens</i>	Fruit	Charred													Yellow foxtail					
<i>Setaria viridis</i>	Fruit	Charred										2.0	1.0		Yellow foxtail					
<i>Setaria viridis</i>	Fruit	Waterlogged													Green foxtail					
<i>Setaria</i> sp.	Fruit	Charred	159.0	1.0	5.5	52.0	1196.5	4.0	36688.5	2.0	15.0	2658.5	8.0	26.0	Foxtail grasses					
Wild plants and weeds (combined)																				

Table 7b (continued).

Period		Late Bronze Age (1500/1450-800/750 cal. BC)				Late Bronze Age (1500/1450-800/750 cal. BC)				Common name						
Archaeological group	No. of sites	No. of samples	Preservation	Scientific name	Plant part	Count	Kyjatice	Tumulus			Urnfield		Unknown			
								Count	No. of occurring sites		Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites
			</													

Table 7c.

Period		Late Bronze Age (1500/1450-800/750 cal. BC)				Late Bronze Age (1500/1450-800/750 cal. BC)						
Scientific name	Plant part	Preservation	Gáva	Kyjatice	Tumulus	Urnfield		Unknown				
			No. of sites	1	6	4		3				
			No. of samples	2	231	>53	23		7			
			Count	Count	No. of occur- ring sites	Count	No. of occurring sites	Count	No. of occurring sites			
<i>Triticum timopheevii</i>	Spikelet fork	Charred					32.0	1.0				New Type Glume (NTG) wheat
<i>Triticum</i> sp.	Grain	Charred										Wheat species
<i>Cerealia indet</i>	Grain	Charred		6017.0	25.0	2.0	2608.0	3.0	4.0	1.0		Indeterminate cereals
<i>Cerealia indet</i>	Spikelet fork	Charred			1.0	1.0						Indeterminate cereals
Millet												
<i>Panicum miliaceum</i>	Grain	Charred	4000.0	775.0	811.0	5.0	291.0	3.0	784.0	2.0		Broomcorn millet
<i>Panicum miliaceum</i>	Grain	Waterlogged			67.0	2.0						Broomcorn millet
Pulses												
<i>Cicer arietinum</i>	Seed	Charred										Chick-pea
<i>Lathyrus sativus</i>	Seed	Charred			37.0	1.0						Grass pea
<i>Lens culinaris</i>	Seed	Charred	40.0	32.0	9.0	3.0	61.0	4.0				Lentil
<i>Pisum sativum</i>	Seed	Charred		134.0	678.0	1.0	9.0	2.0				Pea
<i>Vicia ervilia</i>	Seed	Charred		990.0	261.0	1.0						Bitter vetch
<i>Vicia faba</i>	Seed	Charred		2.0								Broad bean
<i>Vicia sativa</i>	Seed	Charred										Common vetch
<i>Leguminosae sativae indet</i>	Seed	Charred					2.0	1.0				Indeterminate pulses
Oil, fibre and dye crops												
<i>Camelina sativa</i>	Seed	Charred					359.0	1.0				Gold-of-pleasure
<i>Carthamus tinctorius</i>	Achenium	Charred										Safflower
<i>Linum usitatissimum</i>	Seed	Charred										Flax/linseed
<i>Linum usitatissimum</i>	Seed	Waterlogged										Flax/linseed
<i>Linum usitatissimum</i>	Fruit + perianth	Charred										Flax/linseed

Table 7c (continued).

Period		Late Bronze Age (1500/1450-800/750 cal. BC)				Late Bronze Age (1500/1450-800/750 cal. BC)						
Scientific name	Plant part	Preservation	Count	Kyjatice	Tumulus		Urnfield		Unknown		Common name	
					Gáva	Count	No. of occur- ring sites	Count	No. of occurring sites	Count		No. of occurring sites
No. of sites	1	1	6	4	3							
No. of samples	2	231	>53	23	7							

Table 7c (continued).

Scientific name	Plant part	Period	Late Bronze Age (1500/1450-800/750 cal. BC)					Late Bronze Age (1500/1450-800/750 cal. BC)				
			Gáva		Kyjatice		Tumulus		Urnfield		Unknown	
			Count	Count	Count	Count	Count	No. of occurring sites	Count	No. of occurring sites	Count	No. of occurring sites
<i>Sambucus ebulus</i>	Fruit endocarp	Waterlogged			28.0		2.0					
<i>Sambucus nigra</i>	Fruit endocarp	Charred		2.0								
<i>Sambucus nigra</i>	Fruit endocarp	Waterlogged						1.0	2.0	1.0		
<i>Sambucus</i> sp.	Fruit endocarp	Waterlogged			30.0		1.0					
<i>Valerianella dentata</i>	Fruit	Charred										
<i>Vitis vinifera</i>	Seed	Charred		2.0								
<i>Vitis vinifera</i>	Seed	Waterlogged			5.0		2.0	1.0	1.0	1.0		
Panicoidae												
<i>Digitaria ischaemum</i>	Fruit	Charred		2.0	3.0		2.0				22.0	1.0
<i>Digitaria sanguinalis</i>	Fruit	Charred		24.0					1.0	1.0	2.0	1.0
<i>Echinochloa crus-galli</i>	Fruit	Waterlogged			2.0		1.0					
<i>Echinochloa crus-galli</i>	Fruit	Charred		26.0	1.0		1.0		2.0	1.0		
<i>Setaria lutescens</i>	Fruit	Charred		2.0					41.0	1.0		
<i>Setaria viridis</i>	Fruit	Charred										
<i>Setaria viridis</i>	Fruit	Waterlogged							1.0	1.0		
<i>Setaria</i> sp.	Fruit	Charred		26.0	4.0		2.0		2.0	1.0		
Wild plants and weeds (combined)			249.0	403.0	3160.0	6.0	1772.0	3.0	65.0	3.0		

Table 7c (continued).




Figure 2. Absolute quantities (%) of plant groups plotted per phase, archaeological group and archaeological site.




Figure 3. Frequencies of taxa (%) plotted per phase, archaeological group and archaeological site.

Considerations of the introduction of millet in Hungary

Within the Bronze Age as a whole, a diversity has been observed in the macro-botanical assemblages of all three phases, with site-specific variations in the predominant plant groups and taxa. The most important observations with respect to changes in the assemblages concern the diversification of the plant spectra from the Middle Bronze Age onwards and the shift in the major cereal taxa during the Late Bronze Age, including the more prominent role of millet. The pollen record supports the intensification in agricultural activity, showing increased signs of human disturbance from the Middle Bronze Age onwards. It must be noted that at least part of the (increasing) diversity in the assemblages reflects differences in preservation (*i.e.* charred vs. waterlogged), sampling strategies, and the number of samples and settlements that have been studied per phase and archaeological group.

According to the published data (Figs. 2 and 3, Table 7), there are finds of millet from all three phases of the Bronze Age. However, the results of the radiocarbon dating of individual charred millet grains demonstrate that the earliest of these grains originate from the mid-15th century BCE and thus from the onset of the Late Bronze Age in Hungary (Filipović *et al.* 2020, 6). For the remainder of the early finds of millet in the Bronze Age, it is likely that these also originate from this later phase, as has been demonstrated for the entire Carpathian Basin by Filipović *et al.* (2020).

In general, macro-botanical assemblages from Hungary confirm the introduction of millet as a staple crop during the Late Bronze Age, as the crop then occurs in higher frequency and greater quantity than before and is the predominant cereal taxon at several settlement sites. It is interesting to note the related increase in the frequency and quantity of weedy species of *Panicoideae*, which have been observed to co-occur with grains of millet in several macro-botanical studies (*e.g.* Motuzaitė-Matuzeviciute *et al.* 2012; Stobbe *et al.* 2019). These weedy species have been suggested to have been a source of nutrition prior to the introduction of cultivated millet into Europe (Kroll 2017).

The earliest definite indications for cultivation of millet in Hungary seem to be related to the occurrence of the Tumulus community during the first part of the Late Bronze Age. As described above, Santa (2010, 522) suggested that the more dispersed and generally smaller settlements of the Tumulus community could possibly be related to a shift from the intensive agricultural practices of the Middle Bronze Age (*i.e.* high input of labour within a restricted plot of land located close to the settlement; Bogaard 2005, 180; Jones 2005, 165; Van der Veen 2005, 159) to extensive agricultural practices (*i.e.* low input of labour on large fields located farther from the settlements; Bogaard 2005, 179; Van der Veen 2005, 159) and a greater reliance on animal husbandry. This hypothesis is especially interesting because it proposes a relationship between Tumulus groups, millet cultivation and animal husbandry, a characteristic that is also attributed to central Asian pastoralists who practised low-investment millet cultivation and are believed to have contributed to the spread of the crop from East Asia into mainland Europe (Spengler 2019, 74), although Tumulus groups migrated from the west to the east of Europe (*e.g.* Csányi 2003). Defining cultivation intensity, however, requires a more detailed set of data than has been presented in the current contribution (*e.g.* analysis of individual samples based on weed assemblages; Charles and Jones 1997; Charles *et al.* 2002) and, to my knowledge, no macro-botanical studies of Late Bronze Age assemblages from Hungary have attempted such a reconstruction. Furthermore, as has also been emphasised by Bartosiewicz (this volume) for the zooarchaeological remains, there exists a diversity in the plant spectra and this diversity seems to be more strongly related to local factors than to 'generalised cultural habits'. Although all Tumulus group settlements in Hungary contain finds of cultivated millet, it does not prevail

at every settlement; at the Tumulus sites of Isztimér-Csőszpuszta (Fejér County), Mosonmagyaróvár-Németdűlő and Polgár 31 (Hajdú-Bihar County), other cereals were more prominent than millet (Fig. 2). This topic requires the support of more systematic studies; the suggestion of Sánta (2010) remains an interesting hypothesis until those become available.

The environmental records obtained from stable isotopes in cave stalagmites and bivalve shells in western and southern Hungary indicate that the humidity peak characterising the Middle Bronze Age gave way to a drier and warmer climate during the Late Bronze Age (Demény *et al.* 2019). Although millet is known to tolerate low amounts of water (Filipović *et al.* 2020), it is, in my opinion, doubtful that its introduction is related to this relatively subtle climatic shift. The climatic deterioration suggested as part of the explanation for the change in settlement patterns at the beginning of the Late Bronze Age (Fischl *et al.* 2013) may be based on the data whose interpretations have changed. Namely, the more recent studies by Demény *et al.* (2019) indicate an opposite situation: the entire Bronze Age is characterised as mild and suitable for arable agriculture (Demény *et al.* 2019, 13, Fig. 9). Climatic conditions may, however, have varied locally in the Carpathian Basin, which today is still known for the mosaic-like character of its climate and geology (Sümegei and Bodor 2000, 84; Nagy-Bodor *et al.* 2000, 121; Mezősi 2017). From the data at hand, it is unclear whether potential local climatic deterioration may have caused the movement of people throughout Hungary and whether this coincided with the introduction of millet cultivation.

Regardless of the exact circumstances in which millet cultivation was introduced, it would certainly have been an advantageous addition to the plant economies of the groups that cultivated the crop, especially due to its tolerance of a variety of growing conditions and its short growing season (Filipović *et al.* 2020, 1). It is beyond the scope of this paper to address whether the cultivation of millet could have favoured the general developments of Late Bronze Age society in Hungary. The potential relationship between the inclusion of millet into the diet, the widespread bronze metallurgy of the Urnfield groups and the distribution of their settlements into marginal areas, such as mountain ranges, may be a topic of interest for future, more elaborate considerations of the macro-botanical and archaeological archives of these groups.

For now, it can be concluded that the macro-botanical assemblages and the pollen records indicate increasing agricultural activity in Hungary from the Middle Bronze Age onwards. The continuity in the main cereals cultivated during the Early and Middle Bronze Age terminated during the Late Bronze Age, when a different set of main cereals appears, including millet. It is unlikely that the appearance of millet is related to unfavourable climatic conditions, especially since the recently studied isotopic records have shown the opposite. The circumstances of the introduction of millet into Hungary and the likely advantages that this had for the Bronze Age groups remains a topic for future investigations.

Acknowledgements

A warm thank-you to Dragana Filipović, Marta Dal Corso and Wiebke Kirleis for organising the conference around this fascinating topic and for inviting me to write this contribution to the proceedings. I am very grateful to Robert Staniuk and Dragana Filipović for providing valuable corrections to the text and to László Bartosiewicz for the friendly and honest discussion about Bronze Age archaeology and environment.

References

- Bogaard, A., 2005. 'Garden agriculture' and the nature of early farming in Europe and the Near East. *World Archaeology*, 37 (2), 177-196. <https://doi.org/10.1080/00438240500094572>
- Bóna, I., 1975. *Die mittlere Bronzezeit Ungarns und ihre südöstlichen Beziehungen*. Budapest: Akadémiai Kiadó.
- Bóna, I., 1992. Bronzezeitliche Tell-Kulturen in Ungarn. In: W. Meier-Arendt, ed., *Bronzezeit in Ungarn: Forschungen in Tell-Siedlungen an Donau in Theiss*. Frankfurt am Main: Museum für Vor- und Frühgeschichte, 9-39.
- Charles, M. and Jones, G., 1997. FIBS in archaeobotany: functional interpretation of weed floras in relation to husbandry practice. *Journal of Archaeological Science*, 24, 1151-1161. <https://doi.org/10.1006/jasc.1997.0194>
- Charles, M., Bogaard, A., Halstead, P., 2002. Towards the archaeobotanical identification of intensive cereal cultivation: present-day ecological investigation in the mountains of Asturias, northwest Spain. *Vegetation History and Archaeobotany*, 11, 113-142. <https://doi.org/10.1007/s003340200015>
- Csányi, M., 2003. The Tumulus culture: invaders from the west. In: Zs. Visy, ed., *Hungarian Archaeology at the Turn of the Millennium*. Budapest: Ministry of National Cultural Heritage Teleki László Foundation, 161-163.
- Demény, A., Kern, Z., Czuppon, G., Németh, A., Schöll-Barna, G., Siklósy, Z., Leél-Össy, S., Cook, G., Serlegi, G., Bajnóczi, B., Sümegi, P., Király, Á., Kiss, V., Kulcsár, G., Bondár, M., 2019. Middle Bronze Age humidity and temperature variations, and societal changes in east-central Europe. *Quaternary International*, 504, 80-95. <https://doi.org/10.1016/j.quaint.2017.11.023>
- Duffy, P., Parditka, G.M., Giblin, J.I., Paja, L., 2019. The problem with tells: lessons learned from absolute dating of Bronze Age mortuary ceramics in Hungary. *Antiquity*, 93 (367), 63-79. <https://doi.org/10.15184/aqy.2018.179>
- Figler, A., Bartosiewicz, L., Fülek, G., Hertelendi, E., 1997. Copper Age settlement and the Danube water system: a case study from north western Hungary. In: J. Chapman and P. Dolukhanov. *Colloquia Pontica 3: Landscapes in Flux: Central and Eastern Europe in Antiquity*. Oxford: Oxbow Books, 209-230.
- Filatova, S., 2020. *Kakucs-Turján III: a Case Study of Macro-Botanical Formation Processes and Plant Economy from the Hungarian Bronze Age*. Bonn: Dr. Rudolf Habelt GmbH.
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Kloof, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H., Stobbe, A., Tolar, T., Wasylukowa, K., Wiethold, W., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Fischl, K., Kiss, V., Kulcsár, G., Szeverényi, V., 2013. Transformations in the Carpathian Basin around 1600 B.C. In: H. Meller, F. Bertemes, H. Bork, R. Risch, eds., *1600 – Cultural Change in the Shadow of the Thera-Eruption?* Tagungen des Landesmuseums für Vorgeschichte Halle (Saale) 9. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 355-370.
- Fischl, K., Kiss, V., Kulcsár, G., Szeverényi, V., 2015. Old and new narratives for Hungary around 2200 BC. In: H. Meller, H.W. Arz, R. Jung, R. Risch, eds., *2200 BC – Ein Klimasturz als Ursache für den Zerfall der Alten Welt? 2200 BC – A climatic breakdown as a cause for the collapse of the Old World?* 7. Mitteldeutscher

- Archäologentag vom 23. bis 26. Oktober 2014 in Halle (Saale). Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 503-524.
- Fülöp, K., 2017. The birth of wells: a Late Bronze Age well from Pusztataskony-Ledence. In: G. Kulcsár, G. Szabó, V. Kiss, G. Vácsi, eds., *State of the Hungarian Bronze Age Research*. Proceedings of the conference held between 17th and 18th of December 2014. Budapest: Institute of Archaeology, Research Centre for the Humanities, Hungarian Academy of Sciences; Institute of Archaeological Sciences, Faculty of Humanities, Eötvös Loránd University; Ősrégészeti Társaság / Prehistoric Society, 309-336.
- Gamarra, B., Howcroft, R., McCall, A., Dani, J., Hajdu, Zs., Gyöngyver Nagy, E., Szabó, L., Domboróczki, L., Pap, I., Raczky, P., Marcsik, A., Zoffmann, Zs., Hajdu, T., Feeny, R.N.M., Pinhasi, R., 2018. 5000 years of dietary variations of prehistoric farmers in the Great Hungarian Plain. *PLoS ONE*, 13 (5), e0197214. <https://doi.org/10.1371/journal.pone.0197214>
- Duffy, P., 2010. *Complexity and Autonomy in Bronze Age Europe: Assessing Cultural Developments in Eastern Hungary*. Doctoral dissertation, University of Michigan.
- Earle, T. and Kolb, M. J., 2010. Regional settlement patterns. In: T. Earle and K. Kristiansen, eds., *Organizing Bronze Age Societies: the Mediterranean, Central Europe and Scandinavia Compared*. Cambridge: Cambridge University Press, 57-86.
- Gogâltan, F., 2008. Fortified Bronze Age tell settlements in the Carpathian Basin: a general overview. In: J. Czebreszuk, S. Kadrow, J. Müller, eds., *Defensive Structures from Central Europe to the Aegean in the 3rd and 2nd Millennia B.C*. Bonn: Dr. Rudolf Habelt GmbH, 39-56.
- Gyulai, F., 1993. *Environment and Agriculture in Bronze Age Hungary*. Budapest: Archaeolingua.
- Gyulai, F., 2010. *Archaeobotany in Hungary: Seed, Fruit, Food and Beverage Remains in the Carpathian Basin from the Neolithic to the Late Middle Ages*. Budapest: Archaeolingua.
- Gyulai, F., 2014. The history of broomcorn millet (*Panicum miliaceum* L.) in the Carpathian-Basin in the mirror of archaeobotanical remains I: From the beginning until the Roman Age. *Columella – Journal of Agricultural and Environmental Sciences*, 1 (1), 29-38. <https://doi.org/10.18380/SZIE.COLUM.2014.1.1.29>
- Horváth, A., 2005. Régészeti kutatás a tervezett csepeli Központi Szennyvíztisztító telep területén [Archaeological investigations on the territory of the planned Central Sewage Plant in Csepel]. In: P. Zsidi, ed., *A BTM Aquincumi Múzeumának ásatásai és leletmentései 2004-ben*. Budapest: Historical Museum, 177-185.
- Hungarian National Museum Archaeology Database, 2020. <https://archeodatabase.hnm.hu/en>
- Jaeger, M., 2016. *Bronze Age Fortified Settlements in Central Europe*. Bonn: Dr. Rudolf Habelt GmbH.
- Jones, G., 2005. Garden cultivation of staple crops and its implications for settlement location and continuity. *World Archaeology*, 37 (2), 164-176. <https://doi.org/10.1080/00438240500094564>
- Juhász, I., 2007a. The pollen sequence from Alsópáhok. In: C. Zatykó, I. Juhász, P. Sümegi, eds., *Environmental Archaeology in Transdanubia*. Budapest: Archaeological Institute of the Hungarian Academy of Sciences, 29-35.
- Juhász, I., 2007b. Comparison and correlation of four pollen sequences from the Little Balaton region (Alsópáhok, Főneyd, Keszthely and Zalavár). In: C. Zatykó, I. Juhász, P. Sümegi, eds., *Environmental Archaeology in Transdanubia*. Budapest: Archaeological Institute of the Hungarian Academy of Sciences, 36-50.
- Juhász, I. and Szegvári, G., 2007. The pollen sequence from Mezőlak. In: C. Zatykó, I. Juhász, P. Sümegi, eds., *Environmental Archaeology in Transdanubia*. Budapest: Archaeological Institute of the Hungarian Academy of Sciences, 316-325.

- Kienlin, T.L., 2012. Patterns of change, or: perceptions deceived? Comments on the interpretation of Late Neolithic and Bronze Age tell settlement in the Carpathian Basin. In: T.L. Kienlin and A. Zimmerman, eds., *Beyond Elites: Alternatives to Hierarchical Systems in Modelling Social Formations*. International Conference at the Ruhr-Universität Bochum, Germany October 22-24 2009. Bonn: Dr. Rudolf Habelt GmbH.
- Kienlin, T.L., 2015. *Bronze Age Tell Communities in Context: an Exploration into Culture, Society, and the Study of European Prehistory – Part 1*. Oxford: Archaeopress.
- Knipl, I. and Sümegi, P., 2012. Life at the interface of two distinct landscapes – relationship of humans and environment in the periphery of the Danube-Tisza interfluvium between Hajós and Császártöltés. *Central European Journal of Geosciences*, 4 (3), 439-447. <https://doi.org/10.2478/s13533-011-0071-x>
- Kordos, L., 1987. Climatic and ecological changes in Hungary during the last 15,000 years. In: M. Pésci and L. Kordos, eds., *Holocene Environment in Hungary: Contribution of the INQUA Hungarian National Committee to the XIIth INQUA Congress*. Budapest: Geographical Research Institute Hungarian Academy of Sciences, 11-24.
- Kroll, H., 2017. Ökologie? – Ökonomie! Zu den Beifunden der neolithischen Getreidewirtschaft. In: J. Lechterbeck, E. Fiscer, eds., *Kontrapunkte: Festschrift für Manfred Rösch*. Bonn: Rudolf Habelt GmbH, 159-163.
- Kulcsár, G., 2009. *The Beginnings of the Bronze Age in the Carpathian Basin: the Makó-Kosihy-Čaka and the Somogyvár-Vinkovci Cultures in Hungary*. Budapest: Archaeolingua.
- Magyari, E.K., Chapman, J.C., Passmore, D.G., Allen, J.R.M., Huntley, J.P., Huntley, B., 2010. Holocene persistence of wooded steppe in the Great Hungarian Plain. *Journal of Biogeography*, 37, 915-935. <https://doi.org/10.1111/j.1365-2699.2009.02261.x>
- Marková, K. and Ilon, G., 2013. Slovakia and Hungary. In: A. Harding and H. Fokkens, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 813-836.
- Matuzaitė-Matuzevičiute, G., Telizhenko, S., Jones, M.K., 2012. Archaeobotanical investigation of two Scythian-Sarmatian period pits in eastern Ukraine: implications for floodplain cereal cultivation. *Journal of Field Archaeology*, 37 (1), 51-60. <https://doi.org/10.1179/0093469011Z.0000000004>
- Mezősi, G., 2017. *The Physical Geography of Hungary*. Cham: Springer International Publishing.
- Nagy-Bodor, E., Járαι-Komlódi, M., Medve, A., 2000. Late glacial and post-glacial pollen records and inferred climatic changes from Lake Balaton and the Great Hungarian Plain. In: M.B. Hart, ed., *Climates: Past and Present*. Geological Society Special Publications 181. London: Geological Society, 121-133.
- Náfrádi, K., Bodor, E., Törőcsik, T., Sümegi, P., 2011. Vegetation history reconstructed from anthracology and pollen analysis at the rescue excavation of the MO Motorway, Hungary. *Central European Journal of Geosciences*, 3 (4), 358-367. <https://doi.org/10.2478/s13533-011-0036-0>
- Poroszlai, I., 2003. Fortified centres along the Danube. In: Zs. Visy, ed., *Hungarian Archaeology at the Turn of the Millennium*. Budapest: Ministry of National Cultural Heritage Teleki László Foundation, 151-155.
- Reményi, L., 2012. The defensive settlements of the Vátya culture and the central European Bronze Age exchange system. In: M. Jaeger, J. Czebreszuk, K. Fischl, eds., *Enclosed Space – Open Society: Contact and Exchange in the Context of Bronze Age Fortified Settlements in Central Europe*. Bonn: Rudolf Habelt GmbH, 275-286.
- Reuter, A., in press. *Einheit in der Vielfalt? Zur Kulturpflanzennutzung im Byzantinischen Reich unter besonderer Berücksichtigung archäobotanischer Untersuchungen in Caričin Grad (Justiniana Prima)*.
- Sánta, G., 2010. Settlements of the Tumulus culture in Hungary. *ANTAEUS*, 31-32, 513-528.

- Somogyi, S., 1987. Relationship between environmental changes and human impact until the 9th century. In: M. Pécsi and L. Kordos, eds., *Holocene Environment in Hungary: Contribution of the INQUA Hungarian National Committee to the XIIth INQUA Congress*. Budapest: Geographical Research Institute Hungarian Academy of Sciences, 25-36.
- Spengler, R.N., III 2019. *Fruits from the Sands: the Silk Road Origins of the Foods We Eat*. California: University of California Press.
- Staniuk, R., 2020. *Kakus-Turján II: Tradition and Practice: Study on Pottery, Chronology and Social Dynamics of the Hungarian Bronze Age*. Studien zur Archäologie in Ostmitteleuropa/ Studia nad Pradziejami Europy Środkowej 19. Bonn: Dr. Rudolf Habelt GmbH.
- Stika, H. and Heiss, A., 2013. Seeds from the fire: charred plant remains from Kristian Kristiansen's excavations in Sweden, Denmark, Hungary and Sicily. In: S. Bergerbrant and S. Sabatini, eds., *Counterpoint: Essays in Archaeology and Heritage Studies in Honour of Professor Kristian Kristiansen*. BAR International Series 2508. Oxford: Archaeopress, 77-86.
- Stobbe, A., Bujskikh, A., Rühl, L., Seuffert, L., 2019. *An Amphora with Millet from the Roman Citadel at Olbia Pontica, Ukraine*. In: J. Meurers-Balke, T. Zerl, R. Gerlach, eds., *Auf dem Holzweg... Eine Würdigung für Ursula Tegtmeier*. Kerpen-Loogh: Deutsche Gesellschaft für Ur- und Frühgeschichte, 251-258.
- Sümege, P. and Bodor, E., 2000. Sedimentological, pollen and geoarchaeological analysis of core sequence at Tököl. In: I. Porozslai and M. Vicze, eds., *SAX: Százhalombatta Archaeological Expedition Annual Report – Field Season 1998*. Budapest: Archaeolingua, 83-90.
- Sümege, P., Gulyás, S., Persaits, G., Szelepcsényi, Z., 2012. Long environment change in forest steppe habitat of the Great Hungarian Plain based on paleoecological data. In: J. Rakonczai and J. Zs. Ladányi, eds., *Review of Climate Change Research Program at the University of Szeget (2010-2012)*. Szeget: Institute of Geography and Geology, 7-24.
- Szabó, G., 2003. The expanding world: masters of bronzeworking in the Carpathian Basin. In: Zs. Visy, ed., *Hungarian Archaeology at the Turn of the Millennium*. Budapest: Ministry of National Cultural Heritage Teleki László Foundation, 163-167.
- Szeverényi, V. and Kulcsár, G., 2012. Middle Bronze Age settlement and society in central Hungary. In: M. Jaeger, J. Czebreszuk, K. Fischl, eds., *Enclosed Space – Open Society: Contact and Exchange in the Context of Bronze Age Fortified Settlements in Central Europe*. Bonn: Rudolf Habelt GmbH, 287-354.
- Van der Veen, M., 2005. Gardens and fields: the intensity and scale of food production. *World Archaeology*, 37 (2), 175-163. <https://doi.org/10.1080/004382405000130731>
- Vicze, M., 2013. Koszider: bread or continuity? In: M. Vicze, I. Porozslai, P. Sümege, eds., *Koszider: Hoard, Phase, Period? Round Table Conference on the Koszider Problem*. Százhalombatta: Matrica Museum, 15-30.

Bronze Age novelties in animal exploitation in the Carpathian Basin in a European context

László Bartosiewicz

Abstract

Bronze Age animal exploitation was very diverse in Europe due to broad chronological boundaries and extensive adaptations to local geographical and socio-economic environments. Demographic changes and the emergence of settlements with high concentrations of increasingly stratified human populations stimulated the diversification of animal-related resources beyond mundane meat consumption. This is directly reflected in ordinary food refuse brought to light at settlement excavations. Subsistence hunting had lost its significance compared with the preceding Neolithic, while domestic animals gained in importance not only as a source of meat: the use of renewable, 'secondary' animal products also became widespread. Innovations such as dairying, wool production and using cattle for draught had already developed earlier. However, the roles of these resources in local economies stabilised during the Bronze Age. Quantitative comparisons between animal remains and macrobotanical finds are made difficult by taphonomic and methodological differences. Reconstructions of mobility, as reflected in archaeozoological assemblages, may be profitably compared with occurrences of millet, a special crop of short growing season and also a C4 plant, detectable by means of stable isotope analysis in both human and animal diets.

Introduction

The study of animal and plant remains began at prehistoric pile-dwellings in Switzerland, where both were preserved in remarkable quantities (Rütimeyer 1861; Heer 1866). Archaeobotany and archaeozoology are disciplines that cross-cut cultural and temporal divisions in archaeology. However, deposition, preservation, sampling, and recovery biases affect plant and animal assemblages differently, making the integrated quantitative analysis of botanical and zoological finds difficult. Raw abundance data cannot be compared directly, methods of standardising joint

datasets are only possible in projects where the finds are collected in a systematically coordinated manner with integrated analyses in mind. Major syntheses are prone to inter-observer bias even in archaeozoology itself. Re-analysing published data is difficult due to uncertain identifications (e.g. distinctions between bones of sheep and goat) and inconsistent reporting of potentially shed antler – a renewable wild animal product – and its unclear relation to hunting (Bartosiewicz 2005, 51). In spite of all these difficulties, however, close cooperation between experts has always produced valuable results (e.g. Gross *et al.* 1990; Schibler *et al.* 1997; VanDerwarker 2010; Jacomet and Schibler 2015; Bleicher *et al.* 2018).

The Bronze Age (BA) in Europe has conventionally been dated to between the mid-3rd and early 1st millennium BCE. If we imagine this interval in terms of CE dates, the challenge in this chapter would be similar to characterising animal keeping ‘during Christianity’, beginning with the fall of the Roman Empire up until today. Although the impacts of technical innovation and religion on daily life should not be confused, the length of this timespan is thought-provoking. A notable difference highlighted by this sketchy analogy is the meagre body of written sources regarding prehistory against which animal remains can be interpreted. Although in southwestern Asia, Egypt, and – exceptionally in Europe – Greece there are contemporaneous texts referring to animals, such evidence is unknown from all other parts of Europe. Therefore, the continent’s BA animal exploitation cannot be appraised in such cognitive terms as linguistic, religious or ‘ethnic’ identities. In spite of the complexity of the Eurasian BA, the focus on millet in this volume makes it easier to single out characteristic phenomena in animal exploitation potentially relevant to the spread of this crop.

Keeping domesticates was firmly established by the BA in most of Europe: stocks could be increased without additional domestication (Bökönyi 1971, 647), the protracted ‘secondary products revolution’ (Sherratt 1983) of inventing sustainable animal products was over. Agriculture also benefited from a valuable secondary product previously neglected in research – manure (Bogaard *et al.* 2013). After 1500 BCE, evidence of manuring is known from northern Europe (De Hingh 2000; Robinson 2003), showing the importance of mixed farming (Wijngaarden-Bakker and Brinkkemper 2005, 496). By the BA, subsistence hunting was abandoned, although it retained some role in meat provisioning in the hilly regions of Austria, Slovakia and western Hungary (Bökönyi 1974, 67; Gál 2017, 64), as well as in the middle Volga River Basin. Here, domestic bovids were introduced along with metallurgy around the turn of the 3rd and 2nd millennia BCE (Matolcsi 1982, 77).

Within this extremely variegated scenario, the title ‘*Novelties in animal exploitation...*’ helps me to focus on a few typical BA developments. There will be an inevitable geographical bias in this brief review. In spite of references to the entire continent and adjacent areas, the data analysed originate from central Europe, especially the Carpathian Basin, where most of my research experience is rooted (Bartosiewicz 2017).

Chronological and environmental framework

There is a south-east-north-west temporal cline in the spread of metallurgy across Europe. The Aegean BA began around 3200 BCE, while at the opposite end of the continent, Oscar Montelius’s (1885) chronology for Scandinavia (supported by radiocarbon dating) indicates the beginning around 1700 BCE. In central Europe, recent radiocarbon results often blurred periods in Reinecke’s typo-chronological system, as each region had its own artefact varieties (Kiss *et al.* 2015; Stockhammer *et al.* 2015), and some old and new ‘cultures’ (defined by stylistic criteria) co-existed, whether for a longer or shorter time (Szabó 2017, 101). Dates proposed for the early (2200-1600 BCE), middle (1700/1600-1300 BCE) and late (1300-800/700 BCE) phases




Figure 1. Graph showing humidity (*Arvicola* %), changes in the altitudes of floodplain occupation (settlement) and AMS-dates for millets during the BA in Hungary. Explanation in the text.

are thus subject to change within a broader geographical context. According to interpretations of the New Kingdom Egyptian texts and other documentary sources (Adams and Cohen 2013), the legendary ‘Sea Peoples’, a little-known confederation of seafaring warriors, ransacked the eastern Mediterranean, contributing to the Late Bronze Age (LBA) collapse (1200-900 BCE) in the region. Meanwhile, this period lasted until approximately 600 BCE in northern Europe.

Climatic oscillations also affected various parts of the continent differently. The BA in central Europe began with a relatively dry period, as observed in the intensive formation of alluvial deposits in Bohemia (Dreslerová 1995). The climate turned cooler and increasingly humid by the middle of the Subboreal period, encompassing the BA in the region (Jäger 1997). The percentage of water vole (*Arvicola amphibius* L., 1758) among rodents in Holocene cave deposits correlates to annual precipitation. Following a lull during the Chalcolithic, this value began to progressively increase around 2200 BCE (Kordos 1977, 225, Fig. 1). Steppe elements of the wild fauna, including Holocene lions, gradually withdrew from the Carpathian Basin (Daróczy-Szabó *et al.* 2020). In Greece, they became extinct during the BA (Thomas 2014).

Topographic surveys show that settlements were moved to high terraces in alluvial areas (Horváth 2000, 150, Fig. 1). Floodplains had been intensively inundated, until alluvial dynamics cut the riverbeds deeper. Consequently, low-lying floodplain areas and emerging islands could be re-occupied toward the LBA. This latter trend, indicated with the orange line in Figure 1, is difficult to quantify relative to the *Arvicola* humidity curve, but the oscillation of hydrological conditions is evident (Horváth 2000, 149). By the time of the LBA, the *Arvicola* humidity curve reached an inflexion point (Kordos 1977, 225). This climatic shift was observed across central Europe (Jäger 1997) and was also seen in other indicators used as a proxy for climatic changes in the northern hemisphere (Gronenborn 2009, 99, Fig. 2).

Gyulai (2014) identified millets at a dozen BA sites in Hungary. Due to their small size, however, these kernels are prone to redeposition, which could be the reason behind their occurrence in ten Neolithic provenances (Gyulai 2014, 32-33). Recently calibrated AMS-dates for millets from Fajsz, Százhalombatta and Pécel point to the end of the MBA-LBA transition in the west of Hungary (1510-1400 cal BCE, $p=95.4\%$; Filipović *et al.* 2020). This is the time of the Tumulus culture, which is seen to have developed locally and then spread from the east toward its heartland in Germany. In any case, millets point to the connections of this culture with the Eurasian steppe, as hypothesised by Anthony (2010, 367).




Figure 2. The percentages of livestock and large game in 66 BA archaeozoological assemblages in Hungary based on the number of identifiable specimens (NISP). Sites are grouped by chronological phase and settlement type.

Animal exploitation in general

Recently, the numbers of identifiable specimens (NISP) in 238 BA archaeozoological assemblages from across Europe were summarised (Bartosiewicz 2013a). Their high variability indicates that surplus became an integral economic factor behind the maintenance of increasingly stratified societies in which power relations had to be constantly negotiated both internally and in relation to the outer world. The emphasis on various domesticates and the versatile modes of their exploitation changed over both time and space, in ways that made use of the different properties of each species to different degrees, in accordance with the variegated needs and aspirations of the given community. Within this immense variability, cattle dominate among domestic artiodactyls (on average 49% of NISP) followed by caprines, *i.e.* sheep and goat (31%), and pig (20%). Sixty-six BA assemblages from Hungary (Bökönyi 1974; Choyke 1983, 2000; Bartosiewicz 1996; Choyke and Bartosiewicz 1999, 2000; Gál 2017) showed a higher average contribution by cattle (58%) and smaller relative frequencies of both sheep/goat (24%) and pig (18%). These simple numbers, however, mask considerable diversity both by period and by settlement type (Fig. 2). High variability within individual settlement types also means that there is no demonstrable correlation between settlement type and taxonomic composition. It seems that local environmental conditions had an impact on both animal keeping and the economic role played by hunting. What stands out in this chaotic-looking representation is the general importance of cattle, the role of horse in Early Bronze Age (EBA) meat provisioning and the capricious contribution of large game. Although caprine percentages usually exceed those of pig, great variability is evident in this regard as well. Small altitudinal differences and diverse, mosaic-like habitats in the Carpathian Basin did not limit the keeping of any species: animal husbandry could flexibly adapt to the immediate environment. This is illustrated by the aforementioned high diversity of taxonomic composition seen within settlement types.

The importance of beef in human diets becomes even more evident when bone weights, rather than NISP values, are compared. Cattle remains exceeded 75% of the weight of all bones among several EBA strata of the Jászdózsa-Kápolnahalom tell settlement, Hungary (Bartosiewicz 2016, 306, Fig. 1). In spite of fragmentation, the heavy bones of large ungulates indicate a larger share in meat supplies than their NISP values would indicate. Increased phenotypic diversity in BA cattle shows regional patterns (Bökönyi 1974; Matolcsi 1982, 122; Bopp-Ito *et al.* 2018, 4). The patterns may also be related to various forms of exploitation. Although work-related lesions are fewer in prehistoric cattle than in Roman period and Middle Ages cattle (Bartosiewicz 2006, 261), in northwestern Europe, EBA and MBA plough marks offer evidence of cattle having been used in tillage (Thrane 1990; Tegtmeier 1993).

Horse

BA animal exploitation shows considerable continuity with the Neolithic. The only notable innovation of strategic impact is the emergence and spread of the domestic horse. The status of Holocene equids in Eurasia, however, has long been debated (Duerst 1908; Meadow and Uerpman 1991). Almost 40 Early and Middle Holocene sites yielded horse remains in Austria, Hungary, Romania, Serbia and Slovakia (Vörös 1981, 1994; Bălăşescu *et al.* 2003; Schmitzberger 2009), and some of these are Neolithic. Bökönyi (1971, 643) considered Chalcolithic horses in Hungary domesticated, while Vörös (1994) argued for the presence of wild horses in the Carpathian Basin. Currently, all such European horses are considered wild (Pruvost *et al.* 2011). Németh *et al.* (2016) assume that the wild horse disappeared from the Carpathian Basin when the Yamnaya culture affected much of Europe around 3000 BCE (Allentoft *et al.* 2015).

Horse bones dominated among the food refuse from the 4th millennium BCE onwards in the steppe region, including deposits at Dereivka (Ukraine) and Botai (Kazakhstan). Mare's milk residue shows that at least some horses at Botai were domesticated around 3500 BCE (Outram *et al.* 2009, 1334). Ludwig *et al.* (2009) suggested that the first domestic horses occurred in Europe during the BA, the time when domestic horse remains were also first identified in the Carpathian Basin, along the Danube, with the onset of the Bell Beaker period (Bökönyi 1974, 242). Bell Beaker pottery is found from the western Mediterranean, across western Europe, all the way to Poland. It became widespread around 2750-2500 BCE and disappeared between 2200 and 1800 BCE. However, its distribution area is not contiguous, possibly resulting from a mobile way of life (Schreiber 1973). Genome-wide analyses of 226 humans associated with the Bell Beaker culture (Olalde *et al.* 2018) displayed only limited genetic affinity between individuals from Iberia and those from central Europe, weakening the argument for inter-regional migration. As shown in Figure 2, EBA horses were an important source of meat in Hungary. Broad anatomical representation and patterns of butchery in Bell Beaker assemblages further illustrate this observation. It has been suggested that Iberian native horses share phylogenetic affinities with a single, 3rd millennium BCE specimen from Hungary (Fages *et al.* 2019). Unfortunately, the exact provenance of this find is not specified within the large city of Dunaújváros, where several BA sites were excavated. In the absence of a precise archaeological context, the possible Iberian connection with a presumed late 3rd millennium BCE 'trade centre' in Hungary cannot be interpreted. Whether there was long-distance exchange of horses during the Bell Beaker period, however, remains a valid question.

Horse flesh consumption became sporadic by the MBA. Once people recognised the high value and strategic importance of horses, their meat must have become more of a luxury than a staple. Horses not only revolutionised herding, warfare and trade; they accelerated information exchange in general. Aside from tripling the speed of human movement (even in canter) (Bartosiewicz 2013a, Fig. 18.2), the physical movement of horses was coupled with their unusual mobility within and among societies: they could be traded, rustled, seized as war booty, or presented as high-status gifts, making the study of their circulation unusually complex.

The first signs of horse's special status were discovered in the Catacomb Grave culture of the Lower Volga region (2500-1200 BCE): the skulls of 40 horses were found in a single structured deposit (Anthony 1997). This special treatment, however, does not yet prove the strategic importance of horse. In western Siberia and the adjacent west Asiatic steppe region, the Andronovo culture is a catch-all term for BA groups (ca. 2300-1000 BCE) who kept cattle and caprines as well as horses. Horses buried at Sintashta, near the Ural Mountains in Russia (ca. 2000-1600 BCE), were laid to rest in a quasi-galloping position. In addition, six two-wheeled chariots dated to ca. 1700-1500 BCE were also recovered at this site (Parzinger 2006, 251-257); the Andronovo culture has been tentatively credited with the invention of spoke-wheeled chariots around 2000 BCE (Anthony and Vinogradov 1995). A recent study of two horse skeletons buried with associated cheek-pieces found at Novoil'novskiy, Kazakhstan – located between Sintashta and Botai – suggests that horsemanship was firmly established in the region by the LBA (Chechushkova *et al.* 2020). The 1890-1774 cal BCE date of these finds indicates a shift from chariots to horseback riding toward the turn of the 2nd millennium BCE.

Necrotic deformation resulting from chronic inflammation in the diastema of an MBA horse mandible from Polgár-Kenderföld, Hungary (Fig. 3), is consistent with the excessive use of rope bridle (Bartosiewicz 2013b, 134), which is a crude loop used to control mounting. Among archaeological artefacts, bridle cheek-pieces offer convincing evidence of horseback riding. They are lateral elements of snaffle bits that exert pressure into the horse's cheeks and lips (Brownrigg 2006), which became common in the northern Pontic steppe and then extended toward the west during the mid-2nd millennium (Bökönyi 1953; Mozsolics 1960; Pankovskiy 2016). Horseback




Figure 3. MBA horse mandible oral fragment deformed by traumatic indentations of the diastema (top) and a healthy reference specimen (bottom).

riding played a pivotal role in the Noua-Sabatinivka cultural complex, a tradition seen as an ‘eastern intrusion’ from the northern Pontic region (Bartosiewicz and Gál 2010a; Boroffka 2013, 889).

Horses thus became a formidable means of cross-regional connectivity, associated with BA socio-economic transformation. Controlling horses created superior military might, a platform in confronting foot soldiers, and horses also became an imposing medium for self-representation for millennia to come (Olsen *et al.* 2006). Among the earliest equestrian peoples in written history are the Cimmerians and the Saka/Scythians mentioned in Assyrian records (Deller 1984). They inhabited the steppe north of the Caucasus and the Black Sea around 1000 BCE. Their first mounted troops were formed when the use of a new type of bridle cheek-piece became common between central Europe and the Caucasus (Olsen *et al.* 2006, 157). Their influence west of eastern Europe is unknown; aside from the circulation of luxury artefacts (weapons, horse tackle, jewellery), their westward movements cannot be confirmed (Metzner-Nebelsick 2000, 165).

Sheep, goat and pig

In southwestern Asia, the Late Chalcolithic-EBA transition is marked by a shift to large-scale caprine herding (Clason and Buitenhuis 1998). Urban centres, palaces and temples often developed on tells, foci of power that ruled extensive hinterlands. Mutton had already attained overwhelming importance in the EBA Late Uruk period (3350-3000 BCE) at Arslantepe, Anatolia (Bartosiewicz 2010, 138), a trend even reflected in the food refuse of individual households (Piccione *et al.* 2015, 16). The concomitant decrease in the share of pig remains is remarkable. Reduced diversity in meat diets may indicate declining self-reliance in the face of increasing centralisation: large caprine herds are easier to inventory and tax than pigs, which would have been easier to keep in individual households. In northeastern Anatolia, the continuing shift from pork to mutton may also be attributable to strengthening relations with steppe regions in Transcaucasia around 3000-2800 BCE, culminating in probable intrusions by mobile pastoralists between 2750-2500 BCE (Siracusano and Bartosiewicz 2012, 114, Fig. 1).

By the Iron Age, the effort to centrally control meat resources may have contributed to pork prohibition in the Biblical Lands (Diener and Robkin 1978, 527). Pig was already poorly represented in the LBA (last third of the 2nd millennium BCE) of Area G at Tel Dor, Israel (Bartosiewicz and Lisk 2018, 283, Fig. 27.2), and was completely absent from Iron Age proveniences in Area D2 (Raban-Gerstel *et al.* 2008, 36, Table 2). Purely ecological explanations are of no help in understanding the origin, presence and change in the status of domestic animals. At Tel Dor, the humid climate of the marshy coastal plain would have been ideal for pig herding, also shown by the presence of wild pig among the remains of game (Raban-Gerstel *et al.* 2008, 38). Sheep also became important in the Balkans, but this trend is less pronounced in the rest of Europe: patterning apparent in southwestern Asia tapers off toward the northwest in increasingly different natural and social environments. When domestic artiodactyls are compared with each other in Hungary, the dominance of cattle is evident, approaching 60% in all phases of the BA (Table 1). Caprine remains identifiable to species show that, in Hungary, the number of bones of sheep is five to six times greater than that of goat among the finds in all periods (Bartosiewicz 1999). Thus, the term caprine may be taken as roughly synonymous with sheep within this context.

Although Table 1 shows parameters obtained for all sites, calculations using 27 assemblages containing at least 500 identifiable specimens shows the same trend: the three BA phases overlap in the top quarter of the ternary diagram in Figure 4, and no statistically significant difference is apparent. Among the large assemblages, only the EBA site of Ravaszd-Villibald-domb (Bartosiewicz 1996) and the MBA assemblages from Kaposvár-Road 61 (Gál 2017, 144) and Kakucs-Turján (Biller 2018, 160) stand out due to low percentages of cattle and relatively high contributions of caprine remains. Meanwhile, sheep and goat are only missing in small datasets along the upper right margin of the diagram: extreme percentages are often a reflection of small assemblage size. In general, the great variability among individual assemblages overshadows minor differences among the mean percentages of phases.

The main trend in Table 1 supports the opinion of Bökönyi (1974, 65) that EBA sheep and goats gained ground at the expense of pigs, although this trend slowed down by the LBA in central Europe. He attributed the decelerating increase in caprine remains to climatic change, pointing out that '*pigs like dampness*', whereas sheep prefer dry grassland habitats. Natural preferences by livestock, however, are known to have been frequently disregarded by their owners. Caprine remains dominated (70-80% of NISP) at the Early Neolithic Körös culture sites in marshland habitats of the Great Hungarian Plain (Bartosiewicz 2012, 198, Table 2). Environmental conditions clearly impacted the health of these herds (Bartosiewicz

Phase	n sites			Cattle %		Caprine %		Pig %	
	<500 NISP	≥ 500 NISP	Total	mean	sd	mean	sd	mean	sd
EBA	21	10	31	55	14	28	12	16	8
MBA	13	13	26	59	15	23	16	17	9
LBA	5	4	9	57	9	25	5	18	7
Pooled	39	27	66	56	14	27	14	17	9

Table 1: Mean values and standard deviations (sd) of percentages of artiodactyl NISP values in the 66 Bronze Age assemblages in Hungary shown in Figure 2.




Figure 4. The percentage contribution of domestic artiodactyls to 66 BA assemblages in Hungary. White dots indicate large (NISP≥500) assemblages.

2008). Climatic factors may also have influenced subsequent caprine introductions to Europe (Gronenborn 2009; Grömer and Saliari 2018).

At the tell site of Százhalombatta-Földvár, Hungary, the contribution of sheep rose from below 20% before 2000 BCE to over 40% in subsequent periods (Vretemark 2010). Terramare pile structures in the plains of MBA northern Italy (ca. 1700-1150 BCE) show comparable phenomena (Bernabò-Brea *et al.* 1997). By around 1600/1500-1300 BCE, the contribution of caprines rose to 40% at the Terramare site of Montale (Emilia-Romagna). Studer (1900, 107) noted an increase in BA sheep remains in Switzerland, a trend stronger in the west of the country (Hescheler and Kuhn 1949). Recent studies have shown that the proportion of caprine remains rose from ca. 20% to 30-40% at BA pile-dwellings, at the expense of pig and game. At inland settlements, this increase was 30-70% (Schibler and Studer 1998, 182, Fig. 72). Quantitative increase was accompanied by qualitative changes in sheep populations. By the end of the 3rd millennium BCE, different population types occurred in central Europe, where withers height estimates show considerable regional variation (Bökönyi 1974, 648, Fig. 4; Grömer and Saliari 2018; Schmölcke *et al.* 2018).

A basic form of renewable exploitation shown by longevity is reproduction. Although possibly slaughtered for meat, the lamb is a secondary product of the ewe, the latter potentially having been kept alive until capable of lambing. Reproduction rates differ among domesticates. In contrast to proliferous pigs (four to six piglets per litter), sheep are uniparous. Twin lambing developed only in modern sheep breeds. An effort to maintain and increase flocks thus results in general longevity in females.

Mutton: Proxy for wool production?

According to Leuzinger and Rast-Eicher (2011, 5), an increase in sheep remains may have been a sign of wool production in Switzerland from the Corded Ware culture onwards. Direct evidence, in the form of a small fragment of wool, however, is only available from the BA (Rast-Eicher and Reinhard 1997, 287). Excessive-looking supplies of mutton at BA rural sites in the inner Alpine region (possibly also related to the provisioning of mining sites: Schmölcke *et al.* 2018, 110) clearly illustrate the complex relationships among various sheep products.

Along with regional upsurges in the quantity of caprine remains, age distributions indicative of longevity may support hypotheses concerning renewable animal products, such as wool (Becker *et al.* 2016). At Százhalombatta-Földvár, caprine age distributions indicate a pattern of slaughtering younger individuals before 2000 BCE, whereas the remains of adult and mature individuals dominate in the subsequent Koszider phase of the Vátya culture at the site (1500-1400 BCE) (Vretemark 2010; Vicze 2013). At Kakucs-Turján, on the Great Hungarian Plain, also assigned to the Vátya culture (1900/1800-1700/1600 cal BCE), caprine bones make up half of the 1981 identifiable mammalian bones. However, only 15% of the 747 ageable sheep remains represented fully grown individuals (Biller 2018, 163). This low value is comparable to the BA sheep mortality pattern observed at Arslantepe, where in spite of the massive contribution of mutton to the diet, no marked longevity was observed to support hypotheses concerning specialised wool production (Siracusano and Bartosiewicz 2012, 119-120). According to Ryder (1987, 114), domestic sheep with primitive fleece evolved by about 3000 BCE in southwestern Asia and by 1500 BCE in Europe.

Written sources suggest that wool had an increasing impact on many aspects of life in southwestern Asia. Rare direct evidence of wool manufacturing is preserved only under special circumstances. An early 3rd millennium BCE mixture of sheep wool and camel hair was found in a saline layer at Shahr-I Sokhta, Iran (Compagnoni and Tosi 1978, 97, Fig. 4). In central Europe, remains of woollen textile found in the Hallstatt salt mines, Austria (Grömer 2012), and at Pustopolje, Bosnia and Herzegovina (Marić Baković and Car 2014), were dated to the mid-2nd millennium BCE, the latter being the earliest dyed woollen fabric find in Europe (1520-1510 BCE, 1σ) (Grömer *et al.* 2018, 354). Garments of the salt miners in Hallstatt were nearly all made of wool by between 1460 and 1245 BCE (Grömer 2012; Bender Jørgensen and Rast-Eicher 2016). In spite of the good preservation of organic materials, animal fibres rarely survive in the alkaline deposits at circum-Alpine waterlogged sites (Rast-Eicher and Reinhard 1997). A rare exception is the piece of dyed textile found at the Pfäffikon-Irgenhausen lake dwelling, Switzerland, dated to the EBA-MBA transition (Grömer 2013, 83). In the north, textile finds from the oak-log coffin burials in Denmark (Broholm and Hald 1940; Bergerbrant 2007) date to the mid-2nd millennium BCE as well (Holst *et al.* 2001).

Almost all BA textiles in northern Europe were made of relatively thick wool yarns (Sofaer *et al.* 2013, 479), resulting in coarse and less evenly made fabric, indicative of a different technology to that in regions to the south (Grömer *et al.* 2018, 355). In the absence of finds of wool manufacturing equipment, the question arises of how many of these ca. 1600-1100 BCE wool finds were locally produced. Gleba (2017)

suggested that textile working in the European continent differed from that of the Mediterranean region. Her observations are consonant with the hypothesis that wool processing followed distinct regional trajectories (Sabatini 2018) that may be reflected in the great variability of caprine representation in animal bone assemblages.

Mobile pastoralism and millet

Broomcorn millet, a drought-tolerant crop of short growing season and rapid seeding (Nesbitt 2005; Spengler *et al.* 2014), has long been a crop of choice for mobile pastoralists. Around 1600 BCE, BA herders in Inner Asia cultivated domestic wheat and barley of southwestern Asian extraction, as well as broomcorn and foxtail millet originating from China (Hermes *et al.* 2019, 1). From the BA perspective, classical *topoi* concerning nomads may be anachronistic, but they do refer to millet within the important context of mobile pastoralism. Herodotus ([1987] IV: 10-18) described the 5th century BCE Scythians as ‘*not cultivators but pastoralists*’, who also grew millet. Bovid herders likely accelerated the trans-Eurasian transmission of domestic cereals during the EBA (2500-2300 BCE) while exploiting foothill graze between the Iranian Plateau, the Altai Mountains and western China (Hermes *et al.* 2019, 7), contributing to the westward dispersal of millets. The direct role of horses in this process remains a question. Plinius Maior ([1938] XVIII: XXIV, 100), however, lends weight to their association with millet in 77 CE:

‘Sarmatarum quoque gentes hac maxime pulte aluntur et cruda etiam farina, equino lacte vel sanguine e cruris venis admixto. Aethiopes non aliam frugem quam milii hordeique novere’ [Sarmatian tribes live chiefly on millet porridge, and even on the raw meal, mixed with mare’s milk or with blood taken from the veins in a horse’s leg. Millet and barley are the only grains known to the Ethiopians]. (Rackham 1938, 253).

The coincidental reference to Ethiopia is interesting because East Africa is one of the few places where, until recently, draining blood from live animals has remained a form of secondary exploitation among cattle herders. According to a 13th century CE source (Ligeti 1962), nomadic Mongols also resorted to consuming blood from live camels.

Domestic chicken?

Domestic chicken is a species of East Asian origins, although the northward distribution of its wild ancestor and thus possible places of domestication in Asia are still to be studied (Eda *et al.* 2019, 6). Although a compilation of finds by West and Zhou (1988, 522, Table 2) includes putative Neolithic and Bronze Age specimens from central and southeastern Europe, recently there has been a consensus that these unusually early European occurrences may require verification (Pitt *et al.* 2016, 3). The taphonomic challenge that chicken bones pose is quite similar to that of ‘pre-BA’ millet finds: small and often fragmented bird remains are prone to post-depositional transport between layers. With the exception of in situ finds or perfectly sealed contexts, their dating by means of stratigraphy is unreliable and needs to be supported using direct radiocarbon measurements.

The main period for domestic chicken dispersion throughout Europe was the Iron Age (Benecke 1993), with the earliest reliable find reported from the Czech Republic. The hillfort on top of the Rubín hill yielded a partial skeleton of an adult individual radiocarbon dated to 720-690 BCE (2 σ ; Kyselý 2010, 13). This date is consonant with other discoveries of domestic fowl in central Europe attributed to the Hallstatt period C and D phases (ca. 8th century BCE) and the contemporaneous Scythian finds at the settlements of Jászfelsőszentgyörgy (Bökönyi 1974, 371) and Balassagyarmat (Bartosiewicz and Gál 2010b, 116, Table 9.1), in eastern Hungary.

Although verifiable occurrences of domestic chicken post-date the earliest millet finds in Europe, the ‘Scythian connection’ is indicative of the possibility that these birds reached the European continent from the Eurasian steppe region through mediation by mobile pastoralists, whose earlier waves had plausibly played a role in introducing millet through migration and/or trade.

Thoughts on antler manufacture

Following the Neolithic, the quality of bone manufacture began to deteriorate in central Europe, showing strong regional variability attributable to localised traditions manifested in sometimes opportunistic household production (Choyke and Schibler 2007, 58, Fig. 8). Meanwhile, a limited class of high-quality objects, probably made by part-specialists, emerged. Many were made from red deer antler, in spite of the decline in hunting. At the Terramare sites in northern Italy, almost three quarters of MBA osseous objects were carved from antler, often using metal tools (Provenzano 2001). In the Baltic, where red deer antler was difficult to find, skeletal bone was more carefully turned into LBA spearheads, points, buttons and decorated handles (Luik and Maldre 2007).

Many of the artefacts were evidently made of shed antler: the naturally separated antler rose is still visible on typical MBA antler mattocks, *e.g.* on the Great Hungarian Plain and in the Po River valley, Italy (Choyke 1998a, Fig. 2; Provenzano 2001). Antler gathering, rather than hunting, explains the positive correlation ($r=0.663$) between the relative frequency of manufactured red deer remains and the contribution of this game to the refuse bone assemblages in BA Hungary (Choyke 1984, Table 3). This relationship is far weaker between bone tools and the skeletal remains of livestock among the food remains, where the choice of raw material was more diverse, the manufacturing less planned, and the use of bone frequently opportunistic (Choyke 1998b, 67). The appearance of such mundane bone tools is in stark contrast to sophisticated decorative antler objects, suggesting that the latter may have been manufactured for an emerging elite by artisans possessing specialised skills as well as rare metal tools (Provenzano 2003). Special-purpose antler objects, often parts of complex multi-media artefacts, included harpoons, polished and decorated handles, toggles and pinheads. However, it is horse harness elements, such as bridle cheek-pieces, that are most directly relevant to the spread of horseback riding at the time (Fig. 5).

Due to their function in mobility and the association with elite horsemanship, they were made in part for display, travelled long distances and may have inspired local artisans over broad areas. These objects represent a spectacular intersection among dimensions of raw material, function and relations with animals: before the BA, there were no horses to ride, whereas after the BA, bridles were made of metal; hunting was no longer important, but antler working still flourished.

An odd ‘footnote’

A previously unparalleled type of BA bone object is represented by perforated canid and hare metapodia, usually drilled above the distal articular condyle, in a dorsoplantar direction. They attracted considerable attention in the material from Százhalombatta-Földvár (Choyke 2000, 101). Two such hare metatarsal pendants were found in the EBA Nagyrév culture layer of the tell, while the MBA Vátya culture strata yielded nine such perforated hare metapodia. Two similar pendants were found at the Vátya culture hillfort of Pákozd-Várhegy (Choyke *et al.* 2004, 183, Figs. 8, 18). Vretemark and Sten (2010) contextualised these objects within the skeletal manipulation of dogs at Százhalombatta. Recent finds from the contemporaneous



Figure 5. Examples of MBA red deer antler working from Százhalombatta-Földvár.

Table 2. Perforated small mammal metapodia in burials.

Site	Grave	Century	Metapodium finds	Comment	Source
Tara Su, Kyrgyzstan	18	4th CE	17 small mammal	woman	Кибиров 1959
Kiskőrös-Vágóhid, Hungary	74	7th-8th CE	11 hare	young girl	Bartosiewicz et al. 2014
Alattán-Tulát, Hungary	161	7th-8th CE	7 hare, 7 canid, 5 non-identifiable small mammal	-	Bartosiewicz et al. 2014
Kiszbombor-E, Hungary	17	7th-8th CE	16 hare	woman	Bálint 1975
Dmitrievka, Ukraine	-	8th-9th CE	9 hare	woman	Pletnyova 1967
Csanytelek-Újhalastó, Hungary	15	6th-4th BCE	2 hare, plus rest of skeleton	woman	Galántha 1981

Vatya culture settlement of Kakucs-Turján (Gál 2018, 124-125, Figs. 9-10) have directed comparable attention to hare. They include 22 hare metapodia and only 4 dog metapodia, as well as 2 hare phalanges. Aside from these three settlements, no BA sites in Hungary have yielded such objects. However, some more recent analogies point to a marked central Asian steppe tradition, as seen in several burials (Table 2). Unfortunately, no parallels to these highly personalised grave goods can be expected from BA Hungary, as cremation prevailed at the time.

The 7th-8th CE finds from the Great Hungarian Plain are associated with the Avars, a population of central Asian origins, whereas the 6th-4th BCE was the time of the Scythian influence in the region. Lending weight to the gendered nature of these objects, Pletnyova (1967, 172) also mentions that hare phalanx pendants are rare and are usually found in association with women.

Discussion and conclusions

The BA witnessed manifold environmental and cultural changes in Eurasia. This often-tumultuous period was characterised by major migrations and even replacements of human populations (Allentoft *et al.* 2015). A century ago, Childe (1925) outlined substantial BA connections between the Near East and southeastern Europe. Amidst conflicting relative chronologies, there are striking stylistic similarities between art in the Mycenaean Greece and the Nordic BA (Kristiansen and Suchowska-Ducke 2015, 363-364). However, the question remains as to which similarities are due to the diffusion of ideas, the distribution of objects, or major human migrations. Similarities between localised phenomena may even be purely coincidental. Although these possibilities are not mutually exclusive, it is difficult to correlate patterning in material culture, including domesticates, with the movement of peoples (Allentoft *et al.* 2015).

Human-driven diffusion is undeniable when domesticates start occurring outside the natural distribution areas of their ancestral forms. Mitochondrial DNA and stable isotopic ratios in animal remains indicate that, around 2700 BCE, domestic caprines of southwest Asian origins were winter-foddered with millet in the Dzhungar Mountains of Kazakhstan (Hermes *et al.* 2019). This discovery already represents a fusion between domesticates originating from two different regions. However, from a European perspective, domestic bovids arrived to the continent millennia before the earliest known occurrence of millet, exposing the complexity of the equation.

The emergence of large BA settlements with high concentrations of people stimulated the diversification of animal resources beyond primary meat consumption. In addition to arable land, extensive pastures were needed for provisioning around large settlements. The increase in caprine remains in various parts of Eurasia may be explained differently depending on time and place: as economic policy, the preservation of breeding stocks, wool production, or even interrelated combinations of these possibilities.

Millet first occurred in Hungary slightly later than in Ukraine and at about the same time as in northern Italy (Filipović *et al.* 2020, 10, Figs. 4-5). Because they were the only C4 crop in Europe prior to the introduction of maize from the New World, millets can be safely detected using dietary stable isotope analyses. Tafuri *et al.* (2009, 150) noted enriched $\delta^{13}\text{C}$ values in the bones of both humans and livestock from the MBA-LBA (16th-12th centuries BCE) site of Olmo di Nogara and the EBA-MBA site of Sedegliano, in northern Italy. Charred food crusts on shards from Bruszczewo, Poland (Heron *et al.* 2016), differed between the EBA (2100-1650 cal BCE) and LBA-Early Iron Age (1100-800 cal BCE), consistent with the occurrence of residue from a C4 plant. Although the single millet kernel identified among the macro-botanical remains at Kakucs-Turján, Hungary (Filatova *et al.* 2018, 178, Table 1), was unsuitable for direct

dating, it would fit this process, especially at a settlement where sheep and goat were heavily represented in the meat diet.

Superficial similarities, however, are easier to spot than substantial differences. Hypothetical connections between BA phenomena outlined in this chapter need to be continuously tested using new data and methods to benefit proper scholarship and understanding of this important period. Co-occurrences or even correlations do not necessarily mean causal relationships. This summary is thus less of a synthesis than a review of ideas worth pursuing in future analyses.

Acknowledgements

Grateful thanks are due to Dragana Filipović, Wiebke Kirleis and Marta Dal Corso for having invited this topic and me to the workshop ‘Millet and what else?’ held in Kiel on 27-28 November 2019. Insightful comments and linguistic and editorial improvements by Jelena Bulatović, Suzanne Needs-Howarth and an anonymous reviewer have greatly improved the final manuscript. Erika Gál provided important literature on chicken domestication.

References

- Adams, M.J. and Cohen, M.E., 2013. Appendix. The ‘Sea Peoples’ in primary sources. In: A.E. Killebrew and G. Lehmann, eds., *The Philistines and other ‘Sea Peoples’ in Text and Archaeology*. Atlanta: Society of Biblical Literature, 645-664.
- Allentoft, M.E., Sikora, M., Sjörögen, K.G., Rasmussen, S., Rasmussen, M., Stenderup, J., Damgaard, P.B., Schroeder, H., Ahlström, T., Vinner, L., Malaspinas, A.-S., Margaryan, A., Higham, T., Chivall, D., Lynnerup, N., Harvig, L., Baron, J., Della Casa, P., Dąbrowski, P., Duffy, P.R., Ebel, A.V., Epimakhov, A., Frei, K., Furmanek, M., Gralak, T., Gromov, A., Gronkiewicz, S., Grupe, G., Hajdu, T., Jarysz, R., Hartanovich, V., Khokhlov, A., Kiss, V., Kolář, J., Kriiska, A., Lasak, I., Longhi, C., McGlynn, G., Merkevičius, A., Merkyte, I., Metspalu, M., Mkrtchyan, R., Moiseyev, V., Paja, L., Pálfi, G., Pokutta, D., Pospieszny, Ł., Price, T.D., Saag, L., Sablin, M., Shishlina, N., Smrčka, V., Soenov, V.I., Szeverényi, V., Tóth, G., Trifanova, S.V., Varul, L., Vicze, M., Yepiskoposyan, L., Zhitenev, V., Orlando, L., Sicheritz-Pontén, T., Brunak, S., Nielsen, R., Kristiansen, K., Willerslev, E., 2015. Population genomics of Bronze Age Eurasia. *Nature*, 522, 167-172. <https://doi.org/10.1038/nature14507>
- Anthony, D.W., 1997. Let them eat horses. *Newsletter Institute for Ancient Equestrian Studies*, 4 (Summer).
- Anthony, D.W., 2010. *The Horse, the Wheel, and Language: How Bronze-Age Riders from the Eurasian Steppes Shaped the Modern World*. Princeton: Princeton University Press.
- Anthony, D.W. and Vinogradov, N., 1995. Birth of the Chariot. *Archaeology*, 48 (2), 36-41.
- Bălăşescu, A., Udrescu, M., Radu, V., Popovici, D., 2003. *Archéozoologie en Roumanie: corpus de données*. Musée National d’Histoire de Roumanie, Bibliothèque du Musée National – Série Recherches Pluridisciplinaires 5. Târgoviste: Cetatea de Scaun.
- Bálint, Cs., 1975. A szaltovó-majaki kultúra avar és magyar kapcsolatairól [On the Avar and Hungarian relations of the Saltovo-Mayak culture]. *Archaeologiai Értésítő*, 105, 52-63. <http://real-j.mtak.hu/id/eprint/391>
- Bartosiewicz, L., 1996. Bronzkori állattartás az Északnyugat-Dunántúlon [Bronze Age animal keeping in northwestern Transdanubia, Hungary]. *Acta Musei Papensis*, 6, 31-42.

- Bartosiewicz, L., 1999. The role of sheep versus goat in meat consumption at archaeological sites. In: L. Bartosiewicz and H. Greenfield, eds., *Transhumant Pastoralism in Southern Europe*. Budapest: Archaeolingua, 47-60.
- Bartosiewicz, L., 2005. Plain talk: animals, environment and culture in the Neolithic of the Carpathian Basin and adjacent areas. In: D. Bailey and A. Whittle, eds., *(Un)settling the Neolithic*. Oxford: Oxbow, 51-63.
- Bartosiewicz, L., 2006. Mettre le chariot devant le boeuf: anomalies ostéologiques liées à l'utilisation des boeuf pour la traction. In: P. Pétrequin, R.-M. Arbogast, A.-M. Pétrequin, S. Van Willigen, M. Bailly, eds., *Premiers chariots, premiers araires: la diffusion de la traction animale en Europe pendant les IVe et IIIe millénaires avant notre ère*. CRA Monographies 29. Paris: CNRS Editions, 259-267.
- Bartosiewicz, L., 2008. Environmental stress in early domestic sheep. In: Z. Miklíková and R. Thomas, eds., *Current Research in Animal Palaeopathology*. Proceedings of the Second Animal Palaeopathology Working Group Conference. BAR International Series S1844. Oxford: Archaeopress, 3-13.
- Bartosiewicz, L., 2010. Herding in period VIA: development and changes from period VII. In: M. Frangipane, ed., *Economic Centralisation in Formative States: The Archaeological Reconstruction of the Economic System in 4th Millennium Arslantepe*. Studi di Preistoria Orientale 3. Roma: Sapienza Università, 119-148.
- Bartosiewicz, L., 2012. Mammalian remains from Körös culture sites in Hungary. In: A. Anders and Zs. Siklósi, eds., *The First Neolithic Sites in Central/South-East European Transect. Volume III: the Körös Culture In Eastern Hungary*. British Archaeological Reports International Series S2334. Oxford: BAR Publishing, 195-204.
- Bartosiewicz, L., 2013a. Animals in Bronze Age Europe. In: H. Fokkens and A. Harding, eds., *The Oxford Handbook of Bronze Age Europe*. Oxford: Oxford University Press, 320-339.
- Bartosiewicz, L., 2013b. *Shuffling Nags, Lame Ducks: the Archaeology of Animal Disease*. Oxford: Oxbow Books.
- Bartosiewicz, L., 2016. 'Súlyos állatságok': a hatvani kultúra húsfogyasztása Jászdózsza-Kápolnahalmon ['Weighty matters': meat consumption during the Hatvan culture occupation at Jászdózsza-Kápolnahalom, Hungary]. *Tisicum*, XXV, 303-307.
- Bartosiewicz, L., 2017. Zooarchaeology in the Carpathian Basin and adjacent areas. In: U. Albarella, M. Rizzetto, H. Russ, K. Vickers, S. Viner-Daniels, eds., *The Oxford Handbook of Zooarchaeology*. Oxford: Oxford University Press, 99-112.
- Bartosiewicz, L. and Gál, E., 2010a. Pferde in den Steppen nördlich des Schwarzen Meeres von den frühen Bronzezeit bis ins Mittelalter: Pferde und Nomadentum. In: J. Leskovar and M.Ch. Zingerle, eds., *Goldener Horizont: 4000 Jahre Nomaden der Ukraine*. Linz: Oberösterreichische Landesmuseen, 206-212.
- Bartosiewicz, L. and Gál, E., 2010b. Living on the edge: 'Scythian' and 'Celtic' meat consumption in Iron Age Hungary. In: D. Campana, A.M. Choyke, P. Crabtree, S. deFrance, J. Lev-Tov, eds., *Anthropological Approaches to Zooarchaeology: Colonialism, Complexity and Animal Transformations*. Oxford: Oxbow, 115-127.
- Bartosiewicz, L., Biller, A.Zs., Choyke, A.M., 2014. Középvár nyúlláb függők a Magyar Nemzeti Múzeum gyűjteményéből [Middle Avar period hare foot pendants in the collections of the Hungarian National Museum]. In: A. Anders, Cs. Balogh, A. Türk, eds., *Avarok pusztái. Régészeti tanulmányok Lőrinczy Gábor 60. születésnapjára [Avarum solitudines: archaeological studies presented to Gábor Lőrinczy on his sixtieth birthday]*. Opitz Archaeologica 6 (MTA BTK MÓT Kiadványok 2). Budapest: Martin Opitz, 1-11.
- Bartosiewicz, L. and Lisk, E., 2018. Mammalian remains. In: A. Gilboa, I. Sharon, J.R. Zorn, eds., *Excavations at Dor, Final Report: Volume IIB Area G, the Late Bronze and Iron Age: Pottery, Artifacts, Ecofacts and Other Studies*. Quedem Reports 11. Jerusalem: Israel Exploration Society, 277-312.

- Becker, C., Benecke, N., Grabundžija, A., Küchelmann, H.-C., Pollock, S., Schier, W., Schoch, C., Schrakamp, I., Schütt, B., Schumacher, M., 2016. The textile revolution: research into the origin and spread of wool production between the Near East and central Europe. *ETopoi Journal for Ancient Studies*, Special Volume 6, 102-151.
- Bender Jørgensen, L. and Rast-Eicher, A., 2016. Innovations in European Bronze Age textiles. *Praehistorische Zeitschrift*, 91 (1), 68-102. <https://doi.org/10.1515/pz-2016-0004>
- Benecke, N., 1993. On the utilization of the domestic fowl in central Europe from the Iron Age up to the Middle Ages. *Archaeofauna*, 2, 21-31.
- Bergerbrant, S., 2007. *Bronze Age Identities: Costume, Conflict and Contact in Northern Europe 1600-1300 BC*. Stockholm Studies in Archaeology 43. Lindome: Bricoleur Press.
- Bernabò-Brea, M., Cardarelli, A., Cremaschi, M., eds., 1997. *Le Terramare: la più antica civiltà padana*. Milano: Electra.
- Biller, A. Zs., 2018. Archaeozoological results of the pits from Kakucs-Turján. In: M. Jaeger, G. Kulcsár, N. Taylor, R. Staniuk, eds., *Kakucs-Turján: a Middle Bronze Age Multi-Layered Fortified Settlement in Central Hungary*. Studien zur Archäologie in Ostmitteleuropa Studia nad Pradziejami Europy Środkowej 18. Bonn: Dr. Rudolf Habelt GmbH, 159-174.
- Bleicher, N., Antolín, F., Heiri, O., Häberle, S., Hüster-Plogmann, H., Jacomet, S., Pümpin, C., Rentzel, P., Schäfer, M., Schibler, J., Wiemann, P., van Hardenbroek, M., Tóth, M., Schneider, H.R., 2018. Middens, currents and shorelines: unraveling the complex depositional processes of waterlogged prehistoric lakeside settlements at the example of Zurich-Parkhaus Opéra (Switzerland). *Journal of Archaeological Science*, 97, 26-41. <https://doi.org/10.1016/j.jas.2018.06.010>
- Bogaard, A., Fraser, R., Heaton, T.H.E., Wallace, M., Vaiglova, P., Charles, M., Jones, G., Evershed, R.P., Styring, A.K., Andersen, N.H., Arbogast, R.-M., Bartosiewicz, L., Gardeisen, A., Kanstrup, M., Maier, U., Marinova, E., Ninov, L., Schäfer, M., Stephan, E., 2013. Crop manuring and intensive land management by Europe's first farmers. *Proceedings of the National Academy of Sciences*, 110 (31), 12589-12594. www.pnas.org/cgi/doi/10.1073/pnas.1305918110
- Bopp-Ito, M., Deschler-Erb, S., Vach, W., Schibler, J., 2018. Size diversity in Swiss Bronze Age cattle. *International Journal of Osteoarchaeology*, 28 (3), 1-11. <https://doi.org/10.1002/oa.2654>
- Boroffka, N., 2013. Romania, Moldova, and Bulgaria. In: H. Fokkens and A. Harding, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 877-897.
- Broholm, H.C. and Hald, M., 1940. *Costumes of the Bronze Age in Denmark: Contributions to the Archaeology and Textile-History of the Bronze Age*. Copenhagen: Nyt Nordisk Forlag Arnold Busck.
- Bökönyi, S., 1953. Reconstruction des mors en bois de cerf et en os. *Acta Archaeologica Academiae Scientiarum Hungaricae*, III-1 (4), 113-122.
- Bökönyi, S., 1971. The development and history of domestic animals in Hungary: the Neolithic through the Middle Ages. *American Anthropologist*, 73, 640-674. <https://www.jstor.org/stable/671760>
- Bökönyi, S., 1974. *History of Domestic Mammals in Central and Eastern Europe*. Budapest: Akadémiai Kiadó.
- Brownrigg, G., 2006. Horse control and the bit. In: S.L. Olsen, S. Grant, A.M. Choyke, L. Bartosiewicz, eds., *Horses and Humans: the Evolution of Human Equine Relationships*. BAR International Series 1560. Oxford: Archaeopress, 165-171.
- Chechushkova, I.V., Usmanova, E.R., Kosintsev, P.A., 2020. Early evidence for horse utilization in the Eurasian steppes and the case of the Novoil'inovskiy 2 cemetery in Kazakhstan. *Journal of Archaeological Science: Reports*, 32, 1-10. <https://doi.org/10.1016/j.jasrep.2020.102420>

- Childe, V.G., 1925. *The Dawn of European civilisation*. London: Kegan Paul.
- Choyke, A.M., 1983. *An Analysis of Bone, Antler, and Tooth Tools from Bronze Age Hungary*. Doctoral dissertation, State University of New York, Binghamton.
- Choyke, A.M., 1984. Faunal information offered by worked bone assemblages. *Acta Archaeologica Academiae Scientiarum Hungaricae*, 36, 54-58.
- Choyke, A.M., 1998a. Bronze Age red deer: case studies from the Great Hungarian Plain. In: P. Anreiter, L. Bartosiewicz, E. Jerem, W. Meid, eds., *Man and the Animal World: Studies in Memoriam Sándor Bökönyi*. Budapest: Archaeolingua Kiadó, 157-178.
- Choyke, A. M., 1998b. The bone tool manufacturing continuum. *Anthropozoologica*, 25-26, 65-72.
- Choyke, A.M., 2000. Refuse and modified bone from Százhalombatta-Földvár: some preliminary observations. In: I. Poroszlai and M. Vicze, eds., *SAX: Százhalombatta Archaeological Expedition: Annual Report 1: Field Season 1998*. Százhalombatta: Matrica Museum, 97-102.
- Choyke, A.M. and Bartosiewicz, L., 1999. Bronze Age animal keeping in western Hungary. In: E. Jerem and I. Poroszlai, eds., *Archaeology of the Bronze Age and Iron Age: Experimental Archaeology, Environmental Archaeology, Archaeological Parks*. Budapest: Archaeolingua, 239-249.
- Choyke, A.M. and Bartosiewicz, L., 2000. Bronze Age animal exploitation in the central Great Hungarian Plain. *Acta Archaeologica Academiae Scientiarum Hungaricae*, 51 (1), 43-70.
- Choyke, A.M., Vretemark, M., Sten, S., 2004. Levels of social identity expressed in the refuse and worked bone from Middle Bronze Age Százhalombatta-Földvár, Vátya culture, Hungary. In: S.L. O'Day, W. Van Neer, A. Ervynck, eds., *Behaviour Behind Bones: the Zooarchaeology of Ritual, Religion, Status and Identity*. Oxford: Oxbow, 177-189.
- Choyke, A.M. and Schibler, J., 2007. Prehistoric bone tools and the archaeozoological perspective: research in central Europe. In: C. Gates St-Pierre and R. Walker, eds., *Bones as Tools: Current Methods and Interpretations in Worked Bone Studies*. BAR International Series 1622. Oxford: Archaeopress, 51-65.
- Clason, A.T. and Buitenhuis, H., 1998. Patterns in animal food resources in the Bronze Age in the Orient. In: H. Buitenhuis, L. Bartosiewicz, A.M. Choyke, eds., *Archaeozoology of the Near East III*. Proceedings of the Third International Symposium on the Archaeozoology of Southwestern Asia and Adjacent Areas. Groningen: Centrum voor Archeologische Research and Consultancy, 233-242.
- Compagnoni, B. and Tosi, M., 1978. The camel: its distribution and state of domestication in the Middle East during the 3rd millennium B.C. in the light of finds from Shahr-I Sokhta. In: R.H. Meadow and M.A. Zeder, eds., *Approaches to the Faunal Analysis in the Middle East*. Peabody Museum Bulletin 2. Cambridge, MA: Harvard University Press, 91-103.
- Daróczy-Szabó, M., Kovács, Zs.E., Raczky, P., Bartosiewicz, L., 2020. Pending danger: recent Copper Age lion (*Panthera leo* L., 1758) finds from Hungary. *International Journal of Osteoarchaeology*, 30 (4), 469-481. <https://doi.org/10.1002/oa.2875>
- de Hingh, A.E., 2000. *Food Production and Food Procurement in the Bronze Age and Early Iron Age (2000-500 BC)*. Archaeological Studies Leiden University 7. Leiden: Leiden University.
- Deller, K., 1984. Ausgewählte neuassyrische Briefe betreffend Urartu zur Zeit Sargons II. In: P.E. Pecorella and M. Salvini, eds., *Tra lo Zagros e l'Urmia: ricerche storiche ed archeologiche nell'Azerbaigian Iraniano* [Between Zagros and Urmia: historical and archaeological research in Iranian Azerbaijan]. Incunabula Graeca 78. Roma: Edizione Dell'Ateneo, 97-122.

- Diener, P. and Robkin, E.R., 1978. Ecology, evolution, and the search for cultural origins: the question of Islamic pig prohibition. *Current Anthropology*, 19, 495-540. <https://doi.org/10.1086/202138>
- Dreslerová, D., 1995. The prehistory of the middle Labe (Elbe) floodplain in the light of archaeological finds. *Památky archeologické*, 86, 105-136.
- Duerst, J.U., 1908. Animal remains from the excavations at *Anau* and the horse of *Anau* in its relation to the races of domestic horses. In: R. Pumpelly, ed., *Exploration in Turkestan, Expedition of 1904: Prehistoric Civilizations of Anau, Origins, Growth, and Influence of Environment: Volume 2*. Carnegie Institution of Washington Publication 73. Washington, DC: Carnegie Institution of Washington, 341-431.
- Eda, M., Kikuchi, H., Sun, G., Matsui, A., 2019. Were chickens exploited in the Neolithic early rice cultivation society of the lower Yangtze River? *Archaeological and Anthropological Sciences*, 11, 6423-6430. <https://doi.org/10.1007/s12520-019-00783-x>
- Fages, A., Hanghøj, K., Khan, N., Gaunitz, C., Seguin-Orlando, A., Leonardi, M., McCrory Constantz, C., Gamba, C., Al-Rasheid, K.A.S., Albizuri, S., Alfarhan, A.H., Allentoft, M., Alquraishi, S., Anthony, D., Baimukhanov, N., Barrett, J.H., Bayarsaikhan, J., Benecke, N., Bernáldez-Sánchez, E., Berrocal-Rangel, L., Biglari, F., Boesenkool, S., Boldgiv, B., Brem, G., Brown, D., Burger, J., Crubézy, E., Daugnora, L., Davoudi, H., de Barros Damgaard, P., de los Ángeles de Chorro y de Villa-Ceballos, M., Deschler-Erb, S., Detry, C., Dill, N., do Mar Oom, M., Dohr, A., Ellingvåg, S., Erdenebaatar, D., Fathi, H., Felkel, S., Fernández-Rodríguez, C., García-Viñas, E., Germonpré, M., Granado, J.D., Hallsson, J.H., Hemmer, H., Hofreiter, M., Kasparov, A., Khasanov, M., Khazaeli, R., Kosintsev, P., Kristiansen, K., Kubatbek, T., Kuderna, L., Kuznetsov, P., Laleh, H., Leonard, J.A., Lhuillier, J., Lettow-Vorbeck, C.L. von, Logvin, A., Lõugas, L., Ludwig, A., Luis, C., Arruda, A.M., Marques-Bonet, T., Matoso Silva, R., Merz, V., Mijiddorj, E., Miller, B.K., Mochlov, O., Mohaseb, F.A., Morales, A., Nieto-Espinet, A., Nistelberger, H., Onar, V., Pálsdóttir, A.H., Pitulko, V., Pitskhelauri, K., Pruvost, M., Rajić Šikanjić, P., Rapan Papeša, A., Roslyakova, N., Sardari, A., Sauer, E., Schafberg, R., Scheu, A., Schibler, J., Schlumbaum, A., Serrand, N., Serres-Armero, A., Shapiro, B., Sheikhi Seno, S., Shevnina, I., Shidrang, S., Southon, J., Star, B., Sykes, N., Taheri, K., Taylor, W., Teegen, W.-R., Trbojević Vukićević, T., Trixl, S., Tumen, D., Undrakhbold, S., Usmanova, E., Vahdati, A., Valenzuela-Lamas, S., Viegas, C., Wallner, B., Weinstock, J., Zaibert, V., Clavel, B., Lepetz, S., Mashkour, M., Helgason, A., Stefánsson, K., Barrey, E., Willerslev, E., Outram, A.K., Librado, P., Orlando, L., 2019. Tracking five millennia of horse management with extensive ancient genome time series. *Cell*, 177, 1-17. <https://doi.org/10.1016/j.cell.2019.03.049>
- Filatova, S., Gissel, C., Filipović, D., Kirleis, W., 2018. The plant economy at the Bronze Age site of Kakucs-Turján: first archaeobotanical results. In: M. Jaeger, G. Kulcsár, N. Taylor, R. Staniuk, eds., *Kakucs-Turján: a Middle Bronze Age Multi-Layered Fortified Settlement in Central Hungary*. Studien zur Archäologie in Ostmitteleuropa. Studia nad Pradziejami Europy Środkowej 18. Bonn: Dr. Rudolf Habelt GmbH, 175-190.
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Klooß, S., KohlerSchneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A.M., MuellerBieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H.P., Stobbe, A., Tolar, T., Wasylkowa, K., Wiethold, J., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>

- Galántha, M., 1981. Előzetes jelentés a Csanytelek újhalastói szkítakori temető ásátásáról [Preliminary report on the excavation of the Scythian burial ground of Csanytelek-Újhalastó]. *Communicationes Archaeologicae Hungariae*, 1, 43-58.
- Gál, E., 2017. *Animals at the Dawn of Metallurgy in South-Western Hungary: Relationship between People and Animals in Southern Transdanubia during the Late Copper to Middle Bronze Ages*. Budapest: Archaeolingua.
- Gál, E., 2018. Preliminary results concerning the Middle Bronze Age (Vatya culture) bone tools from Kakucs-Turján, central Hungary. In: M. Jaeger, G. Kulcsár, N. Taylor, R. Staniuk, eds., *Kakucs-Turján: a Middle Bronze Age Multi-Layered Fortified Settlement in Central Hungary*. Studien zur Archäologie in Ostmitteleuropa/Studia nad Pradziejami Europy Środkowej 18. Bonn: Dr. Rudolf Habelt GmbH, 119-136.
- Gleba, M., 2017. Tracing textile cultures of Italy and Greece in the early first millennium BC. *Antiquity*, 91 (351), 1205-1222. <https://doi.org/10.15184/aqy.2017.144>
- Gross, E., Jacomet, S., Schibler, J., 1990. Stand und Ziele der wirtschaftsarchäologischen Forschung an neolithischen Ufer- und Inselsiedlungen im unteren Zürichseeraum (Kt. Zürich, Schweiz). In: J. Schibler, J. Sedlmeier, H.P. Spycher, eds., *Festschrift für Hans R. Stampfli: Beiträge zur Archäozoologie, Archäologie, Anthropologie, Geologie und Paläontologie*. Basel: Helbing & Lichtenhahn, 77-100.
- Grömer, K., 2012. Austria: Bronze and Iron Age. In: M. Gleba and U. Mannering, eds., *Textiles and Textile Production in Europe from Prehistory to AD 400*. Ancient Textile Series 11. Oxford: Oxbow Books, 27-88.
- Grömer, K., 2013. Tradition, Kreativität und Innovation: textiltechnologische Entwicklung von der Bronzezeit zur Hallstattzeit/Tradition, creativity and innovation: the development of textile expertise from the Bronze Age to the Hallstatt period. In: K. Grömer, A. Kern, H. Reschreiter, H. Rösler-Mautendorfer, eds., *Textiles from Hallstatt: Weaving Culture in Bronze Age and Iron Age Salt Mines/ Textilien aus Hallstatt: gewebte Kultur aus dem bronze- und eisenzeitlichen Salzbergwerk*. Budapest: Archaeolingua, 53-98.
- Grömer, K., Jørgensen, L.B., Marić Baković, M., 2018. Missing link: an early wool textile from Pustopolje in Bosnia and Herzegovina. *Antiquity*, 92 (362), 351-367. <https://doi.org/10.15184/aqy.2018.18>
- Grömer, K. and Saliari, K., 2018. Dressing central European prehistory – the sheep's contribution: an interdisciplinary study about archaeological textile finds and archaeozoology. *Annalen des Naturhistorischen Museums in Wien*, 120, 127-156.
- Gronenborn, D., 2009. Climate fluctuations and trajectories to complexity in the Neolithic: towards a theory. *Documenta Praehistorica*, XXXVI, 97-110. <https://doi.org/10.4312/dp.36.5>
- Gyulai, F., 2014. The history of broomcorn millet (*Panicum miliaceum* L.) in the Carpathian Basin in the mirror of archaeobotanical remains from the beginning until the Roman Age. *Columella – Journal of Agricultural and Environmental Sciences*, 1 (1), 29-38. <https://doi.org/10.18380/SZIE.COLUM.2014.1.1.29>
- Hermes, T.R., Frachetti, M.D., Doumani Dupuy, P.N., Mar'yashev, A., Nebel, A., Makarewicz, C.A., 2019. Early integration of pastoralism and millet cultivation in Bronze Age Eurasia. *Proceedings of the Royal Society B: Biological Sciences*, 286, 20191273, 1-9. <https://doi.org/10.1098/rspb.2019.1273>
- Herodotus, 1987. *The History*. Translated by D. Gren. Chicago: University of Chicago Press.
- Heron, C., Shoda, S., Breu Barcons, A., Czebreszuk, J., Eley, Y., Gorton, M., Kirleis, W., Kneisel, J., Lucquin, A., Müller, J., Nishida, Y., Son, J., Craig, O.E., 2016. First molecular and isotopic evidence of millet processing in prehistoric pottery vessels. *Scientific Reports*, 6, 38767. <https://doi.org/10.1038/srep38767>

- Hescheler, K. and Kuhn, E., 1949. *Die Tierwelt der prähistorischen Siedlungen der Schweiz: Sonderdruck aus Otto Tschumi: Urgeschichte der Schweiz, Band I*. Frauenfeld: Verlag Huber & Co AG, 121-368.
- Holst, M.K., Breuning-Madsen, H., Rasmussen, M., 2001. The south Scandinavian barrows with well-preserved oak-log coffins. *Antiquity*, 75 (287), 126-136. <https://doi.org/10.1017/S0003598X00052820>
- Horváth, A., 2000. Hazai újholocén klíma- és környezetváltozások vizsgálata hazai régészeti adatok segítségével [Investigation of middle and late Holocene climatic and environmental changes in Hungary by means of archaeological data]. *Földrajzi Közlemények*, 48, 149-158.
- Jacomet, S. and Schibler, J., 2015. L'alimentation des hommes préhistoriques en Suisse. *Le Globe: Revue genevoise de géographie*, 155, 33-46.
- Jäger, K.D., 1997. Mid- to late Holocene changes in central Europe: climate and man. In: H.N. Dalfes, G. Kukla, H. Weiss, eds., *Third Millennium BC Climate Change and Old World Collapse*. Heidelberg and Berlin: Springer.
- Кибиров, А.К., 1959. Археологические работы в Центральном Тянь-Шане. Труды Киргизской комплексной археолого-этнографической экспедиции 2 [Archaeological work in the central Tien Shan. Proceedings of the Kyrgyz Integrated Archaeological and Ethnographic Expedition 2]. Труды Института языка и литературы Академии наук Киргизской ССР 2. Москва: Наука, 114-116.
- Kiss, V., Fábíán, Sz., Hajdu, T., Köhler, K., Kulcsár, G., Major, I., Szabó, G., 2015. Contributions to the relative and absolute chronology of the Early and Middle Bronze Age in western Hungary based on radiocarbon dating of human bones. In: R.E. Németh and B. Rezi, eds., *Bronze Age Chronology in the Carpathian Basin*. Proceedings of the International Colloquium from Târgu Mureş 2-4 October 2014. Târgu Mureş: Editura MEGA, 23-36.
- Kordos, L., 1977. Changes in the Holocene climate of Hungary reflected by the 'vole-thermometer' method. *Földrajzi Közlöny*, 25 (101), 222-229.
- Kristiansen, K. and Suchowska-Ducke, P., 2015. Connected histories: the dynamics of Bronze Age interaction and trade 1500-1100 BC. *Proceedings of the Prehistoric Society*, 81, 361-392. <https://doi.org/10.1017/ppr.2015.17>
- Kyselý, R., 2010. Review of the oldest evidence of domestic fowl *Gallus gallus* f. *domestica* from the Czech Republic in its European context. *Acta Zoologica Cracoviensia: Series A: Vertebrata*, 53 (1-2), 9-34.
- Leuzinger, U. and Rast-Eicher, A., 2011. Flax processing in the Neolithic and Bronze Age pile-dwelling settlements of eastern Switzerland. *Vegetation History and Archaeobotany*, 20, 535-542. <https://doi.org/10.1007/s00334-011-0286-2>
- Ligeti, L., 1962. *A mongolok titkos története* [The secret history of Mongolian]. Translated by L. Ligeti. Budapest: Gondolat Kiadó.
- Ludwig, A., Pruvost, M., Reissmann, M., Benecke, N., Brockmann, G.A., Castaños, P., Cieslak, M., Lippold, S., Llorente, L., Malaspinas, A.-S., Slatkin, M., Hofreiter M., 2009. Coat color variation at the beginning of horse domestication. *Science*, 324, 485. <https://doi.org/10.1126/science.1172750>
- Luik, H. and Maldre, L., 2007. Bronze Age bone artefacts from Narkunai, Nevieriske and Kereliai fortified settlements: raw materials and manufacturing technology. *Archaeologia Litwana*, 8, 5-39.
- Marić Baković, M. and Car, G., 2014. Konzervatorsko-restauratorski radovi i rezultati najnovijih analiza na tekstilnome plaštu iz prapovijesnoga zemljanog tumula br. 16, Pustopolje, Kupres [Conservation and restoration works and the results of the latest analyses on the textile mantle from the prehistoric earthen Tumulus no. 16, Pustopolje, Kupres]. *Cleuna*, 1, 30-47.
- Matolcsi, J., 1982. Állattartás őseink korában [Animal keeping by our ancestors]. Budapest: Gondolat.

- Meadow, R.H. and Uerpmann, H.-P., eds., 1991. *Equids in the Ancient World: Volume II*. Beihefte zum Tübinger Atlas des Vorderen Orients: Reihe A (Naturwissenschaften) 19/2. Wiesbaden: Dr Ludwig Reichert Verlag.
- Metzner-Nebelsick, C., 2000. Early Iron Age pastoral nomadism in the Great Hungarian Plain: migration or assimilation? The Thraco-Cimmerian problem revisited. In: J. Davis Kimball, E.M. Murphy, L. Koryakova, L.T. Yablonsky, eds., *Kurgans, Ritual Sites and Settlements: Eurasian Bronze and Iron Age*. BAR International Series S890. Oxford: Archaeopress, 160-184.
- Montelius, O., 1885. *Om tidsbestämning inom bronsåldern med särskilt avseende på Skandinavien* [On Bronze Age dating with special regard to Scandinavia]. Stockholm: Akademien Förlag.
- Mozsolics, A., 1960. Die Herkunftsfrage der Ältesten Hirschgeweihtrensen. *Acta Archaeologica Academiae Scientiarum Hungaricae*, XII, 125-135.
- Németh, A., Bárány, A., Csorba, G., Magyari, E., Pazonyi, P., Pálffy, J., 2016. Holocene mammal extinctions in the Carpathian Basin: a review. *Mammal Review*, 47, 38-52. <https://doi.org/10.1111/mam.12075>
- Nesbitt, M., 2005. Grains. In: G. Prance and M. Nesbitt, eds., *The Cultural History of Plants*. London: Routledge, 45-60.
- Olalde, I., Brace, S., Allentoft, M.E., Armit, I., Kristiansen, K., Booth, T., Rohland, N., Mallick, S., Szécsényi-Nagy, A., Mitnik, A., Altena, E., Lipson, M., Lazaridis, I., Harper, T.K., Patterson, N., Broome and Khoshbacht, N., Diekmann, Y., Faltyskova, Z., Fernandes, D., Ferry, M., Harney, E., Knijff, P. de, Michel, M., Oppenheimer, J., Stewardson, K., Barclay, A., Alt, K.-W., Liesau, C., Ríos, P., Blasco, C., Vega Miguel, J., Menduiña García, R., Avilés Fernández, A., Bánffy, E., Bernabò-Brea, M., Billoin, D., Bonsall, C., Bonsall, L., Allen, T., Büster, L., Carver, S., Castells Navarro, L., Craig, O.E., Cook, G.T., Cunliffe, B., Denaire, A., Egging Dinwiddy, K., Dodwell, N., Ernée, M., Evans, C., Kuchařík, M., Francès, J., Farré, Fowler, C., Gazenbeek, M., Garrido Pena, R., Haber-Urriarte, M., Haduch, E., Hey, G., Jowett, N., Knowles, T., Massy, K., Pfrengle, S., Lefranc, P., Lemerrier, O., Lefebvre, A., Heras Martínez, C., Galera Olmo, V., Bastida Ramírez, A., Lomba Maurandi, J., Majó, T., McKinley, J.I., McSweeney, K., Mende, B.G., Modi, A., Kulcsár, G., Kiss, V., Czene, A., Patay, R., Endrődi, A., Köhler, K., Hajdu, T., Szeniczey, T., Dani, J., Bernert, Zs., Hoole, M., Cheronet, O., Keating, D., Velemínský, P., Dobeš, M., Candilio, F., Brown, F., Flores Fernández, R., Herrero-Corral, A.-M., Tusa, S., Carnieri, E., Lentini, L., Valenti, A., Zanini, A., Waddington, C., Delibes, G., Guerra-Doce, E., Neil, B., Brittain, M., Luke, M., Mortimer, R., Desideri, J., Besse, M., Brücken, G., Furmanek, M., Hałuszko, A., Mackiewicz, M., Rapiński, A., Leach, S., Soriano, I., Lillios, K.T., Cardoso, J. L., Parker Pearson, M., Włodarczak, P., Price, T.D., Prieto, P., Rey, P.-J., Risch, R., Rojo Guerra, M.A., Schmitt, A., Serrallongue, J., Silva, A.M., Smrčka, V., Vergnaud, L., Zilhão, J., Caramelli, D., Higham, T., Thomas, M.G., Kennett, D.J., Fokkens, H., Heyd, V., Sheridan, A., Sjögren, K.-G., Stockhammer, P.W., Krause, J., Pinhasi, R., Haak, W., Barnes, I., Lalueza-Fox, C., Reich, D., 2018. The Beaker phenomenon and the genomic transformation of northwest Europe. *Nature*, 555, 190-196. <https://doi.org/10.1038/nature25738>
- Olsen, S., Grant, S., Choyke, A.M., Bartosiewicz, L., eds., 2006. *Horses and Humans: the Evolution of Human Equine Relationships*. BAR International Series 1560. Oxford: Archaeopress.
- Outram, A.K., Stear, N.A., Bendrey, R., Olsen, S., Kasparov, A., Zaibert, V., Thorpe, N., Evershed, R.P., 2009. The earliest horse harnessing and milking. *Science*, 323 (5919), 1332-1335. <https://doi.org/10.1126/science.1168594>
- Pankovskiy, V. 2016. Deer antler cheek piece from Ushkalka-I: expertise and advice on typological classification. In: A. Zanoci, E. Kaiser, M. Kashuba, E. Izbitser, M. Băţchişinău, eds., *Mensch, Kultur und Gesellschaft von der Kupferzeit bis zur frühen Eisenzeit im nördlichen Eurasien: Beiträge zu Ehren Zum 60. Geburtstag*

- von Eugen Sava. Tyragetia International 1. Chişinău: National Museum of History of Moldova, 225-241.
- Parzinger, H., 2006. *Die frühen Völker Eurasiens: vom Neolithikum bis zum Mittelalter*. Historische Bibliothek der Gerda-Henkel-Stiftung 1. München: Beck.
- Piccione, P., Alvaro, C., Bartosiewicz, L., Lemorini, C., Masi, A., Sadori, L., 2015. Distribution of artefacts and ecofacts in an Early Bronze Age house in Eastern Anatolia: Space use and household economy at Arslantepe VI B2 (2900-2750 BCE). *Journal of Archaeological Science: Reports*, 4, 8-22. <https://doi.org/10.1016/j.jasrep.2015.08.035>
- Pitt, J., Gillingham, P.K., Maltby, M., Stewart, J.R., 2016. New perspectives on the ecology of early domestic fowl: an interdisciplinary approach. *Journal of Archaeological Science*, 74, 1-10. <https://doi.org/10.1016/j.jas.2016.08.004>
- Pletnyova: Плетнёва, С.А., 1967. От кочевий к городам. Салтово-Маяцкая культура. [From nomads to cities: the Saltovo-Mayatsk culture]. Москва: Наука.
- Rackham, H., 1938. *Natural History by Pliny the Elder*. Loeb Classical Library 330. Cambridge, MA: Harvard University Press.
- Provenzano, N., 2001. Worked bone assemblages in northern Italy Terramare: a technological approach. In: A.M. Choyke and L. Bartosiewicz, eds., *Crafting Bone: Skeletal Technologies through Time and Space*. Proceedings of the 2nd meeting of the (ICAZ) Worked Bone Research Group Budapest, 31 August-5 September 1999. BAR International Series S937. Oxford: Archaeopress, 93-103.
- Provenzano, N., 2003. Interactions entre hiérarchisation sociale et spécialisation artisanale dans l'âge du Bronze d'Italie septentrionale. In: P. Brun and P. de Miroschedji, eds., *Thème transversal no. 2: évolution des structures et dynamiques sociales: les moyens d'action sur les hommes*. Cahier des Thèmes Transversaux ArScAn 4. Nanterre: MAE Nanterre.
- Pruvost, M., Bellone, R., Benecke, N., Sandoval-Castellanos, E., Cieslak, M., Kuznetsova, T., Morales-Muñiz, A., O'Connor, T., Reissmann, M., Hofreiter, M., Ludwig, A., 2011. Genotypes of predomestic horses match phenotypes painted in Paleolithic works of cave art. *Proceedings of the National Academy of Sciences*, 108, 18626-18630. <https://doi.org/10.1073/pnas.1108982108>
- Raban-Gerstel, N., Bar-Oz, G., Zohar, I., Sharon, I., Gilboa, A. 2008. Early Iron Age Dor (Israel): a faunal perspective. *Bulletin of the American Schools of Oriental Research*, 349, 25-59. <https://www.jstor.org/stable/25067055>
- Rast-Eicher, A. and Reinhard, J., 1997. Gewebe und Geflechte. In: S. Hochuli, U. Niffeler, V. Rychner, eds., *Die Schweiz vom Paläolithikum bis ins Mittelalter: SPM III Bronzezeit*. Basel: Schweizerische Gesellschaft für Ur- und Frühgeschichte, 285-290.
- Robinson, D.E., 2003. Neolithic and Bronze Age agriculture in southern Scandinavia – recent archaeobotanical evidence from Denmark. *Environmental Archaeology*, 8, 145-165. <https://doi.org/10.1179/env.2003.8.2.145>
- Rütimeyer, L., 1861. *Die Fauna der Pfahlbauten der Schweiz*. Basel: Verlag von Bohnmaier's Buchhandlung (C. Detloff).
- Ryder, M.L., 1987. The evolution of the fleece. *Scientific American*, 256 (1), 112-119. <https://www.jstor.org/stable/24979302>
- Sabatini, S., 2018. Wool economy during the European Bronze Age. *Światowit*, 56, 43-55.
- Schibler, J., Hüster-Plogmann, H., Jacomet, S., Brombacher, C., Gross-Klee, E., Rast-Eicher, A., 1997. Ökonomie und Ökologie neolithischer und bronzezeitlicher Ufersiedlungen am Zürichsee: Ergebnisse der Ausgrabungen Mozartstrasse, Kanalisationssanierung Seefeld, AKAD/Pressehaus und Mythenschloss in Zürich. Monographien der Kantonsarchäologie Zürich 20. Zürich: Egg.
- Schibler, J. and Studer, J., 1998. Haustierhaltung und Jagd während der Bronzezeit in der Schweiz. In: S. Hochuli, U. Niffeler, V. Rychner, eds., *Die Schweiz vom Paläo-*

- lithikum bis zum frühen Mittelalter: Bronzezeit*. Basel: Verlag Schweizerische Gesellschaft für Ur- und Frühgeschichte, 171-191.
- Schmitzberger, M., 2009. *Haus- und Jagdtiere im Neolithikum des österreichischen Donauraumes*. Doctoral dissertation, Universität Wien.
- Schmölcke, U., Gross, D., Nikulina, E.A., 2018. The history of sheep husbandry in Austria from the Neolithic to the Roman period. *Annalen des Naturhistorischen Museums in Wien, Serie A*, 120, 101-126.
- Schreiber, R., 1973. *A harangedények népe Budapest* [The Bell Beaker people in Budapest]. Budapest: Budapesti Történeti Múzeum.
- Sherratt, A., 1983. The secondary products revolution of animals in the Old World. *World Archaeology*, 15, 90-104. <https://doi.org/10.1007/s10963-011-9045-7>
- Siracusano, G. and Bartosiewicz, L., 2012. Meat consumption and sheep/goat exploitation in centralised and non-centralised economies at Arslantepe, Anatolia. *Origini*, XXXIV, 111-123.
- Sofaer, J., Jørgensen, L.B., Choyke, A.M., 2013. Craft production: ceramics, textiles, and bone. In: H. Fokkens and A. Harding, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 469-491.
- Spengler, R.N., Frachetti, M., Doumani, P., Rouse, L., Cerasetti, B., Bullion, E., Mar'yashev, A., 2014. Early agriculture and crop transmission among Bronze Age mobile pastoralists of central Eurasia. *Proceedings of the Royal Society B: Biological Sciences*, 281, 20133382. <https://doi.org/10.1098/rspb.2013.3382>
- Stockhammer, P.W., Massy, K., Knipper, C., Friedrich, R., Kromer, B., Lindauer, S., Radosavljević, J., Wittenborn, F., Krause, J., 2015. Rewriting the central European Early Bronze Age chronology: evidence from large-scale radiocarbon dating. *PLoS ONE*, 10, 1-32. <https://doi.org/10.1371/journal.pone.0139705>
- Studer, Th., 1900. Entwicklung der Haustierzucht bei den Pfahlbauern. *Mitteilungen der Anthropologischen Gesellschaft in Wien* 30, 106-108.
- Szabó, G., 2017. A kárpát-medencei kora bronzkor periodizációjának nehézségei a régi és az újabb AMS radiokarbon adatok tükrében [Problems with the periodisation of the Early Bronze Age in the Carpathian Basin in light of the older and recent AMS radiocarbon data]. *Archeometriai Műhely*, XIV (2), 99-115.
- Tafuri, M.A., Craig, O.E., Cancim A., 2009. Stable isotope evidence for the consumption of millet and other plants in Bronze Age Italy. *American Journal of Physical Anthropology*, 139, 146-153. <https://doi.org/10.1002/ajpa.20955>
- Tegtmeier, U., 1993. *Neolithische und bronzezeitliche Pflugspuren in Norddeutschland und den Niederlanden*. Archäologische Berichte 3. Bonn: Holos.
- Thomas, N.R., 2014. A lion's eye view of the Greek Bronze Age. In: G. Touchais, R. Laffineur, F. Rougemont Peeters, eds., *Physis: l'environnement naturel et la relation homme-milieu dans le monde égéen protohistorique*. Actes de la 14e rencontre égéenne internationale, Paris, Institut National d'Histoire de l'Art (INHA), 11-14 décembre 2012. Aegaeum, Annales liégeoises et PAS Piennes d'archéologie égéenne 37. Leuven and Liège: Université de Liège, 375-390.
- Thrane, H., 1990. Bronzezeitlicher Ackerbau-Beispiel Dänemark. In: V. Furmánek and F. Horst, eds., *Beiträge zur Geschichte und Kultur der mitteleuropäischen Bronzezeit*. Berlin and Nitra: Zentralinstitut für Alte Geschichte und Archäologie der Akademie der Wissenschaften der DDR and Archeologický ústav Slovenskej akadémie vied, 483-493.
- VanDerwarker, A.M., 2010. Correspondence analysis and principal components analysis as methods for integrating archaeological plant and animal remains. In: A.M. VanDerwarker and T.M. Peres, eds., *Integrating Zooarchaeology and Paleoethnobotany: a Consideration of Issues, Methods, and Cases*. New York: Springer, 75-96.
- van Wijngaarden-Bakker, L.H. and Brinkkemper, O., 2005. All-round farming: food production in the Bronze Age and the Iron Age. In: L.P. Louwe Kooijmans, P.W.

- van den Broeke, H. Fokkens, A.L. van Gijn, eds., *The Prehistory of the Netherlands* 2. Amsterdam: Amsterdam University Press, 491-512.
- Vicze, M., 2013. Expecting the unexpected: Százhalombatta-Földvár surprises once again. In: S. Bergerbrant and S. Sabatini, eds., *Counterpoint: Essays in Archaeology and Cultural Heritage in Honour to Professor Kristian Kristiansen*. BAR International Series 2508. Oxford: BAR Publishing, 71-76.
- von Heer, O., 1866. *Die Pflanzen der Pfahlbauten*. Separatabdruck aus dem Neujahrsblatt der Naturforschenden Gesellschaft auf das Jahr 1866. Zürich: Naturforschende Gesellschaft.
- Vörös, I., 1981. Wild equids from the Early Holocene in the Carpathian Basin. *Folia Archaeologica*, 32, 37-68.
- Vörös, I., 1994. Animal husbandry and hunting in the Middle Neolithic settlement at Tiszavasvári-Deákalmi-dűlő (Upper Tisza region). *Jósa András Múzeum Évkönyve*, 36, 167-184.
- Vretemark, M., 2010. Subsistence strategies. In: T. Earle and K. Kristiansen, eds., *Organizing Bronze Age Societies: the Mediterranean, Central Europe, and Scandinavia Compared*. Cambridge: Cambridge University Press, 155-184.
- Vretemark, M. and Sten, S., 2010. Skeletal manipulation of dogs at the Bronze Age site of Százhalombatta-Földvár in Hungary. In: D. Campana, P. Crabtree, S. deFrance, J. Lev-Tov, A.M. Choyke, eds., *Anthropological Approaches to Zooarchaeology: Colonialism, Complexity and Animal Transformations*. Oxford: Oxbow, 210-213.
- West, B. and Zhou, B.X., 1988. Did chickens go north? New evidence for domestication. *Journal of Archaeological Science*, 15, 515-533. [https://doi.org/10.1016/0305-4403\(88\)90080-5](https://doi.org/10.1016/0305-4403(88)90080-5)

Unearthing millet in Bronze and Iron Age Croatia

*Kelly Reed, Jacqueline Balen, Ivan Drnić, Sara Essert,
Hrvoje Kalafatić, Marija Mihaljević, Emily Zavodny*

Abstract

New AMS ^{14}C -dates have been tracking the arrival and spread of millet across prehistoric Europe. They suggest that millet arrived in the Bronze Age at a time when significant socio-economic changes were underway. This paper presents new radiocarbon dates from the region of central and eastern Croatia where grains of both broomcorn (*Panicum miliaceum*) and foxtail millet (*Setaria italica*) were sampled from four prehistoric sites. The radiocarbon dating revealed a date of between the 16th to 12th century cal BCE for two of the sites, corresponding with stable isotope analyses in the region that indicate millet consumption occurred from the mid/late Bronze Age. Overall, the data from Croatia is still relatively sparse, so more information is needed to understand how, when and why millet became incorporated within the local agricultural food system.

Introduction

Croatia has a unique geography within which to explore prehistoric farming, with a coastal zone to the southwest and a continental region to the northeast, separated by the Dinaric Alps (Fig. 1). Within the coastal region, a hotter and drier Mediterranean climate is present and is characterised by a limestone geology that is more rugged and less uniform, with areas containing fertile valleys and mountainous areas covered by thin layers of poor quality soil. Inland Croatia in contrast has a continental climate with greater rainfall and cooler temperatures. As the region moves east it becomes relatively flatter with fertile alluvial plains that are roughly bordered by the Drava River to the north, the Sava to the south and the Danube to the east (Fig. 1). Agriculture arrived in Croatia ca. 6000 cal BCE through two routes; along the coast and through an inland route (*e.g.* Forenbaher and Miracle 2005; Orton *et al.* 2016). The first farmers cultivated einkorn (*Triticum monococcum*), emmer (*Triticum dicoccum*), barley (*Hordeum vulgare*), pea (*Pisum sativum*), lentil (*Lens culinaris*),




Figure 1. Map of Croatia showing the location of Bronze and Iron Age sites with archaeobotanical remains of millet referred to in this paper
 1. Nadin-Gradina 2. Monkodonja
 3. Lasinja 4. Sisak-Pogorelac
 5. Kalnik-Igrišće 6. Kaptol-Gradići 7. Mačkovac-Crišnjevi
 8. Mačkovac-Oštrovi
 9. Orubica-Veliki Šeš.

and flax (*Linum usitatissimum*). These main crops dominate assemblages for over three and a half millennia (Reed 2015, 2017). The Bronze Age, however, witnesses the influx of new crops, cereals, pulses and oil plants, in Europe. These crops do not seem to arrive simultaneously in the form of a ‘package’ but appear at different times during the Bronze Age (Valamoti 2016).

One of these ‘new’ crops was millet, i.e. broomcorn (*Panicum miliaceum*) and foxtail (*Setaria italica*) millet, fast-growing summer annuals. From the archaeobotanical remains in Croatia, we see individual grains of millet appearing in the Neolithic (Reed 2015), small clusters in the Eneolithic (Reed 2017) and large storage deposits in the Late Bronze Age (Mareković *et al.* 2015; Reed *et al.* 2021) and Early Iron Age (Reed and Drnić 2016). From this evidence alone we may surmise an early prehistoric introduction of millet to Croatia but possibly as a weed rather than as a crop. However, studies such as Motuzaite Matuzeviciute *et al.* (2013) and Filipović *et al.* (2018) have shown the importance of dating grains instead of relying on contextual information. Their studies generally showed that many of the millet grains were younger than their contexts, suggesting millet did not arrive in parts of Europe until the Bronze Age, or Late Bronze Age in the case of Germany. These finds highlight the potential for these small grains to move through archaeological contexts. Thus, in recent years singular grains found in contexts dated to before the Bronze Age, when large deposits of grains appear (e.g. Stika and Heiss 2013), have been interpreted as intrusive. Isotopic and biomolecular evidence in Europe also suggests that the preparation and consumption of millet only occurred from the Bronze Age onwards (e.g. Lightfoot *et al.* 2013; Heron *et al.* 2016).

In comparison to other regions, little is known about the development of agriculture and the introduction of these new species in the Croatian Bronze (ca. 2500 to 900/800 cal BCE) and Iron Age (ca. 900/800 to end of the 1st century BCE)

and how agriculture connected with the changing socio-economic environment. Thus, how, when and where millet was introduced to Croatia is still unclear. This paper presents recent AMS-radiocarbon dating of broomcorn and foxtail millet from four prehistoric sites in central and eastern Croatia. These dates were undertaken as part of the Kiel study funded by the German Research Foundation (Deutsche Forschungsgemeinschaft (DFG), Project number 2901391021-SFB 1266), in order to understand the precise chronology and character of the spread of broomcorn millet into Europe and the mechanism for its uptake and final incorporation within food production, diet and the wider socio-economy (Filipović *et al.* 2020).

Evidence of millet cultivation and consumption in prehistoric Croatia

Evidence of broomcorn or foxtail millet in Croatia is rather sporadic at prehistoric sites. Singular finds have been noted from the Neolithic and Eneolithic (Reed 2013), except at the multi-period site of Lasinja where 48 grains of foxtail millet and 6 broomcorn millet grains were identified from an area thought to belong to an Eneolithic settlement (Reed 2017). The site had no other crop remains other than one badly preserved barley grain (cf. *Hordeum vulgare*), with only singular finds of chinese lantern (*Physalis alkekengi*), blackberry (*Rubus* sp.) and elder (*Sambucus ebulus*) and a few weed/wild seeds.

For the Bronze Age only five sites have finds of millet: Monkodonja, Orubica-Veliki Šeš, Mačkovac-Crišnjevi, Mačkovac-Oštrovi and Kalnik-Igrišće. However, Monkodonja, which dates to the Early/Middle Bronze Age (2000-1200 BCE) only had two broomcorn millet grains (Kroll 2015) and Orubica-Veliki Šeš only had one grain (Reed 2013). These few grains have not been dated and so are not considered to indicate crop cultivation at this time. Instead, the most secure finds come from Kalnik-Igrišće where 37,000 broomcorn millet grains have so far been identified within a burnt down building (Mareković *et al.* 2015; Essert 2019). This site is particularly interesting as there are clear storage areas within the building, along with large deposits of other crops including barley, naked wheat (*Triticum aestivum/durum*), emmer (*Triticum dicoccum*), spelt (*Triticum spelta*), broad bean (*Vicia faba*) and lentil (*Lens culinaris*) (Reed *et al.* 2021). At Mačkovac-Crišnjevi, a settlement site, only 10 broomcorn millet grains were identified along with 4 foxtail millet grains. They were generally singular finds in general occupation layers except for one pit where five broomcorn millet grains were recovered. Interestingly, this site also had over 300 oat grains from two general occupation layers. Only one grain of emmer and one of lentil were also identified along with a range of grasses and sedges and other wild/weed species (Reed 2013). At Mačkovac-Oštrovi, a Bronze Age necropolis, 15 broomcorn millet grains were recovered from one of the cremation burials (Reed 2013). No other crops were identified and only three wild/weed seeds were recovered.

In the Iron Age, only three sites have archaeobotanical material published and all of these have yielded broomcorn millet grains to a greater or lesser extent. The first is Nadin-Gradina in Dalmatia, where millet was identified from Iron Age and Roman levels (Nye 1996). Although it is unclear from the publication which samples were dated to the Iron Age, no one sample contained more than 12 broomcorn millet grains and overall, the site yielded only 23 grains. Plant remains collected in 2010 from tumulus 14 at Kaptol-Gradci near Požega date to the Early Iron Age (Hallstatt) (Šaić 2014). In total, 3821 fruits, seeds, chaff and other plant remains were recovered. Here, only six broomcorn millet and one foxtail millet grains were identified. The site was dominated by spelt grain, followed by naked wheat and emmer. Other samples have been collected at the site, but millet was not identified

Site name	Monkodonja	Mačkovac-Oštrovi	Mačkovac-Crišnjevi
Reference	Kroll 2009; 2015	Reed 2013	Reed 2013
Date	ca. 1800-1200 cal BCE		ca. 1400-1100 cal BCE
No. of samples	64	3	25

Cereals

<i>Hordeum vulgare</i>	Grain	18		
<i>Triticum aestivum/durum</i>	Grain			
<i>Triticum dicoccum</i>	Grain	15		1
<i>Triticum dicoccum</i>	Chaff	28		
<i>Triticum monoccoccum</i>	Grain	2		
<i>Triticum monoccoccum</i>	Chaff	4		
<i>Triticum spelta</i>	Grain			
<i>Triticum spelta</i>	Chaff			
<i>Triticum spelta/aestivum</i>	Grain			
<i>Triticum spelta/dicoccum</i>	Grain			
<i>Triticum spelta/dicoccum</i>	Chaff			
<i>Triticum</i> sp.	Chaff			1
<i>Avena</i> sp.	Grain	1		343
<i>Avena</i> sp.	Chaff	1		13
<i>Secale cereale</i>	Grain			
<i>Panicum miliaceum</i>	Grain	2	16	10
<i>Setaria italica</i>	Grain			4

Pulses

<i>Lathyrus sativum</i>	Seed	1		
<i>Lens culinaris</i>	Seed			1
<i>Pisum sativum</i>	Seed	2		
<i>Vicia ervilia</i>	Seed	1		
<i>Vicia faba</i>	Seed	1		

Table 1. Cereals and pulses identified from Bronze and Iron Age sites in Croatia that contained millet.

(Šoštarić *et al.* 2008; 2017). The most significant site for this period is Sisak-Pogorelac, where a pot filled with over a million foxtail millet grains was recovered from the floor of a wooden building (Reed and Drnić 2016). Along with foxtail millet, around a hundred thousand broomcorn millet grains were also present, likely as a weed of the crop due to similarities in size. Work is continuing at the site and more samples are yielding both foxtail and broomcorn millet along with large numbers of broad beans and acorns (*Quercus* sp.), as well as a small number of other cereals, pulses, fruits and wild/weed species (Reed 2020).

Recent stable isotope analyses of humans and animal remains add to this evidence. In coastal Croatia, stable carbon and nitrogen isotope analyses were carried out on human and animal bones from three Bronze Age sites: Gusića Gomila, Jukića Gomila, and Brnjica (Miller 2018). Only two individuals from Brnjica showed signs of small consumption of C4 plants and/or marine foodstuffs. While in the neighbouring region of Lika, Zavodny *et al.* (2017) examined the stable isotopes of 36 human and

Orubica-Veliki Šeš	Kalnik-Igrišće	Nadin-Gradina	Sisak -Pogorelac	Kaptol-Gradići
Reed 2013	Mareković 2013; Mareković <i>et al.</i> 2015; Reed <i>et al.</i> 2021	Nye 1996	Reed, Drnić 2016; Reed 2020	Šaić 2014
ca. 1300-1100 cal BCE	ca. 1000-800 cal BCE	ca. 5th-1st century BCE	ca. 6th-4th century BCE	820-410 cal BCE
2	158		3	1

	17,475	+		5
	9576			124
	5847		24	91
	91			9
	410		8	4
				5
	2189	+		1620
	70	+		40
	6			69
	822			138
				9
		+	24	4
	71			2
	7			
1	36,931	+	104,000	6
	1037		1,342,667	1
	4973			
	12			
	9990		580	

30 faunal remains from 12 Bronze and Iron Age sites. They found no C4 signatures within the animal remains suggesting that millet (a C4 plant) was not regularly used for fodder (also Zavodny *et al.* 2019). Of the human remains, they found that most individuals appear to have eaten comparable proportions of millet, approximately 20% of the overall diet regardless of age or sex, from the Middle Bronze Age onwards (at Bezdanjača, 1430-1290 cal BCE and Veliki Vital, 1420-1295 cal BCE). As the Iron Age progresses, millet consumption increases to almost 40% of the diet at the sites of Mala Metaljka, Sultanov Grob and Trošmarija (Zavodny *et al.* 2017).

Lightfoot *et al.* (2015) demonstrated varying quantities of C4 carbon isotope signatures from the bone collagen of individuals dated to the Iron Age in Croatia, also suggesting the consumption of millet or the consumption of livestock foddered with millet. In particular, they concluded that

‘although millet was available to individuals throughout the study area and periods, it was consumed on a significant scale only in inland Croatia during the Iron Age’ (Lightfoot et al. 2015).

Additionally, they identified possible differentiations in the social status at the Iron Age coastal site of Nadin-Gradina between individuals buried in pits, who produced a more pronounced C4 signature, and those buried in stone-lined graves. Research by Nicholls (2017, 311) has also shown that individuals inhabiting central/eastern Slovenia and northeastern Croatia were ingesting a high quantity of C4 plants from the Middle Bronze Age to the Early Iron Age. In addition, the study found high $\delta^{13}\text{C}$ values in cattle, sheep/goat and pig remains at certain sites, suggesting the consumption of C4 plants, probably millet.

New millet radiocarbon dates

Since the paper by Motuzaite-Matuzeviciute *et al.* (2013), pre-Bronze Age finds of millet have generally been treated with caution. Thus, the discovery of a relatively large number of foxtail millet grains within an occupation layer thought to date to an Eneolithic settlement at the site of Lasinja raised questions regarding its provenance in the deposits, *i.e.* can they be securely dated to the Eneolithic? No millet grains have been dated in Croatia before this study and four sites, with relatively secure deposits, were selected for this project. These sites were Eneolithic contexts at Lasinja, Bronze Age Mačkovac-Crišnjevi and Mačkovac-Oštrovi and Iron Age Sisak-Pogorelac. Finds of both broomcorn and foxtail millet grains were selected. Within the present study, radiocarbon dating was conducted on single millet grains at the laboratory in Poznań. The standard protocols for sample processing followed Brock *et al.* 2010 (see Filipović *et al.* 2020 for further details).

The sites

Lasinja

Since 2010, the Archaeological Museum in Zagreb has been conducting systematic excavations at the site of Lasinja-Talijanovo Brdo. The site is situated on a natural hill to the south of the Kupa River and has, so far, yielded material dated to the Copper, Bronze and Iron Ages. Due to a nearby stone quarry, the south-western part of the site has been destroyed (Balen 2013, 399). No permanent structures have been discovered at the site, except working areas in the form of pits, so it seems likely that the position was used seasonally or when local raw materials were exploited. The excavated remains of material culture fall into the known repertoire of the Lasinja culture production of pottery. The recovered vessel fragments are of poor quality, but the most numerous pottery forms include bowls and pots, as well as ceramic spoons with holes for hafting. The site also yielded many chipped stone artefacts. Only two samples (*c.* 22 litres) taken from features thought to belong to the Eneolithic phase yielded archaeobotanical remains (Reed 2017). A grain of both broomcorn and foxtail millet was taken from one occupation layer context for radiocarbon dating.

Mačkovac-Crišnjevi

Systematic excavations began at the site in 1997 by the Nova Gradiška Municipal Museum. The settlement is located 1 km north of Mačkovac, which is situated on the left bank of the Sava River, approximately 15 km south of Nova Gradiška. The settlement is elevated up to 2 m above the floodplain and covers an area of approximately 2 hectares (Karavanić *et al.* 2002). Three trenches were opened in

1997 and 1998, covering around 323m². Numerous features have been identified within the excavated area including house floors, hearths, pits and evidence of a metallurgical work area. Previously, a Bronze Age hoard was recovered at the site in 1985 and, since then, further bronze items as well as pottery and animal remains have been found (Karavanić *et al.* 2002). The identification of several bronze needles and pottery types have dated the site from the Middle to beginning of the Late Bronze Age, concurrent with the Virovitica group and culturally belonging to the Barice-Gredani group (13th-12th century BCE). Between 2000 and 2003, 28 samples (308 litres) were collected from 18 features, with 25 of those containing carbonised plant remains (Reed 2013). A grain of broomcorn millet was taken from one of the pit samples for radiocarbon dating.

Mačkovac-Oštrovi

Mačkovac-Oštrovi is a Late Bronze Age necropolis and is believed to have belonged to the nearby settlement of Mačkovac-Crišnjevi. Between 2003 and 2009, excavations were led by the Nova Gradiška Municipal Museum. The excavations revealed 73 graves, belonging to the Barice-Gredani group of the Urnfield culture (Mihaljević and Kalafatić 2006, 2009, 2010). All the graves show a similar burial ritual, where the burnt bones of the deceased were collected into a vessel which was laid into the ground upside down (Mihaljević and Kalafatić 2009). Three samples (33 litres) were collected from three different grave areas, although two (from area M10) were from two closely associated graves. Here, one grain of broomcorn millet from within one of the cremation graves was radiocarbon-dated (Reed 2013).

Sisak-Pogorelac

The prehistoric settlement located in today's town of Sisak was founded near the confluence of the Kupa and the Sava Rivers in the later phase of the Late Bronze Age, around 1000 BCE. In the Late Hallstatt period, it became a key regional centre, and its importance continued in the Late Iron Age, when the settlement was spreading along both banks of the Kupa River (Drnić and Miletić Čakširan 2014; Drnić and Groh 2018; Drnić 2020). In 35 BCE, the indigenous settlements known from the ancient sources as Segestica and Siscia were conquered by the Romans, who established a military stronghold that was gradually transformed into an important provincial city – Colonia Flavia Siscia.

Ongoing excavations at the Pogorelac site in Sisak commenced in 2012, led by the Archaeological Museum in Zagreb. In trench 1, under a thick layer of house daub, the remains of an above-ground structure that was oriented northeast-southwest was found (Structure 2). It consisted of a yellow clay floor, several irregularly shaped rock fragments and the remains of wooden beams made of oak (*Quercus* sp.). A thick layer of daub and a large amount of charcoal, together with burnt wooden architectural parts and a clay floor suggest that the structure was destroyed by fire. Some typical Late Hallstatt pottery dates Structure 2 to between the end of the 6th and 4th centuries BCE. On the surface of the floor in Structure 2, a pot full of approximately 1343,000 charred grains of foxtail millet and 104,000 broomcorn millet grains were recovered. A grain of both broomcorn and foxtail millet from the pot sample was radiocarbon dated (Reed and Drnić 2016).

Results

The results of the radiocarbon dating of the broomcorn and foxtail millet grains can be seen in Table 2. Interestingly the foxtail and broomcorn millet from Lasinja have an Iron Age date of the 4th century cal BCE. The grain taken from the secure pot sample at Sisak-Pogorelac has a 4th-3rd century cal BCE date, which corresponds

Site name	Material	Sample number	Context	Calibrated age (95.4% / 2σ hpd range)	BP age	Laboratory code
Mačkovac-Crišnjevi	<i>Panicum miliaceum</i> grain	[SJ71]	Pit	1596-1135 BCE	3120 ± 70	Poz-104900
Mačkovac-Oštrovi	<i>Panicum miliaceum</i> grain	30 [SJ55]	Fill of pot in a cremation burial	1411-1223 BCE	3055 ± 35	Poz-104921
Sisak-Pogorelac	<i>Setaria italica</i> grain	60 [SJ105]	Fill of pot in building floor	399-207 BCE	2260 ± 35	Poz-100059
Lasinja	<i>Setaria italica</i> grain	11 [SJ2]	Occupation layer	360-53 BCE	2150 ± 50	Poz-100058
Lasinja	<i>Panicum miliaceum</i> grain	11 [SJ2]	Occupation layer	341-49 BCE	2115 ± 30	Poz-104923

Table 2. New radiocarbon dates for broomcorn and foxtail millet grains at Mačkovac-Crišnjevi, Mačkovac-Oštrovi, Sisak-Pogorelac and Lasinja.

with the archaeological evidence. Mačkovac-Crišnjevi had the oldest date of 16th to 12th century cal BCE and Mačkovac-Oštrovi had a date of 14th to 13th century cal BCE. These dates place the broomcorn millet age to the Late Bronze Age.

Discussion and conclusion

The new radiocarbon dates taken from the broomcorn and foxtail millet grains from Lasinja, Mačkovac-Crišnjevi, Mačkovac-Oštrovi and Sisak-Pogorelac provide important insights into the arrival and adoption of millet in Croatia. The concentration of foxtail and broomcorn millet from a possible Eneolithic occupation layer at Lasinja dates to the Iron Age rather than the Eneolithic. This suggests that the pit may date to the Iron Age, rather than being intrusive as little material culture was found in the pit and the site had evidence of Bronze Age and Iron Age occupations. Both Mačkovac-Crišnjevi and Mačkovac-Oštrovi have two broomcorn millet grains that date to the Late Bronze Age, while the foxtail millet from Sisak-Pogorelac is consistent with the Iron Age date of the site. These dates correspond with the stable isotope evidence that suggest millet began to be consumed during the Middle/Late Bronze Age, but possibly in a minor capacity. By the Late Bronze Age, millet became a more significant part of the diet in inland Croatia, while on the coast this may not have occurred until the early Iron Age.

Millet was not the only crop to have been introduced at this time, since archaeobotanical evidence showing an increase in cultivation of spelt (*Triticum spelta*) (Akeret 2005), free-threshing wheat (such as *Triticum aestivum/durum*) and broad bean (*Vicia faba*). Within Serbia, Hungary and Croatia we also see for the first time the oil plant gold-of-pleasure (*Camelina sativa*) and safflower (*Carthamus tinctorius*) during the Late Bronze Age (Kroll 1990; Medović 2002; Gyulai 2010, 105; Reed *et al.* 2021). The mechanisms behind the adoption of these new crops is still unclear as there are many factors involved in the decision to adopt or reject a new crop variety. Current evidence restricts determination of what these driving mechanisms may have been, but they could have been associated with environmental constraints, cultivation and processing technologies, nutritional and agronomic qualities, social relationships between source and receiving regions, and cultural associations of status, ritual and culinary appeal.

One factor considered recently by Zavodny *et al.* (2017) for the cultivation of millet in the Lika region of Croatia during this period was that locals began to diversify crops to manage food security. Diversification is a strategy that can be used to minimise the risk of crop failure through the cultivation of a wide range of crops with different growing conditions. They propose that the marginal environment, with poor soil and long winters, prompted the adoption of millet by farmers in order to stagger growing seasons with a greater range of crops and thus minimise risk (Zavodny *et al.* 2017). Even during the Roman period, Strabo (5.i.12) remarks that, in the Po River valley, millet was a secure protector against famine when other crops failed. This phenomenon of increased crop varieties was also seen across western

and central Europe (Bakels 1991, 2009) and the Mediterranean (Valamoti and Jones 2003). This suggests that agricultural strategies were intertwined with wider socio-economic or environmental changes that occurred during the Bronze Age.

Overall, the data from Croatia is still relatively sparse and more information is needed to understand when millet first arrived to the region and how millet cultivation and agricultural developments were integrated within other socio-economic developments.

References

- Bakels, C., 1991. Tracing crop processing in the Bandkeramik culture. In: J.M. Renfrew, ed. *New Light on Early Farming. Recent Developments in Palaeoethnobotany*. Edinburgh: Edinburgh University Press, 281-288.
- Bakels, C., 2009. *The Western European Loess Belt: Agrarian History, 5300 B.C.-A.D. 1000*. Dordrecht: Springer.
- Balen, J., 2013. Lokalitet Lasinja-Talijanovo brdo. *Hrvatski arheološki godišnjak*, 9/2012, 398-400.
- Brock, F., Higham, T., Ditchfield, P., Bronk Ramsey, C., 2010. Current pretreatment methods for AMS radiocarbon dating at the Oxford Radiocarbon Accelerator Unit (ORAU). *Radiocarbon*, 52, 103-112. <https://doi.org/10.1017/S0033822200045069>
- Drnić, I., 2020. The Iron Age settlement in Sisak. In: I. Drnić, ed. *Segestica and Siscia – a Settlement from the Beginning of the History*. Zagreb: Archaeological Museum in Zagreb, 41-54.
- Drnić, I. and Groh, S., 2018. Excavations and geophysical prospection at the Iron Age and Roman-period site at Sisak-Pogorelac position (2012-2017), *Vjesnik Arheološkog muzeja u Zagrebu*, 51, 67-140.
- Drnić, I. and Miletić Čakširan, I., 2014. A Late Iron Age Settlement on the Left Bank of the River Kupa in Sisak: Strabo's „Σισκία“, *Vjesnik Arheološkog muzeja u Zagrebu*, 47, 147-251.
- Essert, S., 2019. The results of the plant remains analysis. In: S. Karavanić and Kudelić, A., eds. *Kalnik-Igrišće – Late Bronze Age Settlement*, Monografije Instituta za Arheologiju 14, Zagreb: Institut za arheologiju, 160-166.
- Filipović, D., Meadows, J., Wiethold, J., Jahns, S., Bittmann, F., Kirleis, W., 2018. Before and after: millet cultivation and the transformation of prehistoric crop production in northern Germany. *Antiquity*, 92(365), e3. <https://doi.org/10.15184/aqy.2018.225>
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Kloof, S., KohlerSchneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A.M., MuellerBieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H.P., Stobbe, A., Tolar, T., Wasylkowa, K., Wiethold, J., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Forenbaher, S. and Miracle, P.T., 2005. The spread of farming in the Eastern Adriatic. *Antiquity*, 79, 514-528. <https://doi.org/10.1017/S0003598X00114474>
- Gyulai, F., 2010. *Archaeobotany in Hungary: Seed, Fruit, Food and Beverage Remains in the Carpathian Basin from the Neolithic to the Late Middle Ages*. Budapest: Archaeolingua.
- Heron, C., Shoda, S., Breu Barcons, A., Czebreszuk, J., Eley, Y., Gorton, M., Kirleis, W., Kneisel, J., Lucquin, A., Müller, J., Nishida, Y., Son, J., Craig, O.E., 2016. First molecular and isotopic evidence of millet processing in prehistoric pottery vessels. *Scientific Reports*, 6, 38767. <https://doi.org/10.1038/srep38767>

- Karavanić, S., Mihaljević, M., Kalafatić, H., 2002. The Mačkovac-Crišnjevi settlement as a contribution to identifying the beginnings of Urnfield Culture in Slavonian Posavina. *Prilozi Instituta za arheologiju Zagrebu*, 19, 47-62.
- Kroll, H., 1990. Salford von Feudvar, Vojvodina. *Archäologisches Korrespondenzblatt*, 20, 41-46.
- Kroll, H., 2015. Spuren prähistorischer Nutzpflanzen aus Monkodonja – Die Pflanzensfunde. In: Hänsel, B., Mihovilić, K., Teržan, B., eds. *Monkodonja 1: Forschungen zu einer protourbanen Siedlung der Bronzezeit Istriens*, Monografije i katalogi Arheološkog muzeja Istre 25, Bonn: Dr Rudolf Habelt GmbH, 104-108.
- Lightfoot, E., Liu, X., Jones, M.K., 2013. Why move starchy cereals? A review of the isotopic evidence for prehistoric millet consumption across Eurasia. *World Archaeology*, 45(4), 574-623. <https://doi.org/10.1080/00438243.2013.852070>
- Lightfoot, E., Šlaus, M., Rajić Šikanjić, P., O'Connell, T.C., 2015. Metals and millets: Bronze and Iron Age diet in inland and coastal Croatia seen through stable isotope analysis. *Archaeological and Anthropological Sciences*, 7 (3), 375-386. <https://doi.org/10.1007/s12520-014-0194-3>
- Mareković, S., Karavanić, S., Kudelić, A., Šoštarić, R., 2015. The botanical macroremains from the prehistoric settlement Kalnik-Igrišće (NW Croatia) in the context of current knowledge about cultivation and plant consumption in Croatia and neighbouring countries during the Bronze Age. *Acta Societatis Botanicorum Poloniae*, 84 (2), 227-235.
- Medović, A., 2002. Archäobotanische Untersuchungen in der metallzeitlichen Siedlung Židovar, Vojvodina/Jugoslawien: Ein Vorbericht. *Starinar*, 52, 181-190.
- Mihaljević, M., Kalafatić, H., 2006. Mačkovac-Crišnjevi (naselje i nekropola). *Hrvatski arheološki godišnjak*, 2/2005, 48-49.
- Mihaljević, M., Kalafatić, H., 2009. Mačkovac-Crišnjevi (naselje i nekropola). *Hrvatski arheološki godišnjak*, 5/2008, 117-118.
- Mihaljević, M., Kalafatić, H., 2010. Mačkovac-Crišnjevi (naselje i nekropola). *Hrvatski arheološki godišnjak*, 6/2009, 69-70.
- Miller, D., 2018. *Stable Carbon and Nitrogen Isotope Analysis in Italy and Croatia: Bronze Age Food Practices Across the Adriatic*. Master's thesis, Sapienza University of Rome.
- Motuzaitė-Matuzevičiute, G., Staff, R.A., Hunt, H.V., 2013. The early chronology of broomcorn millet (*Panicum miliaceum*) in Europe. *Antiquity*, 87, 1073-1085. <https://doi.org/10.1017/S0003598X00049875>
- Nicholls, R.A., 2017. *More than Bones: an Investigation of Life, Death and Diet in Later Prehistoric Slovenia and Croatia*. Doctoral dissertation, University of Bradford.
- Nye, S., 1996. The plant remains. In: J. Chapman, R. Shiel, Š. Batović, eds. *The Changing Face of Dalmatia*. London: Leicester University Press, 240-242.
- Orton, D., Gaastra, J., Linden, M.V., 2016. Between the Danube and the Deep Blue Sea: Zooarchaeological meta-analysis reveals variability in the spread and development of Neolithic farming across the western Balkans. *Open Quaternary*, 2 (6), 1-26. <http://doi.org/10.5334/oq.28>
- Reed, K., 2013. *Farmers in Transition: the Archaeobotanical Analysis of the Carpathian Basin from the Late Neolithic to the Late Bronze Age (5000-900 BC)*. Doctoral dissertation, University of Leicester.
- Reed, K., 2015. From the field to the hearth: plant remains from Neolithic Croatia. *Vegetation History and Archaeobotany*, 24 (5), 601-619. <https://doi.org/10.1007/s00334-015-0513-3>
- Reed, K., 2017. Agricultural change in Copper Age Croatia (ca. 4500-2500 cal B.C)? *Archaeological and Anthropological Science*, 9 (8), 1745-1765. <https://doi.org/10.1007/s12520-016-0330-3>
- Reed, K., 2020. The production and preparation of food at the Iron Age settlement in Sisak. In: Drnić, I., ed. *Segestica and Siscia – a Settlement from the Beginning of the History*. Zagreb: Archaeological Museum in Zagreb, 55-62.

- Reed, K. and Drnić, I., 2016. Iron Age diet at Sisak, Croatia: archaeobotanical evidence of foxtail millet (*Setaria italica* [L.] P. Beauv.). *Oxford Journal of Archaeology*, 35 (4), 359-368. <https://doi.org/10.1111/ojoa.12101>
- Reed, K., Kudelić, A., Essert, S., Polonijo, L., Karavanić, S., 2021. House of plenty: Reassessing food and farming in Late Bronze Age Croatia. *Environmental Archaeology*, <https://doi.org/10.1080/14614103.2021.1979385>
- Šaić, N., 2014. *Arheobotaničko istraživanje prapovijesnog lokaliteta Kaptol – Gradci kraj Požege*. Master's thesis, University of Zagreb.
- Šoštarić, R., Potrebica, H., Brigić, A., 2008. Direct dating of botanical samples in an archaeological context – plant remains from the prehistoric site of Kaptol-Gradci near Požega (Croatia). *Prilozi Instituta za arheologiju u Zagrebu*, 24, 79-88.
- Šoštarić, R., Potrebica, H., Hršak, J., Essert, S., 2017. Archaeobotanical components of grave goods in prehistoric tumuli 6 and 7 at the archaeological site of Kaptol-Gradci, near Požega (Croatia). *Acta Botanica Croatica*, 76 (2), 183-190. <https://doi.org/10.1515/botcro-2017-0004>
- Stika, H.-P. and Heiss A.G., 2013. Plant Cultivation in the Bronze Age. In: Fokkens, H. and Harding, A. eds. *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 348-369.
- Valamoti, S.M., 2016. Millet, the late comer: on the tracks of *Panicum miliaceum* in prehistoric Greece. *Archaeological and Anthropological Science*, 8, 51-63. <https://doi.org/10.1007/s12520-013-0152-5>
- Valamoti, S.M. and Jones, G., 2003. Plant diversity and storage at Mandalo, Macedonia, Greece: archaeobotanical evidence from the Final Neolithic and Early Bronze Age. *Annual of the British School at Athens*, 98, 1-35. <https://www.jstor.org/stable/30073203>
- Zavodny, E., Culleton, B.J., McClure, S., Kennett, D.J., Balen, J. 2017. Minimizing risk on the margins: insights on Iron Age agriculture from stable isotope analyses in central Croatia. *Journal of Anthropological Archaeology*, 48, 250-261. <https://doi.org/10.1016/j.jaa.2017.08.004>
- Zavodny, E., McClure, S., Welker, M., Culleton, B.J., Balen, J., Kennett, D., 2019. Scaling up: stable isotope evidence for the intensification of animal husbandry in Bronze-Iron Age Lika, Croatia. *Journal of Archaeological Science: Reports*, 23, 1055-1065. <https://doi.org/10.1016/j.jasrep.2018.10.008>

The earliest finds of millet and possible associated changes in material culture in Slovenia

Tjaša Tolar and Primož Pavlin

Abstract

We present an overview and review of archaeobotanical results on the earliest occurrence of crop plant remains from the area of what is today Slovenia, focusing primarily on the phenomenon of the appearance of millet and associated material culture in the Bronze Age and the Iron Age. In addition, we present an overview of the development of prehistoric harvesting tools and settlement patterns in Slovenia from the Neolithic to the Iron Age. Based on the available results from archaeological sites in Slovenia, the appearance of broomcorn millet and foxtail millet, together with oat, rye, lentil and faba bean, can be placed in the Late Bronze Age. In the Early Iron Age, the importance of millets increases, while the introduction of free-threshing wheat represents another agricultural innovation at this time. Just as emmer and einkorn are typical cereals of the Copper Age in the region, so millets are of the Late Bronze Age and, especially, of the Early Iron Age. Pulses were less common in the Copper Age and became more important in the Late Bronze and Early Iron Age. Barley seems to have been equally important through all of these periods. Collected wild fruits and nuts were more important for the diet during the Copper Age than during the Late Bronze Age and Early Iron Age. However, we emphasise that, the way in which archaeobotanical sampling/collecting, storing and sieving was done (if it was done at all) may have played an important role in the amount and type of results obtained. Even so, some major conclusions on the spectrum of crops used in different periods remain valid.

Introduction

Slovenia is located in southeastern Europe, along northernmost part of the Adriatic Sea, at the intersection of the Alpine, Mediterranean, Pannonian and Dinaric worlds (Fig. 1). It borders Italy to the west, Austria to the north, Hungary to the north-east and Croatia to the east and south. The topography of Slovenia is relatively diverse

(including mountains, mountain plateaus, plains, valleys, karst fields, and the Adriatic coast), and the different landforms alternate over short distances. The transition between landscapes or parts of them is gradual. Due to its geographical position, Slovenia is a distinct transition country. Important transport links between Italy and central and eastern Europe have always passed through its territory. The climate is temperate, being a mixture of influences of the Alpine, Mediterranean and continental climates.

Little is known about the prehistoric plant economy (cultivation and gathering) in Slovenia, with the exception of that of pile-dwellings from the 4th millennium BCE (Copper Age). Therefore, the answers to the questions of when the first signs of cultivation appeared, where this cultivation came from and how it developed after its introduction are still incomplete. This study reviews and extends the earliest archaeobotanical data on cultivation and plant gathering in Slovenia. The aim is to provide a long-term perspective on plant-human interactions, together with archaeological information from specific sites in the area where archaeobotanical analyses were carried out, focusing on the earliest known evidence of broomcorn millet (*Panicum miliaceum*) cultivation.

The cultivation of plants (as well as the material culture) in what is today Slovenia most probably originated from the southeastern regions of the Balkans. It is also very likely that some influences arrived via the Mediterranean route, across the Adriatic Sea and from the Apennine Peninsula (e.g. Ihde 1995). Despite scarce archaeobotanical evidence, it is generally accepted that plant cultivation was practised in the territory of Slovenia by the Late Neolithic. The oldest flint stones with 'gloss' (referring to a characteristic shine that proves that they were used as harvesting tools) come from Late Neolithic layers (5th millennium BCE) at Movernava, in Bela krajina, and at Dragomelj, near Ljubljana (see Results).

Our contribution summarises archaeobotanical and archaeological evidence from Slovenia that could provide hints on when, how and why millet became one of the most common crops. The three main areas covered are the oldest ¹⁴C-dated millet grains and their frequency through time; possible climatic changes in the period when millet appeared; and the oldest finds of harvest implements and their increasing efficacy through time, as a possible indicator for improving agricultural techniques.

Summary of economic activity and settlement patterns from the Neolithic to the Iron Age in Slovenia

The Neolithic in Europe was characterised by some major changes in the way people lived. They settled in permanent settlements, and agriculture and the keeping of livestock became the main economic activities. They cultivated cereals and pulses and raised animals, such as sheep, goats, cattle and pigs. The production and use of ceramic vessels was extremely important because the ability to cook foods radically changed eating habits. Easier access to food led to population growth and an incipient division of labour, while surplus production boosted trade, all of which led to social stratification (e.g. Bogaard and Styring 2017).

The territory of present-day Slovenia was influenced by the developing Neolithic cultures that surrounded it. The interior and the eastern part of Slovenia were connected with the Pannonian Basin, whereas the Karst and the Slovenian coastal area were connected with the eastern Adriatic coast and thus indirectly with the Mediterranean Sea. In western Slovenia, traces of Neolithic settlement have been found mainly during excavations of karst caves, while in central and eastern Slovenia settlement traces have been found either on the plain, on river terraces, at river bends or on naturally protected elevations (Guštin 2005). The settlement of Gradec, near Mirna, occupied in the middle of the 5th millennium BCE, was

protected by a stone wall (Dular *et al.* 1991, 84-90, 119-128, 140-142). The first pile-dwelling settlement appeared on the shore of the lake in what is now known as the Ljubljansko barje [the Ljubljana wetlands] (Velušček 2006).

The first metal age was the Copper Age, also known as the Eneolithic. In addition to stone, horn and bone tools and weapons, copper objects appeared in small quantities. The first finds related to copper metallurgy in Slovenia date back to the middle of the 4th millennium BCE. The objects in question are a piece of copper slag and fragments of crucibles from the wetland pile-dwellings of Hočevarica (Šmit 2004), Maharski prekop (Velušček and Greif 1998, 31-33) and Stare gmajne (Velušček 2009, 18-25). Copper metallurgy experienced a real heyday in the first half of the 3rd millennium BCE. The so-called Dežman pile-dwellings, near Ig, produced numerous copper objects (awls, daggers, axes) and, even more importantly, typical metallurgical accessories, such as moulds, crucibles and clay blowpipes, which indicate the local production of copper objects there (Korošec and Korošec 1969). Compared with the Neolithic, Slovenia showed a much denser settlement pattern in the Copper Age. Settlements developed in a similar environment as in the Neolithic period, and pile-dwelling settlements flourished in the Ljubljansko barje.

Little is known about the settlement pattern of Slovenia in the Early Bronze Age. At that time, the pile-dwelling culture in the Ljubljansko barje declined, while the Prekmurje plains were relatively densely populated, as recent research has shown (Šavel 2009; Kerman 2011a-c; Guštin, Tomaž 2016; Pavlin 2015; Guštin *et al.* 2017). Both regions belonged to the same cultural area, which was characterised by the so-called Litzen pottery.

In the Middle Bronze Age, the Karst and Istria experienced the rise of the Kaštelir culture, which continued uninterrupted into the Iron Age. The culture is named after the *kaštelir*, a settlement fortified with stone walls. The Štajerska (Styria) region was part of the central European Tumulus culture, which was characterised by tumulus burials. The territory of Slovenia was at that time a transit zone on the trade route between Italy and the Danube Basin, as indicated by the characteristic sword and sickles found by a chance in the riverbed of the Ljubljanica (Pavlin 2006; Turk *et al.* 2009, 228) and by the short sword from Lavrica (Šinkovec 1996, 143, Fig 21.1).

Towards the end of the 14th century BCE, Europe experienced great changes related to the mass migrations of peoples. These caused the collapse of the Mycenaean culture in Greece, the Hittite kingdom, and numerous cities in the Middle East. The so-called Sea Peoples pushed as far as Egypt, where they were defeated by Ramses III.

Because of the nature of the burials at this time, the Late Bronze Age is also known as the Urnfield period. Central and eastern Slovenia were inhabited by people who cremated the deceased and buried their remains in urns in flat necropolises. They belonged to the cultural sphere that stretched from western Hungary, through Croatia to northern Bosnia.

In the Late Bronze Age in central and eastern Slovenia, settlements were located on naturally protected, elevated terrain, which was additionally fortified with wooden palisades. Recent excavations during the construction of motorways in northeastern Slovenia have revealed numerous lowland Bronze Age settlements along the rivers Drava and Mura (Dular 2013, 101-110, Figs. 44-48). The shape of these settlements, their spatial context, and the construction of their houses are illustrated by the following examples.

Near Orehova vas, there was a settlement on the plain, by a stream. Here, 27 floor plans of buildings were discovered, which had been built using earth-fast construction (Grahek 2015, 29-41, 349). The scattered settlement of Oloris was surrounded by a moat, over which a wooden fence stood. The walls of the houses consisted of upright beams carrying interwoven wattle and daub. The houses had a fireplace and a storage pit under the floor (Dular *et al.* 2002, 23-46). The settlement

of Ormož had a slightly different kind of protection. In the south, it was protected by the river Drava, while the other three sides were surrounded by a dike and an earthwork. Inside the settlement, a network of roads up to 4 m wide and paved with gravel were documented. These streets were lined with houses of different sizes. They were built in the same way as at Oloris (Dular and Tomanič Jevremov 2010, 83-97). Excavations at the Tribuna site, in Ljubljana, have also brought to light a settlement from the Late Bronze Age (10th-8th century BCE) with an urban design. It consisted of houses of different sizes, also variously spaced, built in earth-fast construction. The houses were arranged in a grid, and the streets between them were paved (Vojaković *et al.* 2011, 26-31).

Contacts with the Mediterranean region also brought knowledge related to iron metallurgy. This and the abundance of easily accessible limonite iron ore formed the basis for the establishment of iron production towards the end of the 9th century BCE, which is thus taken as the start of the Iron Age locally. For a brief insight into Iron Age settlement, we can use data from the most researched region, Dolenjska. Here, new settlements were established either in the vicinity of older Urnfield settlements, which slowly died out, or in previously uninhabited areas. As a rule, they stood on elevated ground and were fortified by 2-3 m thick stone walls that followed the shape of the terrain. The walls were built using the dry-stone wall technique, from unworked stone blocks, and reinforced with vertical wooden beams driven into the ground. They were built in one go, and some were later repaired or supplemented. The surface areas of the settlements differ in size, from just a few to more than 10 ha. With an area of about 20 ha and 2.3 km long walls, Cvinger, near Vir, which is near Stična, was a real metropolis for that time. The houses were built in two ways. In the first method, which was adopted from the Bronze Age, upright posts were driven into the ground (earth-fast construction). In the second method, the upright posts were not driven into the ground, but stood on a horizontal beam (post-pad construction). This was laid on the bare earth or, sometimes, on a stone drainage bed. The floors of the houses were made of beaten clay. Near one of the walls was the fireplace, around which the kitchen equipment was arranged: ceramic vessels, hand mills for grinding grain, firedogs, clay rings and coils. The weights hanging on strings testify that looms were also among the household items (Gabrovec 1994).

The settlement with an urban character at Most na Soči, right on the border with Italy, was different. It stood on a naturally protected plateau above the confluence of two rivers. On the third side, access was protected by a stone wall. The houses were located on a sunny slope. They stood in rows along cobbled streets lined with ditches and canals. They had a stone foundation on which a wooden structure of horizontal and vertical beams and planks stood (Svoljšak and Dular 2016).

Short history of archaeobotanical investigations in Slovenia

The investigation of fossil plant macro-remains from archaeological sites is considered to have started relatively early in Slovenia, *i.e.* when the first pile-dwelling site in the Ljubljansko barje region was discovered, in 1875. The first results on its rich archaeobotanical remains, with a focus on wooden piles, were published soon after (Deschmann 1875, 1878). Since then, archaeobotanical reports from pile-dwellings have been common, but only isolated (occasional) plant macro-remains were collected during the excavations prior to 1998 (Schmid 1910, 1915; Korošec 1953; Šercelj 1981-1982; Culiberg 1984). The wet-sieving technique was first tested in Slovenia in 1989, at the Palaeolithic cave site of Divje Babe I (Turk 2007), where mainly charcoal fragments were extracted (Culiberg 2007). Thereafter, the flotation method was used at some other archaeological sites, *e.g.* at the Early Iron Age sites of southeastern Slovenia (Culiberg and Šercelj 1995; Dular and Tecco Hvala 2007).

Finally, in 1998, the wet-sieving method was employed at a waterlogged site as well – the Hočevarica pile-dwelling site (Velušček 2004). Thanks to the excellent preservation through waterlogging, the results for this site were outstanding (Jeraj 2004; Jeraj *et al.* 2009). In 2007, the method was supplemented with the semi-flotation technique, based on the Swiss model (Tolar *et al.* 2010). The first application of this method was at the Stare gmajne pile-dwelling settlement with rich assemblage of waterlogged plant material (Tolar *et al.* 2011). Since then, 6 cultivated and more than 20 gathered plant taxa have been regularly identified at the lake dwellings of the Ljubljansko barje (4th-3rd millennium BCE; Copper Age), which is comparable to the plant spectrum at other circum-Alpine lake dwellings (Tolar *et al.* 2011, 2016; Tolar 2018a).

Nowadays, most excavations in Slovenia use wet-sieving techniques and various methods of sampling (*i.e.* profile, systematic and judgement; Andrič *et al.* 2016, 62-65). On dry-land sites, plant remains are poorly preserved and mostly charred; charcoal predominates. Unfortunately, careful extraction of archaeobotanical remains is still not a standard, as some archaeologists do not pay enough attention to field sampling and subsequent proper processing of archaeobotanical sediment samples (*i.e.* fine wet sieving with semi-flotation using smaller than 2 mm mesh size). The loss of fragile and small seeds, such as millet grains, is certainly a consequence of negligence. Therefore, there is a great potential to find earlier archaeobotanical evidence of agriculture in Slovenia, especially given the geographical location of its territory.

Materials and methods

The main method used in this study is a review of archaeobotanical and archaeological investigations of selected prehistoric sites in Slovenia. We focused on the earliest evidence of cultivated plants and related agricultural tools, such as harvesting knives. We consider 14 sites in the archaeobotanical part of this study (Table 1; Fig. 1) and we include some additional ones in the archaeological review. They were selected according to the earliest absolutely dated evidence of crop macro-remains and the accessibility of archaeological and archaeobotanical data.


Data sources and study sites

The dataset on archaeological features and archaeobotany comes from existing publications and from manuscripts that are in preparation for publication. We have summarised our own results, including those unpublished (Archive of the Institute of Archaeology ZRC SAZU). Unfortunately, the quantification of the archaeobotanical results is not comparable, because some datasets are presented as absolute numbers of plant remains in an unknown amount of sediment, some as concentrations per litre of sediment, some as grams, and some as frequencies (see Table 2). The main objective is to make an inventory of the earliest finds of cultivated and gathered plant remains, focusing on the appearance of millet.

To track the earliest (Copper Age) absolutely dated traces of plant cultivation (remains of *e.g.* barley and wheat) in Slovenia, this study considers two different site types – dry-land and waterlogged. At the dry-land cave site of Ajdovska jama, excavated several times and over many years (from 1884 to 1990), less reliable archaeobotanical results from older excavations are available. Five cultural horizons were defined here, from the Late Neolithic to the Middle Ages, by different excavators (Horvat 1989; Culiberg *et al.* 1992 and references therein). The second horizon, radiocarbon dated to 5625±130 BP (after Culiberg *et al.* 1992) was determined to be the most intensively visited – the cave was used as a cemetery and ritual site in this period. At the time of the early excavations, flotation and wet sieving were not yet established and the recovered archaeobotanical material is scarce. In this study, we refer to a ¹⁴C-dated

Table 1. The 14 sites with archaeobotanical data considered in this study (excavation years noted in parentheses).

Copper Age	Late Bronze Age	Early Iron Age
Ajdovska jama (1884, 1938, 1967, 1982-1990)	Dragomelj (1997, 2000-2001)	Six sites in southeastern Slovenia (1990-2000)
Strojanova voda (2012)	Orehova vas (2006-2007)	Ljubljana-Tribuna (2008)
Stare gmajne (2007)	Grosuplje (2011)	Most na Soči (1971-1984)



- | | | | |
|-------------------|--------------------------|-----------------------------|----------------------|
| 1 Ajdovska jama | 5 Orehova vas | 9 Cvinger above Korita | 13 Ljubljana-Tribuna |
| 2 Stare gmajne | 6 Grosuplje | 10 Gradec near Vinkov vrh | 14 Most na Soči |
| 3 Strojanova voda | 7 Kincelj above Trbinc | 11 Cvinger near Meniška vas | |
| 4 Dragomelj | 8 Kunkel near Vrhtrebnje | 12 Gradec above Mihovo | |

Figure 1. The location of archaeobotanically-studied sites mentioned in the text. The six Early Iron Age sites in southeastern Slovenia are those marked with numbers 7-12.

stratigraphic unit (labelled SE43), where most of the cultivated plant remains were found and identified (see Table 2). It yielded from 1 to 1050 carbonised grains of crop plants in an unknown amount of sediment. The frequencies of the seeds are listed in Table 2; for the original absolute numbers of seeds/fruits, see Culiberg *et al.* (1992).

The second-earliest crop cultivation sites so far recorded are the two waterlogged Copper Age pile-dwelling sites of Strojanova voda and Stare gmajne. Both were recently investigated and examined with fine flotation methods, using up to 0.355 mm mesh size (see Tolar *et al.* 2010). The results are representative and reliable, expressed in seed/fruit concentrations per litre of sediment sample (Table 2; Tolar *et al.* 2011; Tolar 2018a).

The earliest archaeobotanical evidence of millet cultivation in Slovenia comes from three locations, all from the 12th century cal BCE (Late Bronze Age): Dragomelj, Orehova vas and Grosuplje (Turk and Svetličič 2005; Tolar (in Grahek 2015); Archive of the Institute of Archaeology ZRC SAZU).

For the 'post-millet appearance' period, this study considers the Early Iron Age sites from southeastern Slovenia: Cvinger near Meniška vas; Kunkel near Vrhtrebnje; Kincelj above Trbinc; Gradec above Mihovo; Cvinger above Korita; Gradec near Vinkov vrh (Culiberg, Šercelj 1995; Dular and Tecco Hvala 2007); Ljubljana-Tribuna (available from the Archive of the Institute of Archaeology ZRC SAZU; Vojaković *et al.* 2011); and Most na Soči (Tolar 2018b) (see Table 2; Fig. 1). Numerous other sites either have been archaeobotanically analysed but not yet published due to the limited results and/or absence of absolute-dated archaeobotanical remains (Archive of the Institute of Archaeology ZRC SAZU).

Results and discussion

Overview of the archaeobotanical finds of cultivars in Slovenia from the Late Neolithic to the Early Iron Age

Before the appearance of millet

Wild plants were used throughout Slovenia during and after the Stone Age (Culiberg 1999, 2011). The earliest absolutely dated macro-remains of cultivated plants in Slovenia are naked and hulled barley (*Hordeum vulgare*), einkorn (*Triticum monococcum*) and emmer (*Triticum dicoccum*). They were found at pile-dwelling settlements of the Ljubljansko barje (4th millennium cal BCE) and in the Ajdovska jama cave (5th-4th millennium cal BCE) (Table 2; Culiberg *et al.* 1992; Jeraj *et al.* 2009; Tolar *et al.* 2011; Tolar 2018a).

The earliest plant cultivation in dry locations in Slovenia is poorly understood. An example is the cave cemetery of Ajdovska jama, which was in use from the Late Neolithic onwards and where large assemblages of macro-remains of cultivated plants (such as wheat and barley) were discovered, but which were unfortunately insufficiently documented in terms of stratigraphy and chronology (Culiberg *et al.* 1992). This could be the site with the earliest evidence of plant cultivation in Slovenia. Barley and wheat predominate, but there are also remains of oat (*Avena sativa*) and faba bean (*Vicia faba*) that were not radiocarbon dated.

Dating to the well-researched Late Neolithic, Copper Age and Early Bronze Age (*i.e.* 4600-1700 cal BCE), the waterlogged pile-dwelling sites of the Ljubljansko barje provided a lot of representative data on plant, animal and cultural material. Most of the results come from the 4th millennium BCE (Copper Age) sites, where the cultivation of emmer, einkorn, barley (both naked and hulled), flax (*Linum usitatissimum*), pea (*Pisum sativum*) and opium poppy (*Papaver somniferum*), as well as the domestication of dog, sheep, goat, cattle and pig, has been established (*e.g.* Toškan and Dirjec 2004; Tolar *et al.* 2011). Archaeobotanical results from younger (3rd and 2nd millennium BCE) pile-dwellings are sparse due to the unsuitable recovery methods used prior to 2007 (Tolar *et al.* 2010) and because, subsequently, no excavations have been carried out of the youngest pile-dwelling sites.

Well-dated plant remains from pile-dwellings in Slovenia indicate a continuous cultivation of six crops from at least about 4000-3100 cal BCE (Tolar 2018a; Tolar *et al.* 2011). Even older pile-dwellings in the area of the Ljubljansko barje are known, such as the site of Resnikov prekop, from 4600 cal BCE (Velušček 2006), but with so far no crop macro-remains from this period, likely due to the taphonomic effect of the fluctuating nearby river (Culiberg 2006). It is assumed that plant cultivation in the Ljubljansko barje must have already started in the 5th millennium BCE.

There is a gap in knowledge on the use of plants in Slovenia for the period 1700-1200 cal BCE. Plant macro-remains from this period are poorly preserved (they come from dry sites), and archaeobotanical studies are rare and lacking in information.

Date or time-span	Copper Age		
	4356-3995 cal BCE	4000 cal BCE	3300 cal BCE and 3100 cal BCE
Site	Ajdovska jama, cave cemetery; SE43	Strojanova voda, pile dwelling	Stare gmajne, pile dwelling
Taxon / measure of quantity	frequency	avg. concentration per 1 L of sediment in CL	avg. concentration per 1 L of sediment in CL
<i>Hordeum vulgare</i>	1326	155	102
<i>Triticum monococcum</i>	3049	116	24
<i>Triticum dicoccum</i>	792	142	150
<i>T. durum/turgidum/aestivum</i>	2		
<i>Cerealia</i>	82		
<i>Avena sativa</i>	2		
<i>Avena/Secale</i>			
<i>Avena/Bromus</i>			
<i>Secale cereale</i>			
<i>Papaver somniferum</i>		820	635
<i>Linum usitatissimum</i>		32	75
<i>Pisum sativum</i>			1
<i>Panicum miliaceum</i>			
<i>Sinapis arvensis</i>			
<i>Setaria italica</i>			
<i>Panicum/Setaria</i>			
<i>Lens culinaris</i>			
<i>Vicia faba</i> (fragments)	1		
<i>Lathyrus sativus</i>			
<i>Solanum</i> sp.			1
<i>Physalis alkekengi</i>		115	14
<i>Fragaria vesca</i>		226	92
<i>Rubus idaeus</i>			1
<i>Rubus fruticosus</i> agg.		375	60
<i>Cornus mas</i>		13	1
<i>Cornus sanguinea</i>		4	1
<i>Corylus avellana</i> (fragments)		3	6
<i>Juglans regia</i> (fragments)			
<i>Quercus</i> sp.		33	92
<i>Trapa natans</i>		6	3
<i>Sambucus</i> sp.		550	1
<i>Prunus</i> sp.			1
<i>Maloideae</i>		56	141
<i>Crateaeus</i> sp.		6	2
<i>Vitis vinifera sylvestris</i>		2	1
REFERENCES	Culiberg <i>et al.</i> 1992	Tolar 2018a	Tolar <i>et al.</i> 2011
Notes about the dates	date cited after the reference	radiocarbon and dendro-date	radiocarbon and dendro-date

Table 2. Overview of the archaeobotanical evidence through time from Slovenia.

Late Bronze Age			Early Iron Age		
1190-900 cal BCE	1190-900 cal BCE	1130-970 cal BCE		1000-400 cal BCE	
Dragomelj	Orehova vas	Grosuplje	sites in south-eastern Slovenia (7-12 in Fig. 1)	Ljubljana-Tribuna	Most na Soči
<i>absolute number of grains/seeds</i>	<i>absolute number of grains/seeds</i>	<i>absolute number of grains/seeds</i>	<i>absolute number of grains/seeds or, where presented as words, an estimate</i>	<i>absolute number of grains/seeds</i>	<i>weight or absolute number of grains/seeds</i>
160	1	2	a lot	76	
			some		
1				1	
			63	7	
81	17	2		60	56.63 g of porridge + 5 g of food remains
			9		
					55.37 g of porridge
86					
32			1		
			2		
1			1	3	
80	34	258	many	120	37.12 g
			many		
8			some	28	52.74 g of porridge
3					70 g of food remains
10	5	5	38	78	
1			6	3	5
				1	
				2	
				3	
				1	
				279	
			13	4	
			10	5	
10		2		8	4
					2
				6	
1			4	17	
				1	
1					
Tolar unpublished data	Tolar data published in Grahek 2015	Tolar unpublished data	Culiberg and Šercelj 1995	Tolar and Vojakovič unpublished data	Tolar 2018b
radiocarbon date	radiocarbon date	radiocarbon date		radiocarbon date	

Site	¹⁴ C AMS-date on single (individual) millet grains	Overall quantity of millet grains in the dated deposits
Dragomelj	1192-919 cal BCE (95.4%) [Poz-104925, Poz-104926, Poz-104927]	Altogether 14 grains in two buildings
Orehova vas	1193-923 cal BCE (95.4%) [Poz-104924, Poz-104985]	A total of 25 grains from a fireplace
Grosuplje*	1134-971 cal BCE (83.1%) [Poz-45566: 2880 ± 35 BP]	More than 200 grains in a pit ('pithos')

Table 3. Calibrated radiocarbon dates of carbonised millet grains from three Late Bronze Age sites in Slovenia (for the original data see Filipović et al. 2020; *details on the date for Grosuplje are kept in the Archive of the Institute of Archaeology ZRC SAZU).

Appearance of *Panicum miliaceum* in Slovenia

The earliest finds of millet grains from Slovenia are known from three sites; they were all radiocarbon dated (Table 3).

In addition to millet, the Late Bronze Age inhabitants of Dragomelj cultivated nine more cultivars: barley, emmer, other wheats (*Triticum* spp.), oat, rye (*Secale cereale*), pea, foxtail millet (*Setaria italica*), lentil (*Lens culinaris*) and faba bean. Wild plant gathering seems to have played a less important role, since only three wild plant taxa were recovered: hazel (*Corylus avellana*), elder (*Sambucus* sp.) and grape vine (*Vitis vinifera* ssp.) (Table 2). The sites of Orehova vas and Grosuplje show an absolute predominance of broomcorn millet (Table 2), signalling that it was already a very important, if not the main crop at that period.

After the appearance of millet

The Late Bronze Age and Early Iron Age can be considered as periods of agricultural expansion in Slovenia. The plant macro-remains from the Early Iron Age settlement at Most na Soči show that broomcorn millet and foxtail millet were already two of the main crops in that period. The charred remains of prepared food, i.e. porridge, from those two cultivars as well as from cereals (*Triticum* sp., wheat and *Avena/Secale*, oat/rye), were also found in two Early Iron Age houses at Most na Soči (Tolar 2018b). The find of charred crust resembling porridge has parallels in the find at the Late Bronze Age site of Stillfried, in Lower Austria (Kohler-Schneider 2001). There, the analysis showed a mixture of millet, barley and rye brome (*Bromus secalinus*) – a weed species that was probably tolerated, and therefore not removed, because of its nutritional value. The composition of the find from Most na Soči (Fig. 2) corresponds well to the one described from Stillfried. In addition to the remains of millets and wheats, legumes (i.e. faba bean) and two gathered plant taxa (hazelnut and walnut (*Juglans regia*)) were also recognised at Most na Soči (Table 2).

At all of the six Early Iron Age sites from southeastern Slovenia considered in this study (Fig. 1), systematic sampling and wet sieving of sediments was carried out, representing the second time this had been done in Slovenia (the first time was at the Palaeolithic cave site of Divje Babe I; see Introduction). Systematic excavations were carried out at these sites for more than ten years, starting in the early 1990s, under the supervision of J. Dular from the Institute of Archaeology ZRC SAZU, in Ljubljana (Dular and Tecco Hvala 2007). Colleagues from the Institute of Biology ZRC SAZU carried out the identification of plant macro-remains recovered with the 'initial flotation method' (Culiberg and Šercelj 1995), with which only larger and more resistant charred grains of cultivated plants were obtained (Dular and Tecco Hvala 2007, 30-31). They found that there are no significant differences between the crop spectra at the investigated Late Bronze Age and Early Iron Age sites in southeastern Slovenia. The species they identified were hulled and free-threshing wheats (*Triticum monococcum/dicoccum*, *T. aestivum/durum/turgidum*), barley, broomcorn millet, foxtail millet, flax, oat, rye, pea, faba bean, lentil and other Fabales (e.g. vetch (*Vicia* sp.)); they are all represented in both periods. The frequencies of plant remains differ between the periods, but the small sample and non-systematic recovery preclude any definite conclusions (Dular



Figure 2. Food remains ('porridge') from House 6 at Most na Soči made of foxtail millet (*Setaria italica*) and other cereals (after Tolar 2018b, 448, Fig. 2).

and Tecco Hvala 2007, 209). In any case, it could be stated that among the cereals at these sites, free-threshing wheats prevail, oat and rye are also present, but hulled wheats are almost absent (Table 2). Free-threshing wheat is more productive and highly nutritious, and its flour can be used for making leavened bread. Therefore, its increased presence may be related to a change in culinary practices (Megaloudi 2004, 155). Further, in comparison with hulled wheat, cultivation of free-threshing wheat requires a higher labour input and may reflect more intensive agricultural practices and perhaps also increased population density. Once harvested, however, its processing is easier and faster than that of other prehistoric cereal crops (Nesbitt 1995, 74). Among the gathered plants, only three taxa were recognised at Early Iron Age southeastern Slovenian sites (Table 2): blackberry (*Rubus fruticosus*), raspberry (*Rubus idaeus*) and elderberry (*Sambucus nigra*).

Tribuna is a multi-phase archaeological site in Ljubljana and, among the Early Iron Age sites discussed here, one of the most recently excavated (in 2008). It has layers from the Late Bronze Age (ca. 1100 cal BCE), from the Iron Age to the Roman period, and from modern times. Lots of sediment samples from all four phases were wet sieved, but they were subsequently dried, which caused some loss of fine and fragile plant macro-remains, a factor that should be taken into account when interpreting the archaeobotanical results. The Early Iron Age settlers at Tribuna consumed different kinds of cereals (barley and wheats), legumes (lentil, faba bean, grass pea (*Lathyrus sativus*) and pea) as well as foxtail and broomcorn millet. The last seems to have been very important (see Table 2). Additionally, ten possibly gathered plant taxa were recognised, but except for raspberry, in smaller quantities (Table 2).

Possible climatic changes in Slovenia and adjacent areas in the Bronze Age

Very few palaeoclimatological studies are available for Slovenia and environs. It can be assumed that there were significant differences in past climatic conditions between the regions south and north of the Alps as well as between different altitudes within the Alps, where vegetation is sensitive to climatic fluctuations. Badino *et al.* (2018) have recognised a cold phase in the period ca. 2000-1500 cal BCE in the western Italian Alps. They have reconstructed temperatures suggesting a temperature decrease of

1.8 °C compared with the previous period. On the other hand, glaciers in the Alps experienced a reduction in size (making them smaller than today), during most of the Holocene and until ca. 2000-1000 cal BCE (Ivy-Ochs *et al.* 2009; Solomina *et al.* 2015), indicating melting due to warmer conditions. Badino *et al.* (2018) also detected a warm peak around 1280-1150 cal BCE. In Italy, there are additional indicators of a warm and dry environment during the Bronze Age (*e.g.* Mercuri *et al.* 2012), but this may not apply to neighbouring Slovenia (Andrič *et al.* 2017) or, if it does, can only be proposed for lowland regions in eastern and southern Slovenia.

Few studies so far have focused on environmental changes in the southeastern Alps. A 12 m-long core from the central part of Lake Bohinj, in northwestern Slovenia, reflects environmental conditions and human-environment interactions over the past 6600 years (Andrič *et al.* 2020). It has been suggested that at ca. 6000 cal BP, Lake Bohinj was surrounded by mixed forest (of conifers and deciduous oak (*Quercus*)). Beech (*Fagus*) became dominant after ca. 1300 cal BCE. In the Bronze Age and especially in the Iron Age (1500-500 cal BCE), when the region was densely populated, anthropogenic clearing of forest is detected. It is evident that, around 1000 cal BCE, river floods were not as common as before, but the proportion of beech did not fall, which means there was no increase in aridity, unlike what was documented in the marine core from the central Adriatic (Mercuri *et al.* 2012). The pollen record from that core shows a gradual, irreversible trend towards increasing aridity from ca. 3700 cal BCE and less cool conditions from around 3100 cal BCE. The reduction in precipitation has also been evidenced before and during the Early Bronze Age (ca. 1900-1600 cal BCE; Mercuri *et al.* 2012).

Archaeological evidence: Harvesting tools from the Mesolithic to the Late Bronze Age

Harvesting tools appeared at the same time that agriculture emerged. The predecessor of the sickle is the harvesting knife. Such knives are comprised of flat or slightly curved bone or wooden handles with a groove into which quartz wedges are glued with resin (Fig. 3.1-5). A characteristic shine (known as sickle gloss; German: *Sichelglanz*) is visible on the working edges of the wedges, which proves that they were used as harvesting tools. That this shine is the result of friction that occurs when cutting grass or harvesting cereals has been confirmed through experiments (Steensberg 1943, 10 f.; Unger-Hamilton 1988; Juel Jensen 1996; Petru 1997, 97; Anderson 1999). The oldest preserved harvesting knives belong to the Mesolithic Natufian culture in Palestine (Fig. 3.1). Various shaped Neolithic harvesting knives are known from Egypt (Fig. 3.2), Asia Minor, Cyprus, the Balkans (Fig. 3.4), and pile-dwellings in Italy (Fig. 3.3) and Switzerland (Fig. 3.5) (Steensberg 1943, 126; Mellaart 1961, 45, Pl. 4a; Müller-Karpe 1968, 509; Rageth 1974, 193, Fig. 12, Pl. 103.1-3, Pl. 118.5; Egloff 1984, 64, Fig. 19; Perini 1987, 395, Fig. 186; Speck 1990, 262, Fig. 13; Ramseyer 2000, 218, Fig. 178; Gurova 2005).

The first Neolithic sickles, which have a concavely curved blade, originate from Mesopotamia. They were made of very hard, fired clay. The sharp edge is finely serrated (Fig. 3.6) (Steensberg 1943, 133). No completely preserved Neolithic sickles are known from central Europe. However, a reconstruction of the appearance of a Late Neolithic sickle is made possible by finds from two sites in northwestern Switzerland (Fig. 3.7). The stone wedge was found on the island of Werd, near Eschenz (Hasenfratz 1985, 110), and the wooden handle on the site of Niederwil-Egelsee, which belongs to the Pfyn culture (Pfahlbauquartett 2004, 49, cat. No. 45).

Sickles have been used in Egypt since at least the 1st Dynasty. A set of quartz wedges (Flinders Petrie 1891, 12, 54; Müller-Karpe 1968, 409) was inserted into a curved wooden handle, which may have consisted of one or more parts (Fig. 3.8-9).



Figure 3. Harvesting tools from the Mesolithic to the Late Bronze Age. 1. Kebara; 2. Fajum; 3. Lago di Ledro; 4. Karanovo; 5. Egozwill 3; 6. unknown site in Mesopotamia; 7. Eschenz-Island Werd and Niederwil-Egelsee; 8. Sakkara; 9. Kahun; 10. Barche di Solferino; 11. Fiavé; 12. Dunaújváros-Kosziderpadlás II; 13. Senomaty; 14. Balaton area (after: 1 – Mellaart 1961; 2, 5, 8 – Müller-Karpe 1968; 3 – Rageth 1974; 4 – Gurova 2005; 6 – Steensberg 1943; 7 – Pfahlbauquartett 2004; 9 – Flinders Petrie 1891; 10-11 – Perini 1987; 12 – Mozsolics 1967; 13 – Chvojka *et al.* 2017; 14 – Angeli and Neuninger 1964). For the scale of individual objects, see the cited literature.

Such sickles are also depicted in frescoes with harvest scenes on the walls of tombs (Müller-Karpe 1974, Pl. 40.8, Pl. 47, Pl. 49.9, Pl. 60.3, Pl. 75.3, Pl. 78.3, Pl. 81.3, Pl. 96.5). In northern Italy, sickles similar to the Egyptian ones are known; these are referred to as sickles *a mandibola* (Fig. 3.10) and sickles of the Fiavé type (Fig. 3.11). They were in use until the Middle Bronze Age (Perini 1987, 395, Figs. 185-186, 1988, 59).

At the beginning of the Middle Bronze Age, the first bronze sickles joined the sickles with stone blades. Bronze tanged sickles are typical for southeastern Europe at this time. As the Dunaújváros-Kosziderpadlás II and Senomaty hoards show, the first sickles had a flat tang (Fig. 3: 12-13). The sickle from the former hoard is one of the few made for use with the left hand, and the sickle from the latter was made for use with the right hand (Mozsolics 1967, 135, Pl. 49: 10; Chvojka *et al.* 2017, Pl. 78: 17). At the beginning of the Late Bronze Age, bronze sickles with two to six vertical ribs on the tang (Fig. 3: 14) began to appear in the hoards *en masse*. We encounter

them until the beginning of the Iron Age, and then they disappear. They are replaced by similarly shaped iron sickles with a flat tang. Whereas the number of tanged sickles from the Bronze Age reaches into the thousands, the number from the Early Iron Age is only a fraction of that. Whether this is due to the state of research or something else is a question that cannot be answered at the moment.

In Slovenia, the oldest flint pieces with gloss come from the Late Neolithic layers (5th millennium BCE) of the site of Moverná vas, in Bela krajina, located on a riverbank (Petru 1997, 89), and from the settlement of Dragomelj, near Ljubljana, located on a plain. At Dragomelj, up to 15% of the flint pieces show signs of having been used in harvesting (Petru 2005, 82, Fig. 4; Turk and Svetličič 2005; Turk and Turk 2019, Fig. 139). Of similar age are the three flint pieces with gloss from the settlement of Kamna gorica, near Ljubljana, also located on a plain (Petru pers. comm.).

From the Copper Age, we know of the flint pieces with gloss from the Trhlovca cave, near Divača (Petru 2004, 202), and from the pile-dwelling settlement of Hočevarica, in the Ljubljansko barje (Velušček 2004, 5, Fig. 34.1.34). Two flint pieces from the multi-period settlement of Col 1, situated on a plain near Podgračeno, in the Dolenjska region, are dated to the transition from the Neolithic to the Copper Age (Petru 2020, 104 no. 214, 105 no. 235). Due to the unclear stratigraphic situation, it is not known whether the flint piece with gloss from the Mala Triglavca cave, near Divača, dates to the Neolithic or to the Bronze Age (Petru 2004, 201).

Three completely preserved, left-handed metal sickles found in the Ljubljanica River belong to the Middle Bronze Age (Pavlin 2006, 79-76, Fig. 2.1a-3b). They can also be interpreted as gifts to the gods. No hoards are known from the Middle Bronze Age. In contrast, more than 60 hoards are known from the Late Bronze Age. Sickles, either entire or fragmented, are present in about three quarters of the hoards, and their combined number exceeds 300. Traces of hammering and sharpening prove that they were used primarily as harvesting tools. In addition to being found in hoards, they are found as individual finds both in and outside settlements (Teržan 1995). Particularly noteworthy is the Tribuna site, in Ljubljana. Among other things, at least three fragments of sickles were found there (Vojaković pers. comm.).

Summary and conclusions

Based on the available results from archaeological sites in Slovenia, the appearance of broomcorn millet and foxtail millet, together with oat, rye, lentil and faba bean, can be placed in the Late Bronze Age. In the Early Iron Age, the importance of millet increases, while the introduction of free-threshing wheat represents another agricultural innovation at this time. Just as emmer and einkorn are typical cereals of the Copper Age in the region, so millets are of the Late Bronze Age and, especially, of the Early Iron Age. Pulses were less common in the Copper Age and became more important also in the Late Bronze Age and Early Iron Age. Barley seems to have been equally important through all of these periods. Collected wild fruits and nuts were more important for the diet during the Copper Age than during the Late Bronze Age and Early Iron Age. However, the way in which archaeobotanical sampling/collecting, storing and sieving was done (if it was done at all) may have played an important role in the amount and type of results obtained, especially from excavations prior to 2007. Therefore, the identified remains only give a general impression of the staple crops used at the studied sites.

According to some recent reviews of the Bronze Age archaeobotanical finds from Europe (Stika and Heiss 2013; Filipović *et al.* 2020), broomcorn millet became a major (or even main) crop during the transition from the Middle to the Late Bronze Age. In Slovenia, the same phenomenon may be present, but the scarce archaeobotanical research, especially for the period between 2300-1200 BCE, cannot yet confirm the occurrence of millet before the 12th century BCE. The earliest finds of millet from neighbouring countries date from the Middle Bronze Age, namely to the 16th or 15th centuries BCE in Italy, Hungary, Croatia and to the 15th century BCE in Austria (Filipović *et al.* 2020). Both possible millet transition routes (via the Balkans or the Apennines or both; see Introduction) would thus allow for the possibility of an even earlier arrival of millet in Slovenia.

The crops introduced in the Late Bronze Age and later, in the Iron Age, were probably all used mainly for the preparation of porridge (see Fig. 2) and flat bread, and they are all quite resistant to unfavourable climatic conditions. For Slovenia, there is limited palaeoclimatic evidence for this period (Andrič *et al.* 2017, 2020). The pollen record from the marine core from the central Adriatic (Italy) shows a gradual, irreversible trend towards increasing aridity, less cool conditions and a reduction in precipitation before and during the Bronze Age (Mercuri *et al.* 2012), which could have been one of the reasons for the introduction of more drought-resistant crops in lowland (drier and warmer) regions in Slovenia.

Towards the end of the 14th century BCE, Europe experienced great changes in connection with the mass migrations of people. Central and eastern Slovenia were inhabited by people who belonged to the cultural sphere that stretched from western Hungary through Croatia to northern Bosnia. Large, Late Bronze Age settlements on the plains point to increased population density, which led to more intensive and efficient agricultural practices. At Late Bronze and Iron Age sites in Slovenia, cultivation of millets and free-threshing wheats increased, while that of hulled wheats decreased. Millets and free-threshing wheats are both more productive in comparison with hulled wheats. Although the cultivation of free-threshing wheat requires higher labour input (*i.e.* larger population), its processing is easier and faster. The greater presence of millets and free-threshing wheat may also be related to changes in culinary practices. Interesting research is going on at the moment involving analysis of ceramic forms in relation to food preparation, from various Slovenian archaeological sites dated to the Bronze Age and the Early Iron Age. Results to date reveal that some new forms of vessels (as well as changes in ceramic recipes) developed in the 12th-11th century BCE, which may be connected to changes or innovations in cooking techniques, perhaps linked with the appearance of millet. Later, in the Early Iron Age, these ceramic forms were used widely and regularly (Vinazza 2021).

Concerning the harvesting equipment in the studied region, bronze sickles with two to six vertical ribs on the tang began to appear at the beginning of the Late Bronze Age. We encounter them until the beginning of the Iron Age, and then they disappear. They are replaced by similarly shaped iron sickles with a flat tang.

Acknowledgements

We thank Dr. Lucija Grahek, Dr. Petra Vojaković, Dr. Peter Turk, Dr. Manca Vinazza and Barbara Hofman for information about archaeological research in progress and for some unpublished results. The project was financed by the Slovenian Research Agency, P6-0064 (B).

References

- Anderson, P.C., ed., 1999. *Prehistory of Agriculture: New Experimental and Ethnographic Approaches*. Monograph 40. Los Angeles: Cotsen Institute of Archaeology.
- Andrič, M., Sabatier, P., Rapuc, W., Ogrinc, N., Dolenc, M., Arnaud, F., von Grafenstein, U., Šmuc, A., 2020. 6600 years of human and climate impacts on lake-catchment and vegetation in the Julian Alps (Lake Bohinj, Slovenia). *Quaternary Science Reviews*, 227, 106043-106060. <https://doi.org/10.1016/j.quascirev.2019.106043>
- Andrič, M., Tolar, T., Toškan, B. 2016. *Okoljska arheologija in paleoekologija: palinologija, arheobotanika in arheozoologija*. Ljubljana: Založba ZRC.
- Andrič, M., Verbič, T., Lomax, J., Tolar, T., 2017. Človekov vpliv na okolje v prazgodovini: primer z obrežja Ljubljanske pri Špici (Ljubljana) [Embankment of the Ljubljana River at Špica (Ljubljana) and human impact on the environment in late prehistory]. *Arheološki vestnik*, 68, 479-498.
- Angeli, W. and Neuninger, H., 1964. Ein urnenfelderzeitlicher Depotfund aus der Gegend des Plattensees. *Mitteilungen der Anthropologischen Gesellschaft in Wien*, 93-94, 77-89.
- Badino, F., Ravazzi, C., Vallè, F., Pini, R., Aceti, A., Brunetti, M., Champvillair, E., Maggi, V., Maspero, F., Perego, R., Orombelli, G., 2018. 8800 years of high-altitude vegetation and climate history at the Rutor Glacier forefield, Italian Alps: evidence of middle Holocene timberline rise and glacier contraction. *Quaternary Science Reviews*, 185, 41-68. <https://doi.org/10.1016/j.quascirev.2018.01.022>
- Bogaard, A. and Styring, A., 2017. Plants, People and Diet in the Neolithic of Western Eurasia. In: J. Lee-Thorp and M.A. Katzenberg, eds., *The Oxford Handbook of the Archaeology of Diet*. Oxford Handbooks online. Oxford: Oxford University Press. <https://doi.org/10.1093/oxfordhb/9780199694013.013.45>
- Chvojka, O., Jiráň, L., Metlička, M., 2017. *Nové české depoty doby bronzové. Hromadné nálezy kovových předmětů učiněné do roku 2013*. České Budějovice: Jihočeská univerzita, Praha: Archeologický ústav AV ČR, Plzeň: Západočeské muzeum.
- Culiberg, M., 1984. Karpološke in ksilotomske raziskave kolišča na Partih [Carpological and xylotomic research of the pile dwelling at Parte]. *Poročilo o raziskovanju paleolita, neolita in eneolita v Sloveniji*, 12, 91-100.
- Culiberg, M., 1999. Palaeobotany in Slovene archaeology [Paleobotanika v slovenski arheologiji]. *Arheološki vestnik*, 50, 323-331.
- Culiberg, M., 2006. Rastlinski ostanki z arheološkega najdišča Resnikov prekop [Plant Remains from the Archaeological Site at Resnikov Prekop]. In: A. Velušček, ed., *Resnikov prekop – najstarejša naselbina na Ljubljanskem barju / Resnikov prekop, the oldest pile-dwelling settlement in the Ljubljansko barje*. Opera Instituti Archaeologici Sloveniae 10. Ljubljana: Založba ZRC, 129-132.
- Culiberg, M., 2007. Paleobotanične raziskave v Divjih babah I [Palaeobotanical research in Divje Babe I]. In: I. Turk, ed., *Divje babe I. Paleolitsko najdišče mlajšega pleistocena v Sloveniji. I. del: Geologija in paleontologija / Divje Babe I: Upper Pleistocene Palaeolithic site in Slovenia: Part I: Geology and Palaeontology*. Opera Instituti Archaeologici Sloveniae 13. Ljubljana: Založba ZRC, 177-184.
- Culiberg, M., 2011. Paleobotanične raziskave v paleolitskih najdiščih v Sloveniji [Palaeobotanical research at Palaeolithic sites in Slovenia]. In: B. Toškan, ed., *Drobci ledenodobnega okolja / Fragments of Ice Age environments*. Opera Instituti Archaeologici Sloveniae 21. Ljubljana: Založba ZRC, 219-234.
- Culiberg, M. and Šercelj, A., 1995. Karpološke in antrakotomske analize iz prazgodovinskih višinskih naselij na Dolenjskem [Karpologische und anthrakotomische Analysen aus den vorgeschichtlichen Höhensiedlungen in Dolenjsko]. *Arheološki vestnik*, 46, 169-176.
- Culiberg, M., Horvat, M., Šercelj, A., 1992. Karpološke in antrakotomske analize rastlinskih ostankov iz neolitske jamske nekropole Ajdovska jama [Karpologische

- und anthrakotomische Analysen pflanzlicher Überreste aus der Höhlennekropole Ajdovska jama]. *Poročilo o raziskovanju paleolita, neolita in eneolita v Sloveniji*, XX, 111-126.
- Deschmann, K., 1875. Die Pfahlbautenfunde auf dem Laibacher Moore. *Verhandlungen der k.k. geologischen Reichsanhalt*, 15, 275-284.
- Deschmann, C., 1878. Ueber die vorjährigen Funde im Laibacher Pfahlbau. *Mitteilungen der Anthropologischen Gesellschaft in Wien*, 3-4, Band VIII, 1-20.
- Dular, J., 2013. *Severovzhodna Slovenija v pozni bronasti dobi / Nordostslowenien in der späten Bronzezeit*. Opera Instituti Archaeologici Sloveniae 27. Ljubljana: Založba ZRC.
- Dular, J. and Tomanič Jevremov, M., 2010. *Ormož: utrjeno naselje iz pozne bronaste in starejše železne dobe / Ormož: befestigte Siedlung aus der späten Bronze- und der älteren Eisenzeit*. Opera Instituti Archaeologici Sloveniae 18. Ljubljana: Založba ZRC.
- Dular, J., Križ, B., Svoljšak, D., Tecco Hvala, S., 1991. Utrjena prazgodovinska naselja v Mirenski in Temeniški dolini [Befestigte prähistorische Siedlungen in der Mirenska dolina und der Temeniška dolina]. *Arheološki vestnik*, 42, 65-198.
- Dular, J., Šavel, I., Tecco Hvala, S., 2002. *Bronastodobno naselje Oloris pri Doljem Lakošu / Bronzezeitliche Siedlung Oloris bei Dolnji Lakoš*. Opera Instituti Archaeologici Sloveniae 5. Ljubljana: Založba ZRC.
- Dular, J. and Tecco Hvala, S., 2007. *South-Eastern Slovenia in the Early Iron Age. Settlement – Economy – Society / Jugovzhodna Slovenija v starejši železni dobi. Poselitev – gospodarstvo – družba*. Opera Instituti Archaeologici Sloveniae 12. Ljubljana: Založba ZRC.
- Egloff, M., 1984. A l'aube du 'design': les manches de faucilles du Bronze final. *Helvetica Archaeologica*, 15, 51-66.
- Flinders Petrie, W.M., 1891. *Illahun, Kahun and Gurob: 1889-90*. London: David Nutt.
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Kloß, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A.M., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M.S., Reed, K., Rennwanz, J., Stika, H.-P., Stobbe, A., Tolar, T., Wasylukowa, K., Wiethold, J., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Gabrovec, S., ed., 1994. *Stična 1: naselbinska izkopavanja / Stična 1: Siedlungsausgrabungen*. Katalogi in monografije 28. Ljubljana: Narodni muzej.
- Grahek, L., ed., 2015. *Orehova vas*. Arheologija na avtocestah Slovenije 46. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije. Available from: https://www.zvkds.si/sites/www.zvkds.si/files/uploads/files/publication/046_orehova_vas.pdf
- Gurova, M., 2005. Elements de fauciles neolithiques en silex de la Bulgarie: evidence et contexte. *Archaeologia Bulgarica*, 9 (1), 1-14.
- Guštin, M., ed., 2005. *Prvi poljedelci: savska skupina lengyelske kulture / First farmers: the Sava group of the Lengyel culture*. Annales Mediterranea. Koper: Založba Annales.
- Guštin, M. and Tomaž, A., 2016. *Zatak pri Lendavi*. Arheologija na avtocestah Slovenije 51. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije. Available from: https://www.zvkds.si/sites/www.zvkds.si/files/uploads/files/publication/zatak_pri_lendavi_i.pdf
- Guštin, M., Tiefengraber, G., Pavlovič, D., Zorko, M., 2017. *Nova tabla pri Murski Soboti. Prazgodovina*. Arheologija na avtocestah Slovenije 52. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije. Available from: https://www.zvkds.si/sites/www.zvkds.si/files/uploads/files/publication/nova_tabla_-_prazgodovina_web.pdf
- Hasenfratz, A., 1985. *Eschenz, Insel Werd 2. Das jungneolithische Schichtpaket III*. Zürcher Studien zur Archäologie. Zürich: Abteilung Ur- und Frühgeschichte der Universität Zürich.

- Horvat, M., 1989. *Ajdovska jama pri Nemški vasi*. Ljubljana: Znanstveni inštitut Filozofske fakultete.
- Ihde, C., 1995. Die Elefantendarstellungen der Hvar-Lisičići-Kultur und das Problem ihrer Herleitung / Upodobitve slonov v hvarski kulturi in problem njihovega izvora. *Arheološki vestnik*, 46, 53-88.
- Ivy-Ochs, S., Kerschner, H., Maisch, M., Christl, M., Kubik, P.W., Schlüchter, C., 2009. Latest Pleistocene and Holocene glacier variations in the European Alps. *Quaternary Science Reviews*, 28 (21), 2137-2149. <https://doi.org/10.1016/j.quascirev.2009.03.009>
- Jeraj, M., 2004. Paleobotanične raziskave na kolišču Hočevarica [Palaeobotanical analyses of the Hočevarica pile dwelling]. In: A. Velušček, ed., *Hočevarica-eneolitsko kolišče na Ljubljanskem barju / An Eneolithic pile dwelling in the Ljubljana marsh*. Opera Instituti Archaeologici Sloveniae 8. Ljubljana: Založba ZRC, 56-64.
- Jeraj, M., Velušček, A., Jacomet, S., 2009. The diet of Eneolithic (Copper Age, fourth millennium cal B.C.) pile dwellers and the early formation of the cultural landscape south of the Alps: a case study from Slovenia. *Vegetation History and Archaeobotany*, 18, 75-89. <https://doi.org/10.1007/s00334-008-0144-z>
- Juel Jensen, H., 1996. TRB sickles and early Danish agriculture: a view from the microscope [Srpi culture lijakastih čaš in zgodnje dansko kmetovanje: pogled skozi mikroskop]. *Poročilo o raziskovanju paleolitika, neolitika in eneolitika v Sloveniji: Neolitske študije*, 23, 129-153.
- Kerman, B., 2011a. *Kotare-Baza pri Murski Soboti*. Arheologija na avtocestah Slovenije 17. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije. Available from: <https://www.zvkds.si/sites/www.zvkds.si/files/uploads/files/publication/kotare1.pdf>
- Kerman, B., 2011b. *Kotare-Krogi pri Murski Soboti*. Arheologija na avtocestah Slovenije 20. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije. Available from: https://www.zvkds.si/sites/www.zvkds.si/files/uploads/files/publication/20_kotare-krogi_pri_murski_soboti.pdf
- Kerman, B., 2011c. *Pod Kotom – sever pri Krogu*. Arheologija na avtocestah Slovenije 24. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije. Available from: https://www.zvkds.si/sites/www.zvkds.si/files/uploads/files/publication/24_pod_kotom-sever_pri_krogu.pdf
- Kohler-Schneider, M., 2001. *Verkohlte Kultur- und Wildpflanzenreste aus Stillfried an der March als Spiegel spätbronzezeitlicher Landwirtschaft im Weinviertel, Niederösterreich*. Wien: Verlag der Österreichischen Akademie der Wissenschaften.
- Korošec, J., 1953. Nova kolišča na Ljubljanskem barju: poročilo [New pile-dwellings on the Ljubljana moor: a report]. *Arheološki vestnik*, 4 (2), 256-263.
- Korošec, P. and Korošec, J., 1969. *Najdbe s koliščarskih naselbin pri Igu na Ljubljanskem barju / Fundgut der Pfahlbausiedlungen bei Ig am Laibacher Moor*. Arheološki katalogi Slovenije 3. Ljubljana: Mladinska knjiga.
- Megaloudi, F., 2004. Agriculture in mainland Greece at the Protogeometric period: a view from the archaeobotanical remains. *Eylimenh*, 5, 151-160.
- Mellaart, J., 1961. Excavations at Hacilar: fourth preliminary report, 1960. *Anatolian Studies*, 11, 39-75.
- Mercuri, A.M., Bandini Mazzanti, M., Torri, P., Vigliotti, L., Bosi, G., Florenzano, A., Olmi, L., Massamba N'siala, I., 2012. A marine/terrestrial integration for mid-late Holocene vegetation history and the development of the cultural landscape in the Po Valley as a result of human impact and climate change. *Vegetation History and Archaeobotany*, 21, 353-372. <https://doi.org/10.1007/s00334-012-0352-4>
- Mozsolics, A., 1967. *Bronzefunde des Karpatenbeckens: Depotfundhorizonte von Hajdúsámson und Kosziderpadlás*. Budapest: Akadémiai Kiadó.
- Müller-Karpe, H., 1968. *Handbuch der Vorgeschichte 2: Jungsteinzeit*. München: Beck.
- Müller-Karpe, H., 1974. *Handbuch der Vorgeschichte 3: Kupferzeit*. München: Beck.

- Nesbitt, M., 1995. Plants and people in ancient Anatolia. *Biblical Archaeologist*, 58 (2), 68-81.
- Pavlin, P., 2006. Srednjebronastodobni levoročni jezičastoročajni srpi in meč iz Ljubljani-
ca [Middle Bronze Age left-handed sickles and a sword from the Ljubljani-
ca River]. *Arheološki vestnik*, 57, 69-83.
- Pavlin, P., 2015. *Pod Kotom – cesta pri Krogu*. Arheologija na avtocestah Slovenije 48. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije. Available from: https://www.zvkds.si/sites/www.zvkds.si/files/uploads/files/publication/48_pod_kotom-cesta_pri_krogu.pdf
- Perini, R., 1987. *Scavi archeologici nella zona palafitticola di Fiavé-Carera 2: campagne 1969-1976: resti della cultura materiale: metallo – osso – litico – legno*. Patrimonio storico e artistico del Trento 9. Trento: Provincia autonoma di Trento Servizio Beni Culturali.
- Perini, R., 1988. Le suppellettili lignee nell'Età del Bronzo Sudalpina e la relativa cronologia. In: R. Perini, ed., *Archeologia del legno: documenti dell'Età del Bronzo dall'area Sudalpina*. Quaderni della Sezione archeologica del Museo provinciale d'arte 4. Trento: Provincia autonoma di Trento Servizio Beni Culturali, 53-64.
- Petru, S., 1997. Metode preiskav sledov uporabe na kamenih orodjih iz najdišč Zakajeni spodmol in Mala Triglavca [Methods of research on use wear on the stone tools from the sites of Zakajeni spodmol and Mala Triglavca]. *Poročilo o raziskovanju paleolitika, neolitika in eneolitika v Sloveniji: Neolitske študije*, 24, 79-97.
- Petru, S., 2004. Usewear analysis of Mesolithic and Neolithic stone tools from Mala Triglavca, Trhlovca and Pupičina peč. *Poročilo o raziskovanju paleolita, neolita in eneolita v Sloveniji: Neolitske študije*, 31, 199-204. <https://doi.org/10.4312/dp.31.14>
- Petru, S., 2005. Kamnita orodja s prazgodovinske naselbine v Dragomlju [Stone tools of the prehistoric settlement in Dragomelj]. In: M. Guštin, ed., *Prvi poljedelci: savska skupina lengyelske kulture / First farmers: the Sava group of the Lengyel culture*. Annales Mediterranea. Koper: Založba Annales, 81-85.
- Petru, S., 2020. Analiza sledi uporabe na kamnitih orodjih. In: M. Horvat, ed., *Col 1 pri Podgračnem*. Arheologija na avtocestah Slovenije 84. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije, 98-109.
- Pfahlbauquartett, 2004. *Pfahlbauquartett: 4 Museen präsentieren 150 Jahre Pfahlbau-Archäologie*. Konstanz: Archäologisches Landesmuseum Baden-Württemberg; Frauenfeld: Amt für Archäologie des Kantons Thurgau.
- Rageth, J., 1974. Der Lago di Ledro im Trentino und seine Beziehungen zu den alpinen und mitteleuropäischen Kulturen. *Bericht der Römisch-Germanischen Kommission*, 55, 73-359.
- Ramseyer, D., ed., 2000. *Muntelier – Fischergässli: un habitat néolithique au bord du lac de Morat (3895 à 3820 avant J.-C.)*. Archéologie fribourgeoise 15. Fribourg: Editions Universitaires.
- Schmid, W., 1910. Archäologischer Bericht aus Krain: II: der Pfahlbau von Notranje gorice am Laibacher Moor. *Jahrbuch für Altertumskunde*, 4, 90-110.
- Schmid, W., 1915. Die Ringwälle des Bachergebietes. *Mitteilungen der Prähistorischen Kommission der Kaiserliche Akademie der Wissenschaften*, 2 (3), 229-305.
- Solomina, O.N., Bradley, R.S., Hodgson, D.A., Ivy-Ochs, S., Jomelli, V., Mackintosh, A.N., Nesje, A., Owen, L.A., Wanner, H., Wiles, G.C., Young, N.E., 2015. Holocene glacier fluctuations. *Quaternary Science Reviews*, 111, 9-34. <https://doi.org/10.1016/j.quascirev.2014.11.018>
- Speck, J., 1990. Zur Siedlungsgeschichte des Wauwilermooses. In: M. Höneisen, ed., *Die ersten Bauern: Pfahlbaufunde Europas 1*. Zürich: Schweizerisches Landesmuseum, 255-270.
- Steensberg, A., 1943. *Ancient Harvesting Implements: a Study in Archaeology and Human Geography*. Nationalmuseets Skrifter, Arkæologisk-Historisk Række 1. København: I kommission hos Gyldendalske boghandel & Nordisk forlag.

- Stika, H.-P. and Heiss, A.G., 2013. Plant cultivation in the Bronze Age. In: H. Fokkens and A. Harding, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 340-361.
- Svoljšak, D. and Dular, J., 2016. Železnodobno naselje Most na Soči: gradbeni izvidi in najdbe / *The Iron Age settlement at Most na Soči: settlement structures and small finds*. Opera Instituti Archaeologici Sloveniae 33. Ljubljana: Inštitut za arheologijo and Založba ZRC.
- Šavel, I., 2009. *Pod Kotom – jug pri Krogu*. Arheologija na avtocestah Slovenije 7. Ljubljana: Zavod za varstvo kulturne dediščine Slovenije. Available from: https://www.zvkds.si/sites/www.zvkds.si/files/uploads/files/publication/007_pod_kotom-jug_pri_krogu.pdf
- Šercelj, A., 1981-1982. Pomen botaničnih raziskav na koliščih Ljubljanskega barja [The importance of botanical research at the pile-dwellings of Ljubljansko barje]. *Poročilo o raziskovanju paleolita, neolita in eneolita v Sloveniji*, IX-X, 101-106.
- Šinkovec, I., 1996. Posamezne kovinske najdbe bakrene in bronaste dobe [Individual metal finds from the Eneolithic and Bronze Ages]. In: B. Teržan, ed., *Depojske in posamezne kovinske najdbe bakrene in bronaste dobe na Slovenskem / Hoards and individual metal finds from the Eneolithic and Bronze Ages in Slovenia 2*. Katalogi in monografije 30. Ljubljana: Narodni muzej, 125-163.
- Šmit, Ž., 2004. Preiskava eneolitskih metalurških sledov s Hočevarice z metodo PIXE [Investigation of copper metallurgy at Hočevarica using the PIXE method]. In: A. Velušček, ed., *Hočevarica: Eneolitsko kolišče na Ljubljanskem barju / Hočevarica: an Eneolithic pile dwelling in the Ljubljansko barje*. Opera Instituti Archaeologici Sloveniae 8. Ljubljana: Inštitut za arheologijo and Založba ZRC, 69-71.
- Teržan, B., ed., 1995. *Depojske in posamezne najdbe bakrene in bronaste dobe na Slovenskem / Hoards and individual finds from the Eneolithic and Bronze Ages in Slovenia*. Katalogi in monografije 29. Ljubljana: Narodni muzej.
- Tolar, T., 2018a. Primerjava različnih metod vzorčenja in priprave arheobotaničnih vzorcev z eneolitskih kolišč Strojanova voda in Maharski prekop na Ljubljanskem barju [Comparison of different sampling and treatment methods in order to reconstruct plant economies at the Eneolithic pile-dwellings of Strojanova voda and Maharski prekop in the Ljubljana marshes]. *Arheološki vestnik*, 69, 461-498.
- Tolar, T., 2018b. Ostanke prehranskih rastlin iz hiš 6 in 15A z Mosta na Soči [Edible plant remains from Houses 6 and 15A at Most na Soči]. In: J. Dular and S. Tecco Hvala, eds., *Železnodobno naselje Most na Soči. Razprave / The Iron Age settlement at Most na Soči: Treatises*. Opera Instituti Archaeologici Sloveniae 34. Ljubljana: Založba ZRC, 445-452. <https://doi.org/10.3986/9789610501091>
- Tolar, T., Jacomet, S., Velušček, A., Čufar, K., 2010. Recovery techniques for water-logged archaeological sediments: a comparison of different treatment methods for samples from Neolithic lake shore settlements. *Vegetation History and Archaeobotany*, 19, 53-67. <https://doi.org/10.1007/s00334-009-0221-y>
- Tolar, T., Jacomet, S., Velušček, A., Čufar, K., 2011. Plant economy at a Late Neolithic lake dwelling site in Slovenia at the time of the Alpine Iceman. *Vegetation History and Archaeobotany*, 20, 207-222. <https://doi.org/10.1007/s00334-010-0280-0>
- Tolar, T., Jacomet, S., Velušček, A., 2016. Cereal chaff used as temper in loom-weights: new evidence from a Slovenian Eneolithic pile-dwelling site (ca. 3100 cal BC). *Vegetation History and Archaeobotany*, 25, 291-301. <https://doi.org/10.1007/s00334-015-0549-4>
- Toškan, B. and Dirjec, J., 2004. Hočevarica: analiza ostankov makrofavne [Hočevarica: an analysis of macrofauna remains]. In: A. Velušček, ed., *Hočevarica: eneolitsko kolišče na Ljubljanskem barju / Hočevarica: an Eneolithic pile dwelling in the Ljubljansko barje*. Opera Instituti Archaeologici Sloveniae 8. Ljubljana: Inštitut za arheologijo and Založba ZRC, 76-132.

- Turk, I., ed., 2007. *Divje babe I. Paleolitsko najdišče mlajšega pleistocena v Sloveniji. I. del: Geologija in paleontologija* / *Divje Babe I: Upper Pleistocene Palaeolithic site in Slovenia: Part I: Geology and Palaeontology*. Opera Instituti Archaeologici Sloveniae 13. Ljubljana: Založba ZRC,
- Turk, P. and Svetličič, V., 2005. Neolitska naselbina v Dragomlju [The Neolithic settlement in Dragomelj]. In: M. Guštin, ed., *Prvi poljedelci: savska skupina lengyelske culture / First farmers: the Sava group of the Lengyel culture*. Annales Mediterranea. Koper: Založba Annales, 65-79.
- Turk, P. and Turk, M., 2019. *Najstarejše zgodbe s stičišča svetov*. Ljubljana: Narodni muzej Slovenije.
- Turk, P., Istenič, J., Knific, T., Nabergoj, T. eds., 2009. *The Ljubljanica – a river and its past*. Ljubljana: Narodni muzej Slovenije.
- Unger-Hamilton, R., 1988. *Method in Microwear Analysis: Prehistoric Sickles and other Stone Tools from Arjoune, Syria*. British Archaeological Reports International Series S435. Oxford: BAR Publishing.
- Vinazza, M., 2021. *Lončarstvo pozne bronaste in starejše železne dobe v zahodni Sloveniji*. Doctoral dissertation, University of Ljubljana. <https://repozitorij.uni-lj.si/IzpisGradiva.php?id=127637&lang=slv&prip=rup:9087111:d3>
- Velušček, A., 2004. Terenske raziskave, stratigrafija in najdbe [Field research, stratigraphy and the material finds]. In: A. Velušček, ed., *Hočevarica: eneolitsko kolišče na Ljubljanskem barju / Hočevarica: an Eneolithic pile dwelling in the Ljubljansko barje*. Opera Instituti Archaeologici Sloveniae 8. Ljubljana: Inštitut za arheologijo and Založba ZRC, 33-55.
- Velušček, A., ed., 2006. *Resnikov prekop – najstarejša naselbina na Ljubljanskem barju / Resnikov prekop, the oldest pile-dwelling settlement in the Ljubljansko barje*. Opera Instituti Archaeologici Sloveniae 10. Ljubljana: Založba ZRC.
- Velušček, A., 2009. Koliščarska naselbina Stare gmajne in njen čas [The Stare gmajne pile-dwelling settlement and its era]. In: A. Velušček, ed., *Koliščarska naselbina Stare gmajne in njen čas: Ljubljansko barje v 2. polovici 4. tisočletja pr. Kr. / Stare gmajne pile-dwelling settlement and its era: the Ljubljansko barje in the 2nd half of the 4th millennium BC*. Opera Instituti Archaeologici Sloveniae 16. Ljubljana: Inštitut za arheologijo and Založba ZRC, 11-34.
- Velušček, A. and Greif, T., 1998. Talilnik in livarski kalup z Maharskega prekopa na Ljubljanskem barju [A crucible and casting mould from Maharski prekop in the Ljubljana marshes]. *Arheološki vestnik*, 49, 31-53.
- Vojaković, P., Novšak, M., Žerjal, T., Verbič, T., Krajšek, J., Hrustel, J., 2011. *Poročilo o predhodnih arheoloških raziskavah na lokaciji Ljubljana – stanovanjska soseska Tribuna* [Report on the previous archaeological research at the location Ljubljana – residential neighbourhood Tribuna]. Ljubljana: Arhej. <http://www.arhej.com/datoteke/Pdf/porocilo-tribuna-08-za-objavo-koncna.pdf>

On the 'ancient' evidence for *Panicum miliaceum* and *Vicia faba* in central Germany (primarily Saxony-Anhalt)

Monika Hellmund

Abstract

Broomcorn millet was domesticated in Asia, and not in the region known as the Fertile Crescent, in the Middle East. Archaeobotanical finds of *Panicum miliaceum* (broomcorn millet) in central Europe have been reported to date as early as the Neolithic; dozens of archaeological sites with finds of *P. miliaceum* dated archaeologically to the Neolithic are mentioned in the literature. AMS radiocarbon dating enables investigation of trace amounts of organic material, such as grains of *P. miliaceum*, to accurately date them and either confirm or refute the archaeological dating. This chapter discusses current research concerning broomcorn millet in central Germany, primarily the state of Saxony-Anhalt. It discusses sites with supposed Neolithic finds of *P. miliaceum* and sites with Late Bronze Age, Pre-Roman Iron Age or Roman period finds. It also discusses the difficulty of separating features of different ages on multi-period archaeological sites. Sites with single finds of broomcorn millet are much more common than those that yielded mass finds. In many cases, *P. miliaceum* and *Vicia faba* were found together in archaeological deposits. The radiocarbon dates prove that *P. miliaceum* became a staple crop during the Late Bronze Age, and they do not support a Neolithic date for the introduction of either *P. miliaceum* or *V. faba* in Saxony-Anhalt.

Introduction

Germany is often mentioned in the context of the earliest finds of broomcorn millet in Europe. In 2006-2007, a new permanent exhibition was prepared and installed at the Landesmuseum für Vorgeschichte (State Museum for Prehistory) in Halle (Saale), in central Germany (Meller 2012). It included the crop species of the earliest (Early Neolithic) farmers in the region, both as herbarium specimens and as carbonised grains from archaeological sites. This exhibition was the impetus to re-investigate

finds of ancient crops like *P. miliaceum* from the sites of the Neolithic and younger periods in Saxony-Anhalt in the last 15 years.

In sites of the Lusatian culture (Late Bronze Age to Pre-Roman Iron Age), lumps of charred grains of broomcorn millet have been preserved. Additionally, some mass finds of broomcorn millet are known from pits in settlement sites from the Late Bronze Age and the Iron Age in Saxony-Anhalt. Stray finds of grains exist as well. Both charred lumps and mass finds of broomcorn millet from eight features as well as individual finds of *P. miliaceum* from 15 settlement pits were the focus of the radiocarbon dating.

Until recently, *P. miliaceum* was said to have first appeared in Europe in the Neolithic (Bertsch and Bertsch 1949; Schultze-Motel 1994; Kroll 2001). It was said to be of minor importance as a crop because only single finds were present in assemblages (Willerding 1980; Kreuz 1990; Van Zeist *et al.* 1991; Körber-Grohne 1994; Zohary and Hopf 1994; Lünig 2000). Hunt *et al.* (2008) compiled data on 41 published sites with ancient finds of *P. miliaceum* and *Setaria italica* (foxtail millet) dated to before 5000 cal BCE in the Old World. The authors emphasised the necessity of providing detailed descriptions of morphological criteria and of the archaeological features that yielded the finds. Motuzaite-Matuzeviciute *et al.* (2013) note that only radiocarbon dating of the archaeobotanical material in question can elucidate when broomcorn millet became a staple crop in central Europe. *P. miliaceum* was the dominant cultivated millet species in the region investigated here. In contrast, *S. italica* has only rarely been proven.¹

The wild progenitor of *P. miliaceum* has never been identified, and it may no longer exist (Miller *et al.* 2016). Some potential domestication regions in Asia have been discussed (Hegi 1979; Geisler 1991). The wild progenitor of *S. italica* is considered to be *Setaria viridis*; this latter species is widespread in central Asia. The region of domestication of *S. italica* is not known either.

Early agriculture of *P. miliaceum* in Asia

Early finds of both crops are documented from the early 6th millennium BCE in northern China, near the Yellow River, and in other regions of northeastern China (Hunt *et al.* 2008). Some authors suggest that this area was a centre for dry-land agriculture and initial millet domestication (Ventresca Miller and Makarewicz 2019). Liu *et al.* (2009) argued that early farming in northern China featured the cultivation of both broomcorn millet and foxtail millet, but there are only a few finds of millet grains and even fewer that are radiocarbon dated. At the Early Neolithic sites in Xinglonggou, the upper Liao River region in northeastern China, along the eastern edge of Inner Mongolia, 1400 grains of *Panicum miliaceum* and 60 grains of *Setaria italica* have been excavated, and remains of *P. miliaceum* were directly dated by AMS to ca. 5720 to 5660 cal BCE (7670-7610 cal BP; Zhao 2011). Zhao states that these caryopses were of domesticated plant species; in contrast, Stevens and Fuller (2017) note that the ‘Xinglonggou *Panicum* grains are small and consistent with an early pre-domestication cultivation stage, not domestication’. Li *et al.* (2020) compiled data on directly dated broomcorn millet grains to reconstruct millet agriculture dispersal from northeastern China to the Russian Far East around the middle of the 4th millennium BCE.

Genetic investigations of recent *P. miliaceum* and *P. ruderales* suggest there may be two independent domestication centres in China, one on the Loess Plateau (cluster A) and one in northeastern China (cluster B). The routes of westward spread

1 The archive of State Office of Heritage Management and Archaeology Saxony-Anhalt in Halle (Saale) mentions a mass find of perhaps *Setaria italica* from the site of Gräfenhainichen, county of Wittenberg. The material has not yet been re-located and the find thus remains unconfirmed.

could have been the prehistoric 'Oasis Route' for cluster A and the 'Steppe Route' for cluster B (Xu *et al.* 2019). Along the migration trail from northeastern China to Europe, the 'Inner Asian Mountain Corridor' has to be overcome (Miller *et al.* 2016). Leipe *et al.* (2019, 2) suggest that the '*most probable route connecting the Upper Yellow River and the western central Asian steppes runs along the southwestern slopes of the Mongolian Altai and the southern Mongolian Plateau*'. Broomcorn millet was present on the western edges of the Eurasian Steppe in southeastern Kazakhstan by 2200 cal BCE (Frachetti *et al.* 2010). Filipović *et al.* (2020) documented the current state of the art on the introduction of broomcorn millet in central Europe, based on new and previously produced radiocarbon dates. Recent dating of charred grains of broomcorn millet indicates that the earliest presence in central and southeastern Europe was in the 16th century BCE.

Cultivation of broomcorn millet: Ecological and climatic demands

At present, broomcorn millet is seldom cultivated in central Europe. The crop has a moderate demand on soil nutrients and low water requirements; it tolerates drought but does not like cold or wet soils. Uniquely, this crop can be cultivated in very different climatic conditions. Varieties of broomcorn millet require a very short growing period and mature quickly, even at higher altitudes (Geisler 1991). In central Europe, the growing period is stated to be about 100 days, from May to August or at the latest from mid-June to September. In some other regions of the world, the vegetation period amounts to 60-90 days (Zohary *et al.* 2012) or even just 30-45 days (Frachetti *et al.* 2010). In contrast to those of other cereals, grains within the panicle of *P. miliaceum* do not all ripen at the same time; the grains at the top ripen first. The panicle is harvested by hand or with a sickle (Steinbrück 1908). A minimum temperature of 10 °C is necessary for germination, and the seedlings have a high frost sensitivity (Mehnert *et al.* 1964).

Broomcorn millet was the first C4 plant to be introduced in central Europe. In contrast to most of the other important world crops, which are characterised by C3-photosynthesis, *P. miliaceum* is characterised by C4-photosynthesis, which gives it an advantage in more arid climates, where water is scarce. *P. miliaceum* leaves a distinct isotopic signal in human and animal bones when used for either human nutrition or as fodder for domestic animals (*e.g.* Tafuri *et al.* 2009). Millet grains have a well-balanced nutritional composition (Bray 1981). The grains can be cooked like rice or consumed as a porridge or in soups, and they can be baked into a flatbread. They are used for human diet as well as for animal feed. It can be assumed that in the past the non-edible residues of the plant were fed to domestic animals.

Material and methods

Archaeological finds of broomcorn millet from already published and unpublished sites in Saxony-Anhalt were radiocarbon dated, with a focus on extant grains of *P. miliaceum* or accompanying cereal grains, preferably originating from large deposits (Table 2). Sometimes the carbonised grains of broomcorn millet themselves were dated, and wherever possible, five or more grains were dated. Because of the very low content of carbon in grains of broomcorn millet, sometimes the co-occurring grains of larger cereals from the same deposit were included in the sample for dating, based on the assumption that other grains in the same sample of stored grains, such as *Hordeum vulgare* or *Triticum dicoccum*, or pulses, such as *Vicia faba*, must be of the same age as the grains of broomcorn millet.



Figure 1. Locations of the sites in Saxony-Anhalt and surrounding regions in central Germany mentioned in the text. Dating (assumed or confirmed): Late Bronze Age, Iron Age, Roman period and unknown/recent (map by B. Parsche, LDA Halle (Saale), based on a design by the author. Map base: D 1000, Bundesamt für Kartographie und Geodäsie, Nr. 01/2002, 1:2,000,000).

Results

The radiocarbon dates of contexts with broomcorn millet in Saxony-Anhalt and the immediate surroundings are described here. The location of the sites mentioned in the text is shown on the map in Figure 1. Unless otherwise mentioned, the sites are located in the German federal state of Saxony-Anhalt. An overview of archaeological phases, cultures and periods used in central Germany from the Bronze Age until the Early Middle Age, together with the placement of the sites in Saxony-Anhalt mentioned in the text, is presented in Table 1. The radiocarbon dates and the calibrated ages are compiled in Table 2. The calibrations were carried out with OxCal v4.4.2² based on the IntCal20 atmospheric curve (Bronk Ramsey 2009; Reimer *et al.* 2020). The calibrated timespans are displayed in multi-plots (Figs. 2-5). Photos of the examined grains and lumps of broomcorn millet are presented in Figure 6.

² <https://c14.arch.ox.ac.uk/oxcal.html>

Temporal indicator		General period	Phase according to regional chronology*	Cultural attribution	Sites in Saxony-Anhalt
410	CE	Early Middle Ages	FMA	Thuringians	
165	CE	Period of Germanic Migration Roman Period	RKZ C-VWZ	Germanic tribes	Wolmirstedt Elbeu Klötze Osterwieck Quedlinburg
60	BCE		Lt D2-RKZ B		
150	BCE	Late Iron Age (La Tène)	Lt D1	Jastorf culture Wahlitz group Naumburg group Oder-Warthe group	Quedlinburg Hundisburg
200	BCE		Lt C2	Jastorf culture Naumburg group Oder-Warthe group	Walsleben Gommern Wulkau-Schönfeld Helfta
275	BCE		Lt C1	Jastorf culture Naumburg group	
300	BCE		Lt B2b		
400	BCE		Lt B1-2a	Jastorf culture Thuringian culture	
450	BCE		Lt A		
525	BCE	Early Iron Age (Hallstatt)	Ha D3	Jastorf culture House urn culture Billendorf culture Thuringian culture	Erdeborn Bösenburg Reppichau Niederröblingen Schlieben (BB)
575	BCE		Ha D2		
625	BCE		Ha D1	Elb-Havel group House urn culture Billendorf culture Thuringian culture	
750	BCE		Ha C		
1075	BCE	Late Bronze Age (Urnfield period)	Ha B1-3 = P IVb-V	Elb-Havel group Saalemündungs group Lusatian culture Unstrut group	Quenstedt Lüdelsen Kemberg Niederröblingen Löberitz Kuckenburg Oechlitz Schönebeck
1250	BCE		Ha A1-2 = P IIIb-IVa		Niederröblingen Piesteritz-Wittenberg Schafstädt Radis
1325	BCE		Bz D = P IIIa	Lusatian culture Unstrut group	
1475	BCE	Middle Bronze Age	Bz C = P II	Lüneburg group Fulda-Werra group	
1550	BCE		Bz B = P II		
2200	BCE	Early Bronze Age	Bz A = P I	Únětice culture	

* Bz = Bronzezeit (Bronze Age)

* Ha = Hallstatt

* Lt = La Tène

* RKZ = Römische Keiserzeit (Roman period)

* VWZ = Völkerwanderungszeit (Migration period)

* FMA = Frühmittelalter (Early Middle Ages)

* P = Period

Table 1. Chronology of archaeological phases, cultures and periods in central Germany and the sites in Saxony-Anhalt mentioned in the text. The time span is Bronze Age to Early Middle Ages (by R. Schwarz, Halle (Saale), modified by the author).

Supposed Neolithic finds of broomcorn millet in central Germany

Until 2008, there was a scientific consensus that broomcorn millet was already present in central Germany in the Early Neolithic, either as a crop or as a weed in cereal fields (Kreuz 1990; Wasylikowa *et al.* 1991; Körber-Grohne 1994; Stika and Heiss 2013; Behre 2008). Lüning (2000) emphasised that *P. miliaceum* was a component of the Linear Pottery culture package of crop species in the region. The same locations with ancient finds of broomcorn millet have been described in different archaeobotanical reports. Willerding (1980) and Rösch (1999) mentioned 10 Linear Pottery sites with *P. miliaceum* in eastern and central Europe, with the locations Hundisburg (Haldensleben) and Eisenberg as the westernmost ones. Körber-Grohne (1994) suggested that the oldest evidence for *P. miliaceum* comes from central Germany and named these two sites: Eisenberg, in Thuringia, and Hadmersleben (correct name: Haldensleben; see Hummel 1968; Hellmund 2012a), in Saxony-Anhalt (Fig. 1). Other authors also described these two sites as yielding ancient finds of broomcorn millet (e.g. Willerding 1980; Kreuz 1990; Körber-Grohne 1994; Schultze-Motel and Gall 1994; Zohary and Hopf 1994). Lüning (2000) listed three further Neolithic sites in central Germany as containing broomcorn millet, as well as five sites in eastern Europe. Wasylikowa *et al.* (1991) mentioned some more localities in eastern Europe. Most of these finds are not radiocarbon dated.

The site of **Eisenberg** (county of Saale-Holzland, federal state of Thuringia) was excavated in 1925 and was archaeobotanically analysed in the 1950s by Rothmaler and Natho (1957). Schultze-Motel re-examined the material. The assemblage consisted of *Triticum dicoccon*, *Triticum monococcum*, *Hordeum distichum/vulgare* (hulled), *P. miliaceum* and *Cannabis sativa*. Both Rothmaler and Natho (1957) and Schultze-Motel and Gall (1994) expressed doubts about the assignment of the hemp finds in the Linear Pottery culture contexts to *C. sativa*. A further re-examination indicates that the attribution of broomcorn millet to the Neolithic is in doubt as well.³

For Saxony-Anhalt, Körber-Grohne (1994) mentioned the site of Hadmersleben. This must be the village of **Hundisburg**, near the town of Haldensleben (county of Börde) (Hummel 1968; Willerding 1980; Hellmund 2012a). The archaeobotanical material was stored in the local Museum Haldensleben and marked with the old labels created by Rothmaler, who did the first analysis. This material was accidentally rediscovered during a student excursion to the Museum in the 1950s. Grains of broomcorn millet (Fig. 6.1) from Hundisburg were radiocarbon dated, and the date indicates a time span of 150-46 cal BCE (1 σ , Table 2, Fig. 2) (Hellmund 2012a). Evidently, Hundisburg is not the site with the oldest find of broomcorn millet in Saxony-Anhalt.

At the site of **Eitzum** (town of Schöppenstedt, county of Wolfenbüttel, federal state of Lower Saxony), two grains of *P. miliaceum* were found in structures of the Linear Pottery culture (Kreuz 1990). In more recent publications, Kreuz doubted the archaeological dating of the small grains of *P. miliaceum* to the Neolithic and wrote: ‘Solitary finds of *P. miliaceum* (broomcorn millet) are certainly later intrusions in the Linear Pottery Culture pits, transported there by earthworms, mice or other animals, as proven by various 14^c dates’ (Kreuz and Marinova 2017, 646).

In Westerhausen, at the site of **Jätchenberg**, near the town of Thale (county of Harz), a stone grave of the Globular Amphorae culture with a human skeleton, as well as two adjacent pits with burials of two and five cattle, respectively, were detected in 2003 (Meller 2012; Döhle and Pape 2021). The contents of some pots

3 B. Zach, Blaubeuren, Jena, and the author examined the material in question in January 2020 at the Institute for Archaeology in Jena, and the grains of *Panicum miliaceum* should be dated soon.

Laboratory *	Site and feature (f.)	Material	Radiocarbon date	Calibrated age 1 σ cal BCE	Calibrated age 2 σ cal BCE
KIA-33854	Bösenburg	1 barley grain	2452 \pm 33 BP	749-434	756-413
MAMS-47955	Gommern-Gerstenberg	1 barley grain	2244 \pm 23 BP	381-229	387-206
KIA-33853	Hundisburg	7 broomcorn millet grains	2080 \pm 30 BP	150-46	175 cal BCE-8 cal CE
MAMS-22384	Kemberg	fragment of wood	2701 \pm 21 BP	896-811	900-809
MAMS-22383	Kemberg	fragment of wood	2752 \pm 20 BP	917-837	969-828
Erl-6712	Löberitz	1 barley grain	2818 \pm 50 BP	1046-906	1118-835
KIA-37253	Niederröblingen, f. 3635	14 grains of broomcorn millet	2951 \pm 31 BP	1221-1115	1261-1051
KIA-34310	Niederröblingen, f. 3636	5 grains of broomcorn millet	2968 \pm 33 BP	1256-1125	1364-1053
KIA-35825	Niederröblingen, f. 3637	1 hulled barley grain	3012 \pm 26 BP	1369-1215	1384-1129
KIA-34311	Niederröblingen, f. 3646	1 hulled barley grain	2476 \pm 28 BP	754-543	770-426
MAMS-47954	Niederröblingen, f. 3709	2 fragments of hulled barley grains	2975 \pm 24 BP	1258-1128	1283-1113
KIA-36818	Niederröblingen, f. 4377	1 spelt grain	2548 \pm 21 BP	791-596	796-567
KIA-34309	Niederröblingen, f. 4718	1 emmer grain	2951 \pm 34 BP	1224-1111	1266-1021
KIA-36817	Niederröblingen, f. 4883	5 grains of broomcorn millet	2790 \pm 59 BP	1010-840	1109-815
MAMS-47959	Niederröblingen, f. 5359	1 spelt grain	2594 \pm 25 BP	800-780	809-770
KIA-34308	Niederröblingen, f. 5377	1 einkorn grain	2813 \pm 64 BP	1051-848	1190-819
MAMS-47961	Niederröblingen, f. 5397	2 fragments of hulled barley grains	2901 \pm 25 BP	1123-1019	1201-1008
KIA-37255	Niederröblingen, f. 9100	4 emmer grains	2973 \pm 31 BP	1259-1126	1369-1055
MAMS-47952	Piesteritz-Wittenberg	fused grains of broomcorn millet (8 mg)	2962 \pm 23 BP	1219-1126	1264-1058
MAMS-47958	Quenstedt, f. 10	1 barley grain	2735 \pm 25 BP	902-833	926-816
KIA-33851	Quenstedt, f. 116	1 fragment of faba bean seed	2743 \pm 31 BP	911-833	977-813
Poz-31957	Quenstedt, f. 12	5 broomcorn millet grains	2805 \pm 35 BP	1005-916	1051-836
Poz-31960	Quenstedt, f. 290	2 fragments of faba bean seeds	2740 \pm 35 BP	910-831	978-810
MAMS-47960	Quenstedt, f. 9	1 emmer grain	2749 \pm 25 BP	916-835	974-822
KIA-38121	Radis	fused grains of broomcorn millet (15 mg)	2994 \pm 36 BP	1285-1128	1386-1113
MAMS-47953	Reppichau	5 grains of broomcorn millet	2499 \pm 23 BP	761-553	773-543
KIA-33852	Schafstädt	1 hulled barley grain	2977 \pm 34 BP	1261-1127	1375-1055
MAMS-39145	Schönfeld-Wulkau	1 hulled barley grain	2251 \pm 25 BP	384-231	391-206
KIA-33855	Walsleben	5 grains of broomcorn millet	2182 \pm 33 BP	353-174	369-116

* KIA = Leibniz-Laboratory for Radiometric Dating and Stable Isotopic Research of Kiel University (Germany)

* Erl = AMS ^{14}C -Laboratory of the Friedrich-Alexander-University Erlangen-Nürnberg (Germany)

* Poz = AMS Laboratory of the Adam Mickiewicz-University in Poznań (Poland)

* MAMS = ^{14}C -Laboratory Curt-Engelhorn-Centre Archaeometry gGmbH in Mannheim (Germany)

Table 2. Radiocarbon dates of the sites and features mentioned in the text. Calibration using OxCal v4.4.2 and IntCal20 (Bronk Ramsey 2009; Reimer et al. 2020).





Figure 2. Radiocarbon dates from the following sites and materials: charred millet 'porridge' from Radis (RD) and Wittenberg-Piesteritz (WB); mass finds of millet from Reppichau (REK), Hundisburg (HUB) and Walsleben (WA); single finds of millet from Löberitz (LOB), Kemberg (KEF; date on wood), Bösenburg (BOE), Wulkau-Schönfeld (WSC), Gommern-Gerstenberg (GOV) and Schafstädt (SCH). Calibration using OxCal v4.4.2 and IntCal20 (Bronk Ramsey 2009; Reimer et al. 2020). Prepared by the author.

found in the grave were analysed. Sample WEH-04 contained uncarbonized grains of *P. miliaceum* and other plants. It was assumed that this is a recent contamination; indeed, the radiocarbon date (KIA-31868) of the subfossil grains of broomcorn millet proved that they post-date 1954 CE.

As part of a scientific project of Kiel University, megalithic tombs in Altmark in the northern part of Saxony-Anhalt were investigated, including the 'king's burial' in **Lüdelsen** (community of Jübar, county of Salzwedel), belonging to the Funnel Beaker culture (Diers 2018). H. Kroll analysed a large number of archaeobotanical samples from this site, detecting grains of *P. miliaceum* in two samples from Sector 7. Their radiocarbon dates fall in the Late Bronze Age, not the Neolithic (KIA-42500: 2783 ± 37 BP, 1σ 997-854 cal BCE, 2σ 1015-830 cal BCE; KIA-42499: 2684 ± 40 BP, 1σ 897-804 cal BCE, 2σ 910-794 cal BCE) (Diers 2018; Filipović et al. 2020).

Although located in the federal state of Brandenburg, the site of **Rathsdorf** (town of Wriezen, county of Märkisch-Oderland) is mentioned because of its published Neolithic age, as emphasised by Jahns et al. (2018) and Effenberger (2017). As described in Diers (2018) and Jahns et al. (2018), the radiometric dating of two grains of *P. miliaceum* from features no. 384 and no. 385 indicate the Neolithic age (KIA-42498: 3826 ± 49 BP, 1σ 2399-2152 cal BCE, 2σ 2459-2142 cal BCE). Therefore, Rathsdorf was thought to have yielded the oldest finds of broomcorn



millet in Brandenburg. However, Filipović *et al.* (2020) did not include this date in their Millet Dating Programme because it was obtained on a pooled sample of grains originating from multiple features and they considered it unreliable. There are other issues with this date too: the $\delta^{13}\text{C}$ value for the dated sample hints at the potential inclusion of C3 plant material (Filipović pers. comm.) and the dates produced by the Leibniz-Laboratorium in Kiel during this period (the dating was done in late 2010) turned out to be too old due to errors in the laboratory procedures.⁴

The town of **Quenstedt** (municipality of Arnstein, county of Mansfeld-Südharz) is situated in the northeastern forelands of the Harz Mountains; ca. 1 km southwest from it, in the Hengstbach valley, the so-called Schalkenburg hillock rises to a height of 20-25 m above the valley. On top of it is a multi-period archaeological site, excavated between 1967 and 1986. An area of 1.2 hectare was uncovered, including a circular palisade enclosure of the late Stroke-Ornamented Ware (*Stichbandkeramik*) culture; burials of different Neolithic cultures and of the Early Bronze Age Únětice culture; settlement pits of the Middle Neolithic Bernburg culture; and pits and postholes of a Late Bronze Age and Early Iron Age settlement, fortified by a rampart and a ditch cutting off the spur from the surrounding flat terrain (Behrens and Schröter 1980; Sosnowski 2006, 2014, 2015). Schultze-Motel analysed the archaeobotanical material and the present author conducted quantitative analysis (Schultze-Motel and Hellmund in press). Most finds of broomcorn millet at the site derive from the Late Bronze Age and Early Iron Age settlement, but in some cases they were also found in pits together with ceramics belonging to the Neolithic Bernburg culture. Grains of *P. miliaceum* were present in the assemblages from pits no. 9, 10, 12 and 116. Fragments of faba bean were found in features no. 116 and 290. Occasionally, *P. miliaceum* and *V. faba* were found in the same pit. One barley grain from pit no. 10, two fragments of *V. faba* from pit no. 290, one emmer grain from pit no. 9, a fragment of *V. faba* from pit no. 116 and several grains of *P. miliaceum* (Fig. 6.2) from pit no. 12 were radiocarbon dated. Four dates point to the 9th century cal BCE, and the radiocarbon date for pit no. 12 (millet) points to the 10th century cal BCE (1 σ , Table 2, Fig. 3).

Figure 3. Quenstedt-Schalkenburg. Radiocarbon dates for the pits suggested to belong to the Bernburg culture. Dated grains: pit no. 10 – barley; pit no. 290 – faba bean; pit no. 116 – faba bean; pit no. 9 – emmer; pit no. 12 – *P. miliaceum*. Calibration using OxCal v4.4.2 and IntCal20 (Bronk Ramsey 2009; Reimer *et al.* 2020). Prepared by the author.

⁴ Meadows *et al.* (2015) discuss the problematic measurements produced by the radiocarbon laboratory in Kiel and state that ‘the late-2010 and 2011 results are too old’ (Meadows *et al.* 2015, 1046).

Broomcorn millet in the Bronze Age in Saxony-Anhalt

Finds of broomcorn millet attributed to the Late Bronze Age are more common in Saxony-Anhalt than those attributed to the Neolithic. Due to the lack of sites from the Middle Bronze Age, little is known about the plant spectra from this period. Botanical finds from the Early Bronze Age (Únětice culture) are rare as well. No remains of *P. miliaceum* from the Únětice culture period are known (see section on the site of Oechlitz, below).

Mass finds and single finds of broomcorn millet in the Lusatian (Lausitz) culture

At the end of the 14th century BCE, communities of the Lusatian culture advanced from eastern Europe westwards. Their presence expanded from Poland into Bohemia, Moravia and Germany, specifically Brandenburg, Saxony and the eastern part of Saxony-Anhalt. In Saxony-Anhalt, they reached the Elbe River and settled in the area of the present-day town of Wittenberg (Schunke 2018). The first traces of the Lusatian culture date to Period III (1325-1150 BCE) of the chronology devised by Oscar Montelius. Settlement activity increased during Periods IV (1150-1000 BCE) and V (1000-750 BCE) and decreased during the Early Iron Age, when the Lusatian culture transformed (from an archaeological point of view) into the Billendorf culture (750-450 BCE) (Schmidt and Göricke 1985; Meller 2015) (Table 1). The Lusatian culture groups cremated their dead and buried them beneath earthen barrows surrounded by broad ditches (Schwarz 2003). The communal cemeteries often comprise dozens of burial mounds (Jockenhövel 2013).

The site of **Radis** (near the town of Kemberg, in the region of Wüste Mark Gemeln), lies in a forested area 12 km south of the Elbe. During rescue excavations in this forest in 1952-1953, a lump of charred millet ‘porridge’ was discovered in barrow no. 31, near a broken, ovoid pot lying next to some metal finds (Schmidt and Göricke 1985). The lump of fused grains of broomcorn millet measured 27.5 ml and weighed approximately 7.67 g. The radiocarbon date on a fragment of the porridge (Fig. 6.5) spans the 13th and 12th centuries cal BCE (1 σ , Table 2, Fig. 2).

Schmidt and Göricke (1985) also mentioned the site of **Piesteritz** (in the town of Lutherstadt Wittenberg, in the county of Wittenberg) as another site of the Lusatian culture. In 1994, salvage excavation took place at a burial location here (Wurda 1994), during which soil samples from pots as well as lumps of fused broomcorn millet grains were recovered. Sample WB-12i (Fig. 6.9) contained 4.525 g and sample WB-12h contained 7.1 g of the grains. Another sample had seeds of *V. faba* in it as well as millet. The grains of *P. miliaceum* (Fig. 6.3) were dated to the period between the mid-13th and 12th century cal BCE (1 σ , Table 2, Fig. 2).

From 1000 BCE onwards, settlements of the Lusatian and Billendorf cultures were often fortified. One of these is the fortification of **Kemberg** (Wittenberg county) (Meller 2015; Nebelsick and Swieder 2018), a waterlogged site from the end of the Late Bronze Age, considered the oldest fortified settlement of the Lusatian culture (Ha B and Ha C-D1). Timber from this site was dated to ca. 968 and 955 den BCE; two radiocarbon dates on wood fragments exist as well, and they point to the 9th century cal BCE (1 σ , Table 2, Fig. 2). Archaeobotanical investigations of Kemberg revealed residues of *P. miliaceum*, including an uncarbonized glume (Fig. 6.10) (Hellmund 2018).

The archive of the State Museum of Prehistory in Halle (Saale) houses plant remains from the site of Malitschkendorf, near **Schlieben**, located east of the Elbe (county of Elbe-Elster, federal state of Brandenburg). They were collected in 1828, by F.A. Wagner, and also in 1931. Other material was kept in different archives and analysed by several researchers. Buschan (1895, 73, 213) mentioned charred lumps

of broomcorn millet and seeds of faba bean. In the archive in Halle (Saale), there are four samples from Schlieben. The most diverse plant spectrum was documented in sample SCB-8286, consisting mostly of *Triticum dicoccon* (emmer grains and chaff), *V. faba*, *Pisum sativum*, *P. miliaceum*, naked wheat and several grains of barley. In addition, fragments of carbonised porridge of broomcorn millet were found. The site has been dated archaeologically to the Late Bronze to the Early Iron Age (Ha B-D1) (Nebelsick and Swieder 2018); no radiocarbon dates have yet been obtained.

Broomcorn millet at the sites of Late Bronze Age archaeological groups in Saxony-Anhalt

During the Late Bronze Age, different regional cultures developed in Saxony-Anhalt, named for a dominant river flowing through their respective areas of settlement: the Unstrut group (ca. 1325-750 BCE), the Saalemündungs group (1300-750 BCE) and the Elb-Havel group (Meller 2015) (Table 1). The Saalemündungs group was partially influenced by the neighbouring Lusatian culture and partially also by the Nordic Bronze Age culture. Cremation was quite common. From about 1000 BCE, the Saalemündungs group expanded farther south, into the areas of settlement of the Unstrut group. The Unstrut group continued the tradition of the Middle Bronze Age barrow culture. Inhumation in wooden coffins persisted for a long time, but around 1000 BCE, cremation became the dominant burial rite (Meller 2015). Further regional subdivision of the Unstrut group is represented by the Helmsdorf group in the northeastern and the Wandersleben group in the southwestern part of its distribution. These two subgroups were separated roughly by the low mountain ranges of the Schmücke and the Finne, which delimit the Thuringian Basin to the north-east (Schwarz pers. comm.). The richness of the bronze hoards found along the middle course of the Saale River (Schunke 2004) mirrors the wealth of the population of the Helmsdorf group, which was accumulated through salt production in the salt mines that existed in and near the city of Halle (Saale).

Mass finds of broomcorn millet in settlement pits

There are some mass finds with carbonised grains of broomcorn millet in the territory of Saxony-Anhalt. Prior to the construction of the A71 motorway from Sangerhausen to Erfurt in Thuringia, rescue excavations were carried out. The site of **Niederröblingen** (municipality of Allstedt, county of Mansfeld-Südharz) is located near the floodplain of the Helme in the southern Harz foreland. A tell-like settlement was detected here, the first known from central Germany, with a height in the centre of about 1.80 m. This site was occupied starting with the Linear Pottery culture. The accumulation of cultural layers accelerated during the Late Bronze Age (Meller 2011).

The archaeological situation at Niederröblingen is highly complex. On a visit to the site during the preparation of a new excavation area, the present author discovered two small pits filled with mass finds of carbonised grains of *P. miliaceum*. A third feature with a mass find of broomcorn millet was detected later. Initially, five grains of *P. miliaceum* from feature no. 3636 (Fig. 6.6) were sent for dating; the response from the Leibniz-Laboratory for Radiometric Dating in Kiel was that the amount of organic material had been too small. The second attempt at dating led to a result, showing that the material dated to the period encompassing the 2nd half of the 13th and the early 12th century cal BCE (1 σ , Table 2, Fig. 4). Within the mass finds, grains of *P. miliaceum* were mostly found in the 1 mm fraction of the sieved sample (Fig. 6.4). Grains from feature no. 3635, located near the pit (no. 3636), date to the same period (Fig. 4). Five grains of broomcorn millet from the third mass find (pit no. 4883) were also dated (Fig. 6.7), and these returned a date in the 10th to 9th centuries cal BCE (1 σ , Table 2, Fig. 4). A further 14 pit silos with charred




Figure 4. Niederröbblingen. Mass finds dominated by *Panicum miliaceum*: Feature nos. 3636, 3635 and 4883 from which broomcorn millet grains were dated; other features with single grains of *Panicum miliaceum* and grains of emmer, barley, einkorn and spelt, which were also dated. Calibration using OxCal v4.4.2 and IntCal20 (Bronk Ramsey 2009; Reimer et al. 2020). Prepared by the author.

cereals were excavated at Niederröbblingen. Finds of *P. miliaceum* are known from 12 dated granary pits (Table 2, Fig. 4). Most of these features date to the period before 780 cal BCE, whereas three of them date to the Early Iron Age (1 σ , Table 2, Fig. 4). Other granary pits mostly contained grains of other cereals, such as barley, spelt, emmer, einkorn and bread wheat. Remains of *Papaver somniferum* and *Camelina sativa* were also found but are relatively rare (Hellmund and Petzschmann 2011).

The exceptionally large find of charred cereals and pulses, about 400 L in volume (sample REK-06i), from the site of **Reppichau** (community of Osternienburger Land, county of Anhalt-Bitterfeld) in central Saxony-Anhalt consists of thousands of grains of broomcorn millet, peas, hulled barley and emmer. Grains of *P. miliaceum* (Fig. 6.8) have been dated to 760-550 cal BCE (1 σ , Table 2, Fig. 2).

Another mass find of charred grains of *P. miliaceum* was recorded in **Walsleben**, a village belonging to the Hanseatic town of Osterburg (Altmark), in the county of Stendal. The site contained primarily remains of a Roman period occupation, located on an elevation near the Uchte River; archaeological traces of the Pre-Roman Iron Age and the Neolithic were less common (Kolb and Müller 1996). The analysed sample consisted of several thousand grains of broomcorn millet and hundreds of faba bean seeds. Barley was the second-most dominant crop species and emmer was present as well. Grains of broomcorn millet have been dated (Fig. 6.12). Contrary to the assumption by the excavator that they come from the Roman period, their absolute age falls between the 4th and 2nd centuries cal BCE (1 σ , Table 2, Fig. 2).

Late Bronze Age sites with single finds of broomcorn millet

The construction of new tracks on the railway between Nürnberg (English: Nuremberg) and Halle (Saale) and Leipzig was preceded by excavations in the south of Halle (Saale). Preliminary results have been published (Hellmund 2017; Meller and Becker 2017). Charred grains of *P. miliaceum* (Fig. 6.11) were detected in settlement pit no. 20394 near **Oechlitz** (town of Mücheln, county of Saalekreis). The preliminary archaeological dating of this find was to the Early Bronze Age, specifically the Únětice culture, and so the material had been considered by the present author (Hellmund 2017) as deriving from this period. However, subsequent inspection of the updated database and discussion with an archaeologist led to the correction of the temporal attribution, since it was established that the pit belongs to the Late Bronze Age (Knoll and Fröhlich 2017) and not to the Únětice culture. For now, there are no radiocarbon dates available for this site.

At the site of **Schönebeck** (county of Schönebeck), several Neolithic and Bronze Age wells were excavated in advance of the construction of a bypass road. A bronze hoard was discovered outside a Late Bronze Age wooden box well (Bogen 2012). At the base of the shaft created for this well, three layers of the wooden box were preserved. According to the dates, the lowest wooden box was constructed in the Early Bronze Age, in 1995±10 den BCE. The middle and upper wooden boxes were dated to 1248±10 den BCE and 1223±10 den BCE, respectively. It can be assumed that the backfill in the well dates to the later phase of use, not to the earlier phase. The carbonised grains of *P. miliaceum*, found in the uppermost wooden box, were therefore assigned to the Late Bronze Age (Hellmund 2012b). Radiocarbon dates on the botanical macro-remains from the fill of the well are pending.

In 1992, a large deposit of cereals was recovered from a pit during the rescue excavation that preceded the building of a commercial area in **Schafstädt**, a village near the town of Bad Lauchstädt (county of Saalekreis). Barley was dominant in the deposit; grains of *P. miliaceum* and remains of a number of segetal and ruderal weeds, such as *S. viridis*, were also present (Hellmund 2012c). One grain of hulled barley returned a radiocarbon date of 13th to 12th century cal BCE (1σ, Table 2, Fig. 2).

In advance of a gas pipeline construction, a prehistoric settlement near **Löberitz**, near the town of Zörbig (county of Anhalt-Bitterfeld), was investigated. A well that had dried out (Leinthal 2004) yielded several grains of *P. miliaceum* (Hellmund 2004). Radiocarbon dating of a barley grain pinpointed the 11th to 10th century cal BCE (1σ, Table 2, Fig. 2) as the time when the well was in use (Hellmund 2012a).

The University of Jena recently excavated the multi-period site of **Kuckenburg** (municipality of Obhausen, county of Saalekreis) (Ettel *et al.* 2016). Plant remains from the Late Bronze Age (Ha A2-B) have been uncovered, including isolated finds of *P. miliaceum*.⁵

Pre-Roman Iron Age sites in Saxony-Anhalt with single finds of broomcorn millet

In the 1960s, a settlement pit with a large amount of charred cereal grains was excavated near **Bösenburg** (village near town of Gerbstedt, county of Mansfeld-Südharz). Hulled barley dominated the deposit; emmer, einkorn, spelt wheat were also represented, along with grains of broomcorn millet, *V. faba* and *Linum usitatissimum* (Schmidt *et al.* 1965; Schultze-Motel and Kruse 1965). The radiocarbon date on a barley grain falls in the period between the 8th and 5th centuries BCE (1σ, Table 2, Fig. 2) – that is, in the Early Iron Age (Hellmund 2012a).

5 The plant material is being processed by B. Zach, Blaubeuren.




Figure 5. Radiocarbon dates on grains of *Panicum miliaceum* from the following sites in Saxony-Anhalt: Hundisburg (HUB), Walsleben (WA), Reppichau (REK), Niederröblingen (NRL), Qenstedt-Schalkenburg (QUN), Wittenberg-Piesteritz (WB), Radis (RD). Calibration using OxCal v4.4.2 and IntCal20 (Bronk Ramsey 2009; Reimer et al. 2020). Prepared by the author.

At the site of **Erdeborn** (municipality of Seegebiet Mansfelder Land, county of Mansfeld-Südharz), a large pit contained numerous fragments of *briquettage* as well as wood charcoal pieces and charred fruits and seeds (Ipach 2016). The ceramics date the site to Hallstatt Ha C to D2, that is, the Early Iron Age. Grains of broomcorn millet were detected here (Hellmund 2016).

In proximity to the construction serving as flood protection east of the River Elbe, an oven containing human bones, charcoal, clay plaster and charred grains was uncovered at the site of **Schönfeld-Wulkau** (municipality of Kamern, county of Stendal) (Petzold in press). The archaeological finds date to La Tène phase C1. The archaeobotanical material was mostly composed of barley, but broomcorn millet was also present (Hellmund in press). A radiocarbon date on a barley grain confirms the timespan, of the 4th to 3rd century cal BCE (1 σ , Table 2, Fig. 2).

In 1982, a rescue excavation took place on a sand dune rising above the floodplain of the Elbe, where the multi-period site of **Gommern-Gerstenberg** is located, near the town of Gommern (county of Jerichower Land) (Weber and Albert 1984). In a pit with a large concentration of barley, some grains of emmer, einkorn and broomcorn millet were also found. Despite the assumption of the excavator that the material derives from the Early Iron Age, the radiocarbon date on a barley grain gave a span of the 4th to 3rd century cal BCE, that is, the Late Iron Age (1 σ , Table 2, Fig. 2). In further excavations at this site, the famous ‘princely grave of Gommern’, dating to the Roman period, was uncovered.

In the area of the so-called ‘**Königshof**’ (royal court) in Helfta, in the town of Lutherstadt Eisleben (county of Mansfeld-Südharz), in addition to remains from the Middle Ages, a structure of the La Tène culture was excavated and investigated by Schoknecht (1986). Here, within a large deposit of barley, grains of broomcorn millet and associated weeds were attested.




Figure 6. Selected finds of broomcorn millet grains from sites in Saxony-Anhalt, some radiocarbon dated: 1. Hundisburg HUB-001, 2. Quenstedt-Schalkenburg QUN-007, 3. Piesteritz-Wittenberg WB-12h, 4. Niederröbblingen NRL-M3635 (1 mm sieve fraction), 5. Radis RD-M01, 6. Niederröbblingen NRL-M3636, 7. Niederröbblingen NRL-M4883, 8. Reppichau REK-06i, 9. Piesteritz-Wittenberg WB-12i (fused carbonised grains), 10. Kemberg KEF-1110-2 (uncarbonised glume), 11. Oechlitz OEC-20394-6, 12. Walsleben WA-07. Scale 1 mm except for 4 and 10, where the scale is 1 cm. Radiocarbon dated: grains 1-3, 5-8, 12 and fragments of 3 and 5. (Photo: Monika Hellmund).

Sites from the Roman period in Saxony-Anhalt with single finds of broomcorn millet

There are several Roman sites in Saxony-Anhalt with small numbers of finds of *P. miliaceum*: Osterwieck, county of Harz; Elbeu⁶, county of Börde (Paysen and Nelle 2020); Wolmirstedt, county of Börde; Quedlinburg, county of Harz; and Klötze, county of Altmarkkreis Salzwedel (Leineweber and Willerding 2000). These finds were not radiocarbon dated, but dendrochronological dates were produced for wood fragments from the wells at Elbeu and Klötze.

Radiocarbon dates on the grains of *P. miliaceum* in Saxony-Anhalt: An overview

The radiocarbon dates obtained on the grains of broomcorn millet from Saxony-Anhalt are compiled in Figure 5. Apart from sample QUN-007 from Quenstedt-Schalkenburg, the samples were mass finds (concentrations). The dating shows that *P. miliaceum* was an important staple crop during the Late Bronze and the Pre-Roman Iron Age, and that it was also cultivated in the Roman period.

Discussion

The results show that, actually, there are no confirmed finds of broomcorn millet from either the Neolithic or the Early Bronze Age in Saxony-Anhalt. None of the recorded grains of broomcorn millet pre-date 1400 BCE. This is in line with the finding by Filipović *et al.* (2020) that *P. miliaceum* arrived in Europe in the 16th century BCE and in north-central Europe specifically in the 13th/12th century BCE. As noted above, the dating of the previously published finds of *P. miliaceum* from Oechlitz (Hellmund 2017) as Early Bronze Age must be revised because the finds have recently been archaeologically re-assigned to the Late Bronze Age.

The cultivation of broomcorn millet began in Saxony-Anhalt at about 1300/1200 BCE, early in the Late Bronze Age (Bz D-Ha A). The oldest records of broomcorn millet relate to burial sites of the Lusatian culture and a settlement site of the Unstrut group (Niederröbblingen). The features at Quenstedt-Schalkenburg containing *P. miliaceum* are younger than those at Niederröbblingen (Table 1). With regard to the seeds of faba bean from Quenstedt, Schultze-Motel already doubted the Neolithic age of their find context, pit no. 116:

‘Charred remains [of faba bean] are completely absent from the Neolithic in our area ... The occurrence of charred faba beans in pit 116 can only be accepted from a botanical point of view with great reservations’ (pers. comm. cited by Behrens and Schröter 1980, 99 translated by the author).

The radiocarbon-dated faba bean from pit no. 116 at Quenstedt-Schalkenburg dates to the 11th to 10th century cal BCE (Figs. 3, 5). Some of the pits at the site contained pottery of the Neolithic Bernburg culture together with finds of broomcorn millet. In five cases, the grains were directly radiocarbon dated, allowing for the exact age of the broomcorn millet to be established and the find to be assigned to the Late Bronze Age. Two of the pits with Bernburg culture pottery contained broomcorn millet and faba bean, another pit had broomcorn millet, and further three pits contained seeds of faba bean. These plant materials cannot automatically be attributed to the Neolithic based solely on the presence of the characteristic Bernburg culture pottery. On the contrary, it can be assumed that other finds of

6 However, dendro-dates on the wood from the well place this site in the Roman period, not in the Pre-Roman Iron Age.

broomcorn millet and faba bean from the pits at Quenstedt-Schalkenburg are also likely from the Late Bronze Age/Early Iron Age.

'Ancient' finds of *P. miliaceum* and *V. faba* in adjacent areas

Another 'early' find of *P. miliaceum* was reported from the ritual site located on the burial grounds of **Drehna**, part of the town of Luckau (county of Dahme-Spreewald, federal state of Brandenburg) (Rösler 1983). K. Kloss identified small pieces of carbonised crusts found in a pit as being composed mainly of the grains of 'Hirse', presumed by the author to refer to *P. miliaceum*, but this material should be reinvestigated. Other features at this site were archaeologically dated to the end of Period III (Tiedtke 2015), that is, the beginning of the Late Bronze Age according to the cultural periodisation applied in central Germany (Table 1).

In the federal state of Hesse, first indications of *P. miliaceum*, proposed to have been associated with the Únětice culture, were discovered at the sites of **Mardorf** (Kreuz 2000) and **Linsenberg**, in the Glauberg area. As far as is known, no ¹⁴C-dates were produced for these grains, but the radiocarbon dating of other plant material found confirmed an Early Bronze Age date for the sites (Kreuz 2016). However, as long as there is no corresponding radiocarbon dating of the grains of broomcorn millet, it remains only a hypothesis that broomcorn millet was already being cultivated in the Early Bronze Age. Given that a broomcorn millet grain from other sites in Hesse – for example **Bruchenbrücken**, county of Friedberg (Kreuz 1990) – was radiocarbon dated to the period between the end of 16th and early 14th century cal BCE (2σ; Motuzaite-Matuzeviciute *et al.* 2013), it seems likely that the cultivation of *P. miliaceum* began earlier in the adjacent regions to the south of Saxony-Anhalt.

Changes in the spectrum of crop plants due to climatic variations?

During the Bronze Age, two climatic deteriorations were recorded, one in the middle of the 2nd millennium, the so-called Lössen fluctuation, with an assumed shift towards cooler and wetter climatic conditions, and a second one around 800 BCE (Göschonen phase 1). Lake-level transgressions led to the final abandonment of wetland settlements on the lake shores in Switzerland and the surrounding regions (Della Casa 2013). The varied climate during the Subboreal period, with alternating wet and dry phases, may have initiated or accelerated the introduction of new crop species. In this context, cultivation of the warm-season plant *P. miliaceum* would have been beneficial during the suggested dry episodes in the Bronze Age. In the so-called 'Mitteldeutsches Trockengebiet' in parts of Thuringia and Saxony-Anhalt, where the present-day rainfall is less than 500 mm per year, cultivation of *P. miliaceum* may have been particularly favoured. It has been suggested that a climate suitable for growing wine grapes is good for broomcorn millet too (Steinbrück 1908). But cultivation in the past was not restricted to such areas, and the large deposits of *P. miliaceum* in settlement pits described above, as well as the charred millet porridge in graves, prove its cultivation in a range of local climates.

Broomcorn millet and faba bean: A meal for the dead?

Some of the described finds of broomcorn millet come from burial sites (*e.g.* Piesteritz-Wittenberg and Radis in Saxony-Anhalt, Drehna in Brandenburg). The question arises as to whether the plants detected in burial contexts were in some

way special, *i.e.* exotic, and/or whether they were part of everyday diet. Frachetti *et al.* (2010), for example, interpreted the millet grain found in a 3rd millennium BCE burial cist in Kazakhstan as an element of the ritual. This is especially relevant where large deposits or fused grains of millet ('prepared food') have been found. There is a mention in the literature of another location with such a find – **Ostro-Jiedlitz** (county of Kamenz, federal state of Saxony), dated to the Early Iron Age (Ha C2-D1) (Nebelsick and Swieder 2018), where Pax and Hofmann (1915) found carbonized lumps of probable *P. miliaceum* grains. Effenberger (2017) and Jahns *et al.* (2018) did not record any remains of charred millet 'porridge' from graves in Brandenburg, but charred grains of broomcorn millet and seeds of faba bean have been found, for example, at the site of **Großbahren** (in the town of Sonnewalde, county of Elbe-Elster, federal state of Brandenburg) (Effenberger 2017). In his old compilation, Netolitzky (1914) mentions other sites where charred broomcorn millet crusts were found. But it is uncertain whether these finds have been saved and stored in archives.

Broomcorn millet in ritual contexts

The grains of broomcorn millet found in burial contexts can be interpreted as grave goods – food for the dead. According to *Handwörterbuch des deutschen Aberglaubens* [*Concise Dictionary of German Superstitions*], millets represented food for the soul that appeared in entertainment at funeral feasts. Therefore, broomcorn millet could be understood as a food for the dead. Moreover, millets were perceived as a symbol of fertility because of the many small grains, and they were also considered as symbolising wealth (Bächtold-Stäubli *et al.* 1987a).

Faba bean in ritual contexts

V. faba is often found in graves of the Lusatian culture and in settlement pits of the Late Bronze Age. Schultze-Motel (1972) and Jäger (1965, 1987) reported that the earliest finds of this species come from central Germany. More finds have since been added, including those from the sites of Quenstedt-Schalkenburg, Wittenberg-Piesteritz, Walsleben, Coswig (Anhalt) and Buroer Feld in Saxony-Anhalt. In the former village of Tornow (town of Lübbenau/Spreewald, county of Oberspreewald-Lausitz, federal state of Brandenburg), *V. faba* has been documented at a site dated to the Bronze Age Period V (Ha B) (Jäger 1965). Finds of *V. faba* were also registered at the ritual site on the burial ground of Drehna, in the state of Brandenburg (Rösler 1983).

Faba bean was also important in the ancient cult of the dead, according to the folk knowledge described in the *Concise Dictionary of German Superstitions*. In the Roman period, it was used in the cult of the soul (Bächtold-Stäubli 1987b). Faba bean may have been as important in prehistoric times in ritual contexts as its presence in the graves suggest. Pliny described how faba bean was used in the celebrations of the dead because of the belief that the beans preserved the soul of the deceased (Lenz 1859).

Broomcorn millet and faba bean in settlement structures

The settlement pits at Niederröblingen yielded three mass finds of cereals dominated by broomcorn millet and dating from the Late Bronze Age; no legumes were detected there. At Quenstedt-Schalkenburg, small amounts of *P. miliaceum* were discovered in more than a dozen features. Here, broomcorn millet was one of the most important crop species, after emmer and barley. It can be assumed that, in the pits, *P. miliaceum* became mixed with the older, Neolithic Bernburg culture pottery

as a result of soil movement or bioturbation. Indeed, during the Late Bronze Age, and again at the end of the Hallstatt period and beginning of the La Tène culture, the site underwent large-scale digging and levelling as part of the construction of the fortification (Sosnowski 2014, 2015). Unlike at Niederröblingen, at the Late Bronze Age site of Quenstedt-Schalkenburg, legumes such as *V. faba* played an important role alongside cereals.

Based on the frequent finds of *P. miliaceum* and *V. faba* in the settlements at Quenstedt-Schalkenburg and Walsleben, and the substantial evidence for broomcorn millet at Niederröblingen, it can be concluded that both species played an important role in human diet. Apparently, no distinction was made between the food of the living and the food for the dead.

Conclusion

The direct ¹⁴C-dating of grains of *P. miliaceum* and seeds of *V. faba* confirms that there are no Neolithic remains of these species in Saxony-Anhalt. Thus far, there is no evidence for the presence of broomcorn millet in the Early and Middle Bronze Age either. Actually, there are now doubts about the age of all 'ancient', non-radiocarbon-dated (single) finds of broomcorn millet from Saxony-Anhalt, especially those stated to be older than the middle of the 2nd millennium BCE based on their archaeological context. In Hesse, a somewhat earlier cultivation of *P. miliaceum* can be assumed, as indicated for example by the radiocarbon date on millet grains from Bruchenbrücken. Climatic changes, namely deterioration caused by cool and humid conditions and/or the intervening drier periods, may have promoted the growing of *V. faba* and *P. miliaceum*. The introduction of new crops could have been a response to the changing climatic conditions or a result of increased cultural connections, exchange and trade with regions in eastern or southeastern Europe, where the evidence for these two species is older, as shown by the radiocarbon dates on broomcorn millet grains. A definite and obvious advantage of the cultivation of new crop species would have been diversification of the subsistence base as a way of reducing the risk of crop failure.

Acknowledgements

The author is indebted to Ralf Schwarz, LDA Halle (Saale) for the chronology table and for details on the archaeological cultures. Thanks are also due to the two reviewers (anonymous and Julian Wiethold, Metz) for helpful suggestions and to Mathilde Schäfer-Buss, Adelaide, and the native-English copy editor Suzanne Needs-Howarth, Toronto, for the linguistic improvements to the manuscript.

References

- Bächtold-Stäubli, H., ed., 1987a (1942). Hirse. In: E. Hoffmann-Krayer and H. Bächtold-Stäubli, eds., *Handwörterbuch des deutschen Aberglaubens*, Band 4. Berlin: Walter de Gruyter, 117-122.
- Bächtold-Stäubli, H., ed., 1987b (1942). Bohne. In: E. Hoffmann-Krayer and H. Bächtold-Stäubli, eds., *Handwörterbuch des deutschen Aberglaubens*, Band 1. Berlin: Walter de Gruyter, 1470-1473.
- Behre, K.-E., 2008. *Landschaftsgeschichte Norddeutschlands: Umwelt und Siedlung von der Steinzeit bis zur Gegenwart*. Neumünster: Wachholtz.
- Behrens, H. and Schröter, E., 1980. *Siedlungen und Gräber der Trichterbecherkultur und Schnurkeramik bei Halle (Saale). Ergebnisse von Ausgrabungen*. Veröffentlichungen des Landesmuseums für Vorgeschichte in Halle (Saale) 34. Berlin: Deutscher Verlag der Wissenschaften.

- Bertsch, K. and Bertsch, F., 1949. *Geschichte unserer Kulturpflanzen*. Stuttgart: Wissenschaftliche Verlagsgesellschaft.
- Bogen, C., 2012. Brunnen und Opfer – ein bronzezeitlicher Kastenbrunnen mit zwei Bronzedepots. In: H. Meller, ed., *Von Egeln bis Schönebeck. Archäologie und Straßenbau in der Magdeburger Börde*. Archäologie in Sachsen-Anhalt Sonderband 20. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 108-122.
- Bray, F., 1981. Millet cultivation in China: a historical survey. *Journal d'agriculture traditionnelle et de botanique appliquée*, 28 (3-4), 291-307.
- Bronk Ramsey, C., 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon*, 51 (1), 337-360. <https://doi.org/10.1017/S0033822200033865>
- Buschan, G., 1895. *Vorgeschichtliche Botanik der Cultur- und Nutzpflanzen der Alten Welt auf Grund prähistorischer Funde*. Breslau: Kern.
- Della Casa, P., 2013. Switzerland and the Alps. In: H. Fokkens and A. Harding, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 706-722.
- Diers, S., 2018. *Mensch-Umweltbeziehungen zwischen 4000 und 2200 cal BC: Vegetationsgeschichtliche Untersuchungen an Mooren und trichterbecherzeitlichen Fundplätzen der Altmark*. Frühe Monumentalität und soziale Differenzierung 15. Bonn: Dr. Rudolf Habelt GmbH.
- Döhle, H.-J. and Pape, J., 2021. Ein bemerkenswertes Grabensemble der Kugelamphorenkultur – Steinkiste und Rindergruben von Westerhausen. In: H. Meller, ed., *Kataloge zur Dauerausstellung im Landesmuseum für Vorgeschichte Halle*. Band 2. Teil 1: *Früh- und Mittelneolithikum*. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale) 435-445.
- Effenberger, H., 2017. *Pflanzennutzung und Ausbreitungswege von Innovationen im Pflanzenbau der Nordischen Bronzezeit und angrenzender Regionen*. Studien zur nordeuropäischen Bronzezeit 4. Neumünster: Wachholtz.
- Ettel, P., Karl, H.-V., Paust, E., Pollock, M., Schmidt, C., Spazier, T., Wellhöfer, R.F., 2016. Vorbericht zu den Grabungen 2005 bis 2011 der Friedrich-Schiller-Universität Jena auf der Kuckenburg bei Esperstedt, Saalekreis. *Archäologie in Sachsen-Anhalt*, 8 (16), 19-38.
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, J., Jakobitsch, T., Kapcia, M., Kloß, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri A.M., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H.-P., Stobbe, A., Tjaša Tolar, T., Wasylukowa, K., Wiethold, J., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Frachetti, M.D., Spengler, R.N., Fritz, G.J., Mar'yashev, A.N., 2010. Earliest direct evidence for broomcorn millet and wheat in the central Eurasian steppe region. *Antiquity*, 84 (326), 993-1010. <https://doi.org/10.1017/S0003598X0006703X>
- Geisler, G., 1991. *Farbatlas landwirtschaftlicher Kulturpflanzen*. Stuttgart: Ulmer.
- Hegi, G., 1979. *Illustrierte Flora von Mitteleuropa*, Band I, Teil 3: *Pteridophyta, Spermatophyta, Monocotyledoneae 1, Gramineae*. Berlin: Parey Verlag, 1-80.
- Hellmund, M., 2004. Pflanzenreste – in den Brunnen gefallen. In: MITGAS Mitteldeutsche Gasversorgung GmbH, H. Meller, J. Oexle, eds., *Von Peißen nach Wiederitzsch: Archäologie an einer Erdgas-Trasse*. Gröbers: MITGAS Mitteldeutsche Gasversorgung GmbH, 45-47.
- Hellmund, M., 2012a. Frühneolithischer Anbau von Rispenhirse in Mitteldeutschland? Der eisenzeitliche Rispenhirsefund von Hundisburg, Lkr. Börde. *Archäologie in Sachsen-Anhalt*, 6, 297-303.

- Hellmund, M., 2012b. Pflanzenfunde aus dem spätbronzezeitlichen Kastenbrunnen von Schönebeck. In: H. Meller, ed., *Von Egelu bis Schönebeck: Archäologie und Straßenbau in der Magdeburger Börde*. Archäologie Sachsen-Anhalt Sonderband 20. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 124-128.
- Hellmund, M., 2012c. Ein spätbronzezeitlicher Getreidevorrat von Schafstädt, Saalekreis. *Archäologie in Sachsen-Anhalt*, 6, 304-315.
- Hellmund, M., 2016. Eisenzeitliche Pflanzenreste von der Salzsieder-Fundstelle Erdeborn, Lkr. Mansfeld-Südharz. In: S. Ipach, ed., *Die Salzsieder-Fundplätze der älteren Eisenzeit von Erdeborn in Sachsen-Anhalt und Steinhaleben in Thüringen*. Jenaer Schriften zur Vorgeschichte 6. Langenweißbach: Beier & Beran, 72-90.
- Hellmund, M., 2017. Pflanzenreste aus archäologischen Befunden der Schnurkeramik-, Glockenbecher- und Aunjetitzer Kultur im südlichen Sachsen-Anhalt. In: H. Meller and M. Becker, eds., *Neue Gleise auf alten Wegen II: Jügendorf bis Gröbers I*. Archäologie in Sachsen-Anhalt Sonderband 26. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 202-218.
- Hellmund, M., 2018. Botanische Befunde vom spätbronzezeitlichen Burgwall bei Kemberg, Lkr. Wittenberg. In: H. Meller and L.D. Nebelsick, eds., *Über den Wallrand geschaut. Der Kemberger Burgwall und sein bronze- und eisenzeitliches Umfeld*. Kleine Hefte zur Archäologie 15. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 119-128.
- Hellmund, M., in press. Das Pflanzenspektrum von Wulkau-Schönfeld – Pflanzenfunde aus der vorrömischen Eisenzeit. *Jahresschrift mitteldeutsche Vorgeschichte*.
- Hellmund, M. and Petzschmann, U., 2011. Verkohlte Getreidereste in großen Mengen. In: S. Friederich and H. Meller, eds., *Kultur in Schichten. Archäologie am Autobahndreieck Südharz (A 71)*. Archäologie in Sachsen-Anhalt Sonderband 14. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 191-200.
- Hummel, H.-S., 1968. Pflanzenfunde aus der Urgeschichte des mitteldeutschen Raumes. *Jahresschrift für mitteldeutsche Vorgeschichte*, 52, 39-66.
- Hunt, H.V., Vander Linden, M., Liu, X., Motuzaite-Matuzeviciute, G., Colledge, S., Jones, M.K., 2008. Millets across Eurasia: chronology and context of early records of the genera *Panicum* and *Setaria* from archaeological sites in the Old World. *Vegetation History and Archaeobotany*, 17, S5-S18. <https://dx.doi.org/10.1007%2Fs00334-008-0187-1>
- Ipach, S., 2016. *Die Salzsieder-Fundplätze der älteren Eisenzeit von Erdeborn in Sachsen-Anhalt und Steinhaleben in Thüringen*. Jenaer Schriften zur Vorgeschichte 6. Langenweißbach: Beier & Beran.
- Jäger, K.-D., 1965. Verkohlte Samen aus einem bronzezeitlichen Grabgefäß von Tornow, Kreis Calau: ein Beitrag zur Anbaugeschichte der Ackerbohne (*Vicia faba* L.) in Mitteleuropa. *Ausgrabungen und Funde*, 10, 131-138.
- Jäger, K.-D., 1987. Zur Rolle der Ackerbohne (*Vicia faba* L.) in Landwirtschaft und Brauchtum der Urnenfelderzeit in Mitteleuropa. In: E. Plesl and J. Hrala, eds., *Die Urnenfelderkulturen Mitteleuropas*. Symposium Liblice 21-25.10.1985. Praha: Archäologisches Institut der Tschechoslowakischen Akademie der Wissenschaften, 27-35.
- Jahns, S., Alsleben, A., Bittmann, F., Brande, A., Christiansen, J., Dannath, Y., Effenberger, H., Giesecke, T., Jäger, K.-D., Kirleis, W., Kloß, S., Kloss, K., Kroll, H., Lange, E., Medović, A., Neef, R., Stika, H.-P., Sudhaus, D., Wiethold, J., Wolters, S., 2018. Zur Geschichte der nacheiszeitlichen Umwelt und der Kulturpflanzen im Land Brandenburg. In: S. Flohr and P. Morgenstern, eds., *Beiträge zur Archäozoologie und Prähistorischen Anthropologie*. Band XI. Langenweißbach: Beier & Beran, 9-35.

- Jockenhövel, A., 2013. Germany in the Bronze Age. In: H. Fokkens and A. Harding, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 723-745.
- Knoll, F. and Fröhlich, M., 2017. Wände, Böden und andere Strukturen – verziegelte Lehmfragmente als Quelle zur Rekonstruktion von Gebäuden. In: H. Meller and M. Becker, eds., *Neue Gleise auf alten Wegen II: Jüdinghof bis Gröbers I*. Archäologie in Sachsen-Anhalt Sonderband 26. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 176-190.
- Kolb, M. and Müller, A. 1996. Ein Suchschnitt durch die Altmark – Grabungen auf der Erdgastrasse. *Archäologische Berichte aus Sachsen-Anhalt 1994*, 134-139.
- Körber-Grohne, U., 1994. *Nutzpflanzen in Deutschland: Kulturgeschichte und Biologie*. Stuttgart: Theiss.
- Kreuz, A., 1990. *Die ersten Bauern Mitteleuropas – eine archäobotanische Untersuchung zu Umwelt und Landwirtschaft der Ältesten Bandkeramik*. *Analecta Praehistorica Leidensia* 23. Leiden: Leiden University.
- Kreuz, A., 2000. Neolithische und bronzezeitliche Pflanzenfunde von Mardorf 3, Stadt Amöneburg, Kreis Marburg-Biedenkopf (Hessen). In: D. Vorlauf, ed. *Eine mehrperiodige Siedlungsstelle in Mardorf, Stadt Amöneburg (Kr. Marburg-Biedenkopf): Neolithikum bis Eisenzeit, unter besonderer Berücksichtigung der Urnenfelderkultur*. Kleine Schriften des Vorgeschichtlichen Seminars der Phillips-Universität Marburg 50. Marburg: Vorgeschichtliches Seminar der Phillips-Universität Marburg, 137-146.
- Kreuz, A., 2016. Eisenzeitlicher Ackerbau und Landnutzungspotential im Umfeld des Glaubergs im überregionalen Vergleich. In: L. Hansen and C.F.E. Pare, eds., *Untersuchungen im Umland des Glaubergs. Zur Genese und Entwicklung eines frühlatènezeitlichen Fürstensitzes in der östlichen Wetterau*. Wiesbaden: Landesamt für Denkmalpflege Hessen, 211-241.
- Kreuz, A. and Marinova, E., 2017. Archaeobotanical evidence of crop growing and diet within the areas of the Karanovo and the Linear Pottery cultures: a quantitative and qualitative approach. *Vegetation History and Archaeobotany*, 26, 639-657. <https://doi.org/10.1007/s00334-017-0643-x>
- Kroll, H., 2001. Literature on archaeological remains of cultivated plants (1999/2000). *Vegetation History and Archaeobotany*, 10, 33-60. <https://doi.org/10.1007/PL00013368>
- Leineweber, R. and Willerding, U., 2000. Ein kaiserzeitlicher Kastenbrunnen aus Klötze, Altmarkkreis Salzwedel: archäologische und paläo-ethnobotanische Befunde. *Jahresschrift für mitteldeutsche Vorgeschichte*, 83, 141-189.
- Leinthal, B., 2004. Wasserversorgung in der Bronzezeit. In: MITGAS Mitteldeutsche Gasversorgung GmbH, H. Meller, J. Oexle, eds., *Von Peißen nach Wiederitzsch: Archäologie an einer Erdgas-Trasse*. Gröbers: MITGAS Mitteldeutsche Gasversorgung GmbH, 42-44.
- Leipe, C., Long, T., Sergusheva, E.A., Wagner, M., Tarasov, P.E., 2019. Discontinuous spread of millet agriculture in eastern Asia and prehistoric population dynamics. *Science Advances*, 5 (9), eaax6225. <https://doi.org/10.1126/sciadv.aax6225>
- Lenz, H.O., 1859. *Botanik der alten Griechen und Römer, deutsch in Auszügen aus deren Schriften und deren Anmerkungen*. Gotha: Verlag E.F. Thienemann.
- Li, T., Ning, C., Zhushchikhovskaya, I.S., Hudson, M., Robbeets, M., 2020. Millet agriculture dispersed from northeast China to the Russian Far East: integrating archaeology, genetics, and linguistics. *Archaeological Research in Asia*, 22, 100177. <https://doi.org/10.1016/j.ara.2020.100177>
- Liu, X., Hunt, H.V., Jones, M.K., 2009. River valleys and foothills: changing archaeological perceptions of north China's earliest farms. *Antiquity*, 83 (319), 82-95. <https://doi.org/10.1017/S0003598X00098100>

- Lüning, J., 2000. *Steinzeitliche Bauern in Deutschland: die Landwirtschaft im Neolithikum*. Universitätsforschungen zur prähistorischen Archäologie 58. Bonn: Dr. Rudolf Habelt GmbH.
- Meadows, J., Hüls, M., Schneider, R., 2015. Accuracy and reproducibility of ^{14}C measurements at the Leibniz-Labor, Kiel: a first response to Lull *et al.*, 'When ^{14}C dates fall beyond the limits of uncertainty: an assessment of anomalies in western Mediterranean Bronze Age ^{14}C series'. *Radiocarbon*, 57 (5), 1041-1047. https://doi.org/10.2458/azu_rc.57.18569
- Mehnert, M., Zierold, R., Stegmann, H., Paulekon, D., 1964. *Richtzahlen und Tabellen für die Landwirtschaft*. Deutsche Akademie der Landwirtschaftswissenschaften zu Berlin. Institut für Agrarökonomik. Berlin: VEG Deutscher Landwirtschaftsverlag.
- Meller, H., ed., 2011. *Kultur in Schichten – Ausgrabungen am Autobahndreieck Südharz (A71)*. Archäologie in Sachsen-Anhalt Sonderband 14. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale).
- Meller, H., ed., 2012. *LebensWandel – Früh- und Mittelneolithikum*. Begleithefte zur Dauerausstellung im Landesmuseum für Vorgeschichte 3. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale).
- Meller, H., ed., 2015. *Glutgeboren. Mittelbronzezeit bis Eisenzeit*. Begleithefte zur Dauerausstellung im Landesmuseum für Vorgeschichte 5. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale).
- Meller, H. and Becker, M., eds., 2017. *Neue Gleise auf alten Wegen II: Jügendorf bis Gröbers I*. Archäologie in Sachsen-Anhalt Sonderband 26. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale).
- Meller, H. and Nebelsick, L.D., eds., 2018. *Über den Wallrand geschaut. Der Kemberger Burgwall und sein bronze- und eisenzeitliches Umfeld*. Kleine Hefte zur Archäologie 15. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale).
- Miller, N.F., Spengler, R.N., Frachetti, M., 2016. Millet cultivation across Eurasia: origins, spread, and the influence of seasonal climate. *The Holocene*, 26 (10), 1566-1575. <https://doi.org/10.1177%2F0959683616641742>
- Motuzaitė-Matuzevičiūtė, G., Staff, R.A., Hunt, H.V., Liu, X., Jones, M.K., 2013. The early chronology of broomcorn millet (*Panicum miliaceum*). *Antiquity*, 87 (338), 1073-1085. <https://doi.org/10.1017/S0003598X00049875>
- Nebelsick, L.D. and Swieder, A., 2018. Der Burgwall von Kemberg, Lkr. Wittenberg. Eine dendrochronologisch datierte spätbronze-/früheisenzeitliche Befestigung am Südrand der Elbaue bei Wittenberg. Vorbericht über die Ausgrabungskampagne im Jahr 2014. In: H. Meller and L.D. Nebelsick, eds., *Über den Wallrand geschaut. Der Kemberger Burgwall und sein bronze- und eisenzeitliches Umfeld*. Kleine Hefte zur Archäologie in Sachsen-Anhalt 15. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 51-118.
- Netolitzky, F., 1914. Die Hirse aus antiken Funden. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften zu Wien, mathematisch-naturwissenschaftliche Klasse*, 123 (I), 725-759.
- Paysen, A. and Nelle, O., 2020. Botanische Analysen am eisenzeitlichen Brunnen von Elbeu. In: S. Friederich and H. Meller, eds., *Elbeu – Die Burg Albrechts des Bären. Aktuelle Ausgrabungen: Neolithikum bis Mittelalter*. Archäologie in Sachsen-Anhalt Sonderband 28. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 83-90.

- Pax, F. and Hofmann, K., 1915. Prähistorische Pflanzen aus Schlesien und der Ober-Lausitz. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*, 52, 346-353.
- Petzold, D., in press. Die Frau im Ofen: eine Sonderbestattung der späten vorrömischen Eisenzeit aus dem nördlichen Elb-Havel-Winkel bei Wulkau, Ldkr. Stendal. *Jahresschrift mitteldeutsche Vorgeschichte*.
- Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., van der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S., 2020. The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0-55 cal kBP). *Radiocarbon*, 62 (4), 725-757. <https://doi.org/10.1017/RDC.2020.41>
- Rösch, M., 1999. Hirse. In: H. Beck, D. Geuenich, H. Steuer, eds., *Johannes Hoops Reallexikon der Germanischen Altertumskunde*. Band 14. Berlin and New York: De Gruyter, 612-614.
- Rösler, H., 1983. Eine mittelbronzezeitliche Opferstelle von Drehna, Kr. Luckau. *Ausgrabungen und Funde*, 28 (2), 57-64.
- Rothmaler, W. and Natho, I., 1957. Bandkeramische Kulturpflanzenreste aus Thüringen und Sachsen. In: W. Rothmaler and W. Padberg, eds., *Beiträge zur Frühgeschichte der Landwirtschaft*. Band 3. Berlin: Akademie-Verlag, 73-98.
- Schmidt, B. and Göricke, G., 1985. Gräber, Siedlungen und Befestigungen der Westgruppe der Lausitzer Kultur. *Ausgrabungen und Funde*, 30 (4), 182-185.
- Schmidt, B., Schultze-Motel, J., Kruse, J., 1965. Früheisenzeitliche Vorratsgrube auf der Bösenburg, Kr. Eisleben. *Ausgrabungen und Funde*, 10, 29-31.
- Schoknecht, T., 1986. Getreidefunde von Helfta, Kr. Eisleben. *Zeitschrift für Archäologie*, 20, 253-267.
- Schultze-Motel, J., 1972. Die archäologischen Reste der Ackerbohne, *Vicia faba*, und die Genese der Art. *Die Kulturpflanze*, 19, 321-358.
- Schultze-Motel, J., 1994. Literature on archaeological remains of cultivated plants (1991/1992). *Vegetation History and Archaeobotany*, 3, 33-61. <https://doi.org/10.1007/s10722-020-00996-3>
- Schultze-Motel, J. and Gall, W., 1994. *Archäologische Kulturpflanzenreste aus Thüringen*. Weimarer Monographien zur Ur- und Frühgeschichte 32. Stuttgart: Konrad Theiss.
- Schultze-Motel, J. and Hellmund, M., in press. Mittelneolithische, spätbronze- und eisenzeitliche Pflanzenfunde von der Schalkenburg bei Quenstedt, Lkr. Mansfeld-Südharz. *Jahresschrift mitteldeutsche Vorgeschichte*.
- Schultze-Motel, J. and Kruse, J., 1965. Spelz (*Triticum spelta* L.), andere Kulturpflanzen und Unkräuter in der frühen Eisenzeit Mitteleuropas. *Die Kulturpflanze*, 13, 586-619.
- Schunke, T., 2018. An der westlichen Peripherie der Lausitzer Kultur: Die mittelbronze- bis früheisenzeitlichen Gräberfelder bei Coswig, Lkr. Wittenberg. In: H. Meller and L.D. Nebelsick, eds., *Über den Wallrand geschaut. Der Kemberger Burgwall und sein bronze- und eisenzeitliches Umfeld*. Kleine Hefte zur Archäologie 15. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 271-282.
- Schunke, T., 2004. Der Hortfund von Hohenweiden-Rockendorf, Saalkreis, und der Bronzekreis Mittelsaale. Ein Beitrag zur jungbronzezeitlichen Kulturgruppengliederung in Mitteleuropa. *Jahresschrift mitteldeutsche Vorgeschichte*, 88, 219-337.

- Schwarz, R., 2003. *Pilotstudien. Zwölf Jahre Luftbildarchäologie in Sachsen-Anhalt*. Halle (Saale): Landesamt für Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale).
- Sosnowski, S., 2014. Der polykulturelle Fundplatz Quenstedt, Flur 'Schalkenburg', Lkr. Mansfeld-Südharz. Neue Ergebnisse zu Besiedlungsstruktur und -abfolge. *Archäologie in Sachsen-Anhalt*, 7 (14), 55-69.
- Sosnowski, S., 2006. *Der polykulturelle Fundplatz Quenstedt 4, Flur 'Schalkenburg', Ldkr. Mansfelder Land. Untersuchungen zu Besiedlungsstruktur und -abfolge anhand der Siedlungsbefunde*. Master thesis, Martin-Luther-Universität Halle-Wittenberg.
- Sosnowski, S., 2014. Der polykulturelle Fundplatz Quenstedt, Flur 'Schalkenburg', Lkr. Mansfeld-Südharz. Neue Ergebnisse zu Besiedlungsstruktur und -abfolge. *Archäologie in Sachsen-Anhalt*, 7 (14), 55-69.
- Sosnowski, S., 2015. *Die Besiedlung der Schalkenburg bei Quenstedt, Lkr. Mansfeld-Südharz während der frühen Bronze- bis älteren Vorrömischen Eisenzeit: Studie zum Besiedlungsablauf einer mehrphasigen Höhengründung im östlichen Mittelgebirgsraum*. Doctoral dissertation, Martin-Luther-Universität Halle-Wittenberg.
- Steinbrück, K., 1908. *Der praktische Landwirt: Handbuch der gesamten Landwirtschaft*. Band 3. Hannover: Max Jänecke-Verlag.
- Stevens, C.J. and Fuller, D.Q., 2017. The spread of agriculture in eastern Asia: archaeological bases for hypothetical farmer/language dispersals. *Language Dynamics and Change*, 7 (2), 152-186. <https://doi.org/10.1163/22105832-00702001>
- Stika, H.-P. and Heiss, A.G., 2013. Bronzezeitliche Landwirtschaft in Europa. Der Versuch einer Gesamtdarstellung des Forschungsstandes. In: K.-H. Willroth, ed., *Siedlungen der älteren Bronzezeit. Beiträge zur Siedlungsarchäologie u. Paläoökologie des zweiten vorchristlichen Jahrtausends in Südsandinavien, Norddeutschland und den Niederlanden*. Workshop vom 7.-9. April 2011 in Sankelmark. Studien nordeuropäischen Bronzezeit 1. Neumünster: Wachholtz, 189-222.
- Tafuri, M.A., Craig, O.E., Canci, A., 2009. Stable isotope evidence for the consumption of millet and other plants in Bronze Age Italy. *American Journal of Physical Anthropology*, 139, 146-153. <https://doi.org/10.1002/ajpa.20955>
- Tiedtke, V., 2015. Grave, pig and pine: animals and plants at the Lusatian urn field of Müllrose, East Brandenburg. In: J. Kneisel, M. Dal Corso, W. Kirleis, H. Scholz, N. Taylor, V. Tiedtke, eds., *The Third Food Revolution? Setting the Bronze Age Table: Common Trends in Economic and Subsistence Strategies in Bronze Age Europe*. Universitätsforschungen zur prähistorischen Archäologie 283. Bonn: Dr. Rudolf Habelt GmbH, 75-85.
- Van Zeist, W., Wasylikowa, K., Behre, K.-E., eds., 1991. *Progress in Old World palaeoethnobotany: a retrospective view on the occasion of 20 years of International Work Group for Palaeoethnobotany*. Rotterdam: Balkema.
- Ventresca Miller, A.R. and Makarewicz, C.A., 2019. Intensification in pastoralist cereal use coincides with the expansion of trans-regional networks in the Eurasian Steppe. *Scientific Reports*, 9, 8363. <https://doi.org/10.1038/s41598-018-35758-w>
- Wasylikowa, K., Cărciumaru, M., Hajnalová, E., Hartyányi, B.P., Pashkevich, G.A., Yanusevich, Z.V., 1991. East-central Europe. In: W. Van Zeist, K. Wasylikowa, K.-E. Behre, eds., *Progress in Old World palaeoethnobotany: a retrospective view on the occasion of 20 years of International Work Group for Palaeoethnobotany*. Rotterdam: Balkema, 207-239.
- Weber, T. and Albert, W., 1984. Eine früheisenzeitliche Siedlung bei Gommern, Kr. Burg (Grabung 1982). *Ausgrabungen und Funde*, 29, 193-195.
- Willerding, U., 1980. Zum Ackerbau der Bandkeramiker. In: T. Krüger, ed., *Beiträge zur Archäologie Nordwestdeutschlands und Mitteleuropas: Klaus Raddatz zum 65. Geburtstag am 19. November 1979*. Materialien zur Ur- u. Frühgeschichte Niedersachsens 16. Hildesheim: Marie Leidorf GmbH, 421-456.

- Wurda, A., 1996. Archäologische Arbeiten in der Lutherstadt Wittenberg. *Archäologische Berichte aus Sachsen-Anhalt* 1994, 337-341.
- Xu, Y., Liu, M., Li, C., Sun, F., Lu, P., Meng, F., Zhao X., He, M., Wang, F., Zhu, X., Hao, X., Zhou, H., 2019. Domestication and spread of broomcorn millet (*Panicum miliaceum* L.) revealed by phylogeography of cultivated and weedy populations. *Agronomy*, 9 (12), 835. <https://doi.org/10.3390/agronomy9120835>
- Zhao, Z., 2011. New archaeobotanic data for the study of the origins of agriculture in China. *Current Anthropology*, 52 (S4), S295-S305. <https://doi.org/10.1086/659308>
- Zohary, D. and Hopf, M., 1994. *Domestication of Plants in the Old World*. Oxford: Clarendon.
- Zohary, D., Hopf, M., Weiss, E., 2012. *Domestication of Plants in the Old World: the Origin and Spread of Domesticated Plants in Southwest Asia, Europe, and the Mediterranean Basin*. (4th edition). Oxford: Oxford University Press.

Millets in Bronze Age agriculture and food consumption in northeastern France

*Françoise Toulemonde, Julian Wiethold,
Emmanuelle Bonnaire, Geneviève Daoulas,
Marie Derreumaux, Frédérique Durand, Bénédicte Pradat,
Oriane Rousselet, Caroline Schaal, Véronique Zech-Matterne*

Abstract

Populations of the Early Bronze Age and the Middle Bronze Age left a small footprint in northeastern France, settlement evidence being very scarce and funerary contexts barely more numerous. Starting from the 14th c. BCE onwards, important changes occurred in settlement and land use. An ongoing development of settlements, expanding into previously unexploited environments, is observed. This major demographic increase was sustained by a new and dynamic agropastoral economy, in which broomcorn millet seems to have played a central role. To better understand this role, as well as the chronology of changes in farming practices, archaeobotanical research on Bronze Age data from northeastern France has been coupled with radiocarbon dating of the earliest records of broomcorn millet. Results suggest that the introduction of broomcorn millet in France did not take place before the 14th c. BCE, a date in line with the recently published chronological model of the spread of millet in Europe (Filipović *et al.* 2020).

The introduction of millet was not the only change occurring in Bronze Age farming systems. Archaeobotanical evidence indicates that the diversification of crops, integrating new species of cereals, pulses and oil plants, together with a less cereal-focused farming production, had generated a resilient agriculture, securing crop production and thus promoting soaring demographics. Consequently, the gathering of wild carbohydrate resources, such as acorns, declined.

If not the only new crop, broomcorn millet was one of the most widespread new crops during the Late Bronze Age in northeastern France (Bouby *et al.* 2017). Its diffusion seems to have been quick and thorough. Millet was part of the daily diet on every small farm; it was also on the menu for festive meals during large

collective gatherings and accompanied the dead as a funerary offering. This small-seeded cereal was indeed the emblematic plant of food identity during the Late Bronze Age.

Introduction

As the star cereal of the European Bronze Age, broomcorn millet (*Panicum miliaceum*) has been the subject of intense interdisciplinary research for the past 15 years, the issues and methods of which have strongly evolved over time (Liu *et al.* 2018). Questions have shifted from the origin of the plant and the hypothesis of multiple domestications, eastward and westward of central Asia (Jones 2004; Hunt *et al.* 2008; Zohary *et al.* 2012), to the tracking and chronology of the paths of the spread of millet from China to Europe. The development of Chinese archaeobotanical investigations (Ren *et al.* 2016), complemented by stable isotopes studies (Lightfoot *et al.* 2013), advances in archaeogenetic research (Hunt *et al.* 2014; Hunt *et al.* 2018) and radiocarbon dating programmes on European prehistoric millets (*e.g.* Motuzaite Matuzeviciute *et al.* 2013) have indeed succeeded in demonstrating both the seniority of Chinese records and the Chinese origin of broomcorn and foxtail millets (Miller *et al.* 2016; Liu *et al.* 2018). The earliest European records have been brought forward by several millennia, to the time of the first material evidence of contact between East and West, during the Bronze Age (Sherratt 2006). Research has subsequently focused on the modelling of the westward dispersal of millet from China and the identification of the hubs and tempo of diffusion in Europe (Miller *et al.* 2016; Liu *et al.* 2018; Filipović *et al.* 2020). In the framework of recent millet research, many direct dating programmes have already been conducted all over Eurasia (*e.g.* Filipović *et al.* 2018; Herrscher *et al.* 2018), while others are still in progress.

Now that the question of the chronology of the arrival of millet in Europe is being resolved, other issues are emerging about the changes possibly brought about by this new cereal crop ('the millet effect'). Innovations may have concerned many different aspects, such as farming systems, human and animal diet, population increase, expansion, and the organisation of societies. These topics were discussed during the Millet International Workshop held in Kiel, November 2019, and this article formed a contribution to it. It provides unpublished information on French data related to the research questions just mentioned.

Starting with the 'when' of the arrival of millet, we present chronological data from northeastern France, incorporating both chrono-cultural and direct dating. An extensive archaeobotanical synthesis was initially carried out based on all available data from Bronze Age settlements in this area. A series of radiocarbon dates was then obtained from the earliest records, to check their antiquity. This set of dates documents a new milestone in the history of millet diffusion in Europe, extending westward the recent works on more eastern regions. Northeastern France had strong connections with central Europe during the Middle Bronze Age and Late Bronze Age (MBA, LBA, from approximately 1650/1600 to 800/750 BCE) (Mordant 2013) and was probably the gateway into France for millet (Bouby *et al.* 2017).

Continuing with the 'how' and 'why', we present factual data on changes in farming systems and subsistence economy that took place in LBA societies after the introduction of millet. We discuss the role of millet in crop husbandry organisation as well as in the daily diet and festive celebrations and underline the fact that this small-seeded cereal of Chinese origin, today a minor crop in Europe, was a major element of the food identity, not only in northeastern France, but in a large part of Europe during the LBA.

Before presenting the archaeobotanical data, we present the archaeological background. Settlement and land use underwent major changes during the Bronze Age, concomitant with those observed in farming systems. It is therefore essential to explore these changes to better understand the overall scheme.

Bronze Age in northeastern France: The archaeological background

Several archaeological overviews concerning the Bronze Age in France have been published in the past decade (Mordant 2013; Carozza *et al.* 2017a; Lachenal *et al.* 2017; Guilaine and Garcia 2018). However, despite the exponential increase in discoveries over the past 25 years, which ensures we have a reliable archaeological dataset, there has been no re-evaluation of knowledge about land use during the Early Bronze Age (EBA) and MBA (2300/2200-1350 BCE) in northeastern France. Overall, finds relating to domestic occupation remain very sparse for these early phases, in striking contrast to the spectacular number of discoveries for the subsequent period, starting at the transition between the MBA and the LBA (14th c. BCE; Carozza *et al.* 2017b, Figs. 5-8). This observation is particularly true for the westernmost part of the study area, in the Île-de-France and Champagne regions (Fig. 1). Even in extensively excavated areas, such as the Upper Seine Valley and the Marne Valley (Peake *et al.* 2017a, 2017b; Peake 2020, 13-15) or the plain of Troyes (Riquier *et al.* 2017a, 2017b), funerary contexts and settlement sites from the EBA and MBA are not visible. Traces of occupation mainly consist of some isolated pits or waste accumulations in oxbow lakes, with an extremely small footprint and no evidence of buildings or organised settlement. Starting from the middle of the 14th c. BCE and the transition to LBA1 (1350-1100 BCE), settlements multiply exponentially, are more visible in the landscape, and more permanent. At first, they come as modest and isolated open-air farms, comprising pits with rich archaeological deposits, sometimes associated with small post-framed buildings. Then, during the five centuries of the LBA (1350-750 BCE), land occupation intensifies, extending out of the valleys and onto the plateaus of Île-de-France and the chalk plain of Champagne, while settlements diversify. During LBA3 (900-750 BCE), although most settlements are still small farmsteads, others develop into hamlets, such as the sites of Buchères, in Champagne (Riquier *et al.* 2012); Rosières-aux-Salines, in Lorraine (Koenig and Klag 2017); and Marolles-sur-Seine, in Île-de-France (Peake *et al.* 2017a). At the very end of the LBA, fortified sites, with a more ostentatious rather than truly defensive character, appear in the Upper Seine Valley, punctuating it spatially at regular intervals, and illustrating the hierarchy of Bronze Age society. These high-status sites interact with smaller settlements and sometimes host seasonal collective gatherings and feasts, such as at Villiers-sur-Seine (Auxiette *et al.* 2015; Peake 2020). Palisaded storage areas start to develop at the beginning of the Early Iron Age (EIA), with dozens of granaries in the Champagne chalklands (Desbrosse and Riquier 2012) and storage pits and granaries on the loess plateaus of Île-de-France (Peake *et al.* 2017a) and in the Moselle Valley (Deffressigne *et al.* 2002).

In Picardy, a similar evolution took place. A low human footprint for the EBA and MBA is followed by a significant rise in the number of settlements starting at LBA2 (1100-900 BCE). During LBA3, some houses are grouped into hamlets, such as in Choisy-au-Bac, and the development of large storage areas is observed (Bucheux *et al.* 2017). In other northeastern regions, such as Alsace (Michler *et al.* 2017), Lorraine (Koenig and Klag 2017), and northern Burgundy (Ducreux 2017), EBA settlements are more visible, but not very numerous. Different types of building plans have been documented, such as large, three-aisled post-framed buildings in the Moselle Valley or two-aisled buildings in Alsace. The MBA seems to be a period of extremely low population size and land use everywhere in northeastern France. This changes at the




Figure 1. (a) Location of 130 Bronze Age settlements with archaeobotanical data in northeastern France (b) Results of the archaeobotanical synthesis per Bronze Age phase. EBA=Early Bronze Age; MBA=Middle Bronze Age; LBA=Late Bronze Age. Mass finds=concentrations of ≥ 100 remains per litre.

beginning of the LBA, when a sudden and impressive increase in human occupation is observed throughout the area. This would continue for several centuries.

The reasons for such a low human footprint during the EBA and MBA are unknown. Hypotheses include erosion of sites, or an economy based on a semi-nomadic way of life, with settlements leaving few traces (Riquier *et al.* 2017b). The sudden increase in settlements in such areas as the Upper Seine Valley, which had been almost deserted during the MBA, has been linked to potential migrations from eastern regions, as the source of cultural influences in funerary material shifted from west to east at the end of the MBA (Peake *et al.* 2017b). The whole phenomenon – sluggishness and then a marked surge in population – seems to have been a Europe-wide occurrence (Stika and Heiss 2013; Kneisel *et al.* 2015a), and thereby may also have been linked to large-scale events affecting agriculture, such as climate change, although showing complex and regionally different developments (Magny *et al.* 2007; Kneisel *et al.* 2015a; Capuzzo *et al.* 2018). In reality, it is likely that a complex combination of many different parameters was the cause.

For whatever reason, a sparse population with a low environmental impact seems to have been the rule in most places in northeastern France until the end of the MBA. From the 14th c. BCE onwards, a new rural society emerges and develops.

It is characterised by a densification of population networks, supported by a very dynamic agropastoral economy. Millets seem to have played a central role in this new economy. An overview of data tracing the time of the arrival and the rhythm of spread of millet cultivation may help us identify its role.

Arrival and adoption of millet cultivation in northeastern France: Synthesis of the archaeobotanical evidence

The overview of archaeobotanical data with a focus on millets that we aim to present in this paper was carried out for the millet workshop held in Kiel, November 2019. As several recent works had demonstrated that millet did not arrive in Europe during the Neolithic (see Introduction and Conclusions to this volume), we chose to consider only finds dated to the Bronze Age. This work constitutes an important update of previous work. Following the very first research on millets in France, published at a time when the finds of this plant were scarce and studies concerned mainly southern France (Marinval 1992, 1995), a national survey gathering all available French data for the Bronze Age and Early Iron Age was conducted in 2011 and published in 2017 (Bouby *et al.* 2017). Information was collected from 272 French sites, 47 of which fall within the geographical and chronological target of our present work. Since this broad synthesis, the activity of rescue archaeology, coupled with the continued expansion of archaeobotany in northeastern France, has more than doubled the available data for Bronze Age in the research area. The quantity of additional data, and the need for a higher chronological resolution than the one used in the 2011 survey, made it necessary to carry out a new study, this time at a regional level.

Material

The geographical framework of this overview, northeastern France (Fig. 1a), includes the Grand Est region (bringing together the former regions of Champagne-Ardenne, Lorraine, and Alsace); the eastern part of the Île-de-France (département of Seine-et-Marne) and Hauts-de-France (département of Aisne in Picardy) regions; and the northern part of the Bourgogne-Franche-Comté region (départements of Yonne, Côte-d'Or, and Haute-Saône). The Bronze Age in this area lasts from approximately 2300/2200 to 850/750 BCE. It is culturally divided into the Early Bronze Age (2300/2200-1600 BCE), the Middle Bronze Age (1600-1350 BCE) and the Late Bronze Age (1350-800/750 BCE), which is itself split into LBA1 (1350-1100 BCE), LBA2 (1100-900 BCE) and LBA3 (900-800/750 BCE). The data presented here derive from 130 distinct occupation phases from a total of 96 settlement sites, some sites having been occupied in several phases (Fig. 1b). The dataset was assembled by 12 archaeobotanists, belonging to different institutions, and represents all available information for the study as of November 2019 (see Table 3 below).

Most settlements are small, isolated and open-air family farms, but there are some exceptions for LBA3, when the range of site types diversifies (see above). For this period, it also includes a hamlet-like site (Buchères; Riquier *et al.* 2012, 2015), fortified aristocratic settlements (such as Villiers-sur-Seine; Auxiette *et al.* 2015; Peake 2020) and palisaded enclosures with numerous granaries (as at Bazancourt; Desbrosse and Riquier 2012).

Sampled features mainly consist of waste pits, storage pits and postholes; they also include several ditches, ovens, hearths and wells. Systematic bulk sampling was carried out, with a minimum volume of 10 litres per sample whenever possible. Samples were processed by water-flotation or wet sieving, using a minimum mesh size of 0.25 to 0.5 mm.

Site	Location	Archaeological context	Number of charred grains dated	Dating by relative chronology (BCE)	Laboratory sample number	Direct dating ¹⁴ C (years BP)	Direct dating (cal BCE) with 2σ probability (95.4%)	AMS δ ¹³ C (‰)	Carbon mass [mg]
Réau <i>Parc d'activités de l'A5</i>	Seine-et-Marne, Île-de-France	Pit	6	1350-1100	Poz-129554	3045 ± 30	1401-1220	−8.5	1.56
Balloy <i>La Haute Borne</i>	Seine-et-Marne, Île-de-France	Pit	8	1350-1200	Poz-129551	3040 ± 30	1401-1216	−10.9	0.64
Gingsheim <i>Steinbrünnen, Aschenbuckel</i>	Bas-Rhin, Alsace	Pit	5	1350-1200	Poz-129549	2980 ± 30	1375-1059	−14.3	1.48
Hérange <i>Gross Eichholz</i>	Moselle, Lorraine	Pit	6	1350-1200	Poz-129550	2960 ± 30	1265-1054	−11.3	2.06
Sézannes <i>Maison de Santé</i>	Marne, Champagne	Pit	10	1350-1200	Poz-129552	2945 ± 35	1262-1020	−15.1	0.19
Courceroy <i>Les Dizaines</i>	Aube, Champagne	Well	1	1350-1100	Poz-129553	2865 ± 30	1125-926	−16	0.68
Réau <i>Centre Pénitencier</i>	Seine-et-Marne, Île-de-France	Pit	3	1350-1100	Poz-133340	2835 ± 30	1109-908	−6.7	1.391
Ville-Saint-Jacques <i>Le Bois d'Echalas</i>	Seine-et-Marne, Île-de-France	Storage pit	10	1350-1100	Poz-129182	2475 ± 30	770-423	−8.4	1.73
Pont-sur-Seine <i>La Gravière</i>	Aube, Champagne	Building postholes	10	1450-1300	Poz-129548	2335 ± 30	514-236	−14	2.25

Table 1. Results of AMS ¹⁴C-dating of early finds of broomcorn millet in northeastern France.

Results

Results of direct AMS-radiocarbon dating

At the time of the presentation in Kiel, nearly all reported dates for the archaeobotanical material came from relative dating, based on the association of fossilised plant remains with cultural artefacts, mainly pottery, in archaeological layers. A few dates derived from radiocarbon dating on charcoal, cereal caryopses or animal bones, but there was no direct dating of millet remains. Being fully aware that this has resulted in misleading ages for intrusive grains of millet, as demonstrated by the results of the radiocarbon dating programmes recently carried out in Europe (Motuzaite Matuzeviciute *et al.* 2013; Filipović *et al.* 2020), we have since undertaken direct dating on broomcorn millet, starting with the earliest (and most questionable) finds. The goal was to remove doubts about the age of these remains and to specify the date of millet arrival in the study area. These first direct dates launch a database that will help, once it is more extensive, to trace the routes of the introduction and spread of millets in France. The database will also allow comparisons with neighbouring countries where large radiocarbon dating programmes have already been conducted.

During the process of selecting the grains to be dated, some of the earliest finds presented in Kiel were discarded for various reasons: dubious taxonomic identification for Pagny-lès-Goings-Aéroport de Moselle, Lorraine (2200-1500 BCE); re-attribution of the feature of the find to a later date for Warcq-Bois de Charnois, Ardennes (from 1450-1300 to 1200-1100 BCE); and material not accessible in the case of Ay-sur-Moselle-Les Vêlers Jacques (1350-1200 BCE). A total of 10 samples of the remaining earliest discoveries of broomcorn millet, 9 of which concern charred grains and 1 of which concerns waterlogged glumes, were submitted to Poznań Radiocarbon Laboratory (Poland) for AMS-dating. The finds come from different



Figure 2. Some of the plant remains sent for radiocarbon dating: (a) Waterlogged lemma/palea of *Echinochloa crus-galli*/*Panicum miliaceum* from Erstein-Parc d'Activités du Pays d'Erstein, Bas-Rhin; Charred grains of *Panicum miliaceum* from (b) Hérange-Gross Eichholz, Moselle, (c) Sézannes-Maison de Santé, Marne, (d) Réau-Parc d'activités de l'A5, Seine-et-Marne, (e) Pont-sur-Seine-La Gravière, Aube, (f) Balloy-La Haute Borne, Seine-et-Marne, (g) Gingsheim-Steinbrünnen, Aschenbuckel, Bas-Rhin, (h) Ville-Saint-Jacques-Le Bois d'Echallas, Seine-et-Marne.




Figure 3. Calibrated AMS ^{14}C dates of broomcorn millet from northeastern France (calibration with OxCal v.4.4.2 and IntCal 2020 software; Bronk Ramsey 2009, 2017; Reimer et al. 2020).

areas, covering the entire northeastern quarter of France, from Alsace to Île-de-France (Fig. 1a). The results of the AMS-dating are presented in Table 1 and Figure 3.


The supposed earliest find consists of six waterlogged lemma/palea (Fig. 2a) recently recovered from a well at the site of Erstein-Parc d'Activités du Pays d'Erstein, Bas-Rhin, in Alsace (Croutsch *et al.* 2016b). They appeared to be an interesting discovery, as the different types of dating performed on the well layer (dendrochronology on the oak fixtures, typo-chronology of the pottery and radiocarbon dating on a human bone) all converge to the 16th c. BCE (Croutsch *et al.* 2016a). But, identification of the glumes as coming from *P. miliaceum* was doubtful and could not be ascertained because of the poor preservation of the glumes; *Echinochloa crus-galli* was a more likely candidate (Kroll pers. comm.). Moreover, the remains could not be radiocarbon dated because of their low carbon content. For both reasons, this find was not taken into consideration.

The charred grains from Pont-sur-Seine-La Gravière and Ville-Saint-Jacques-Le Bois d'Echallas (Fig. 2e, h) turned out to date to a much later period than expected (Table 1, Fig. 3), namely 514-236 cal BCE (Late Iron Age) and 770-423 cal BCE (Iron Age), respectively. These two sites are multi-phase settlements, with occupations whose relative chronology encompasses the radiocarbon dates (Issenmann *et al.* 2009; Dupéré *et al.* 2013; Toulemonde 2013).

The other six radiocarbon dates fall within the expected timeframe according to the cultural chronology (Table 1). The earliest finds are recorded from Île-de-France (Balloy, in the Upper Seine Valley, and Réau, on the Sénart plateau). Given the current state of our direct dating results, we can conclude, at least provisionally, that broomcorn millet did not arrive in northeastern France before the 14th c. BCE.

Results of the archaeobotanical synthesis

Archaeobotanical studies carried out on the 96 Bronze Age sites identified a total of 131,151 charred plant remains from 1052 samples. Of these remains, 58,092 came from waste assemblages and 73,059 from four cereal mass finds. A total of 23,035 waterlogged plant remains have also been recovered, from wells at five different sites. As expected, the distribution of data is uneven, with the Late Bronze Age accounting for more than 95% of the charred material (Fig. 1b).



Early Bronze Age and Middle Bronze Age (2300/2200-1350 BCE)

Nine sites have been dated to the EBA. They are mainly located in the eastern part of the study area, in Alsace and Lorraine, where more settlements are known for this period than farther west (Fig. 4a). The archaeological material dates four of these sites to 2200-1650 BCE, and five others to 2000-1650 BCE. Altogether, they yielded 2,483 charred and 2,017 waterlogged plant remains. No millet is reported from these assemblages. Seven more sites are less accurately dated to the EBA/MBA (Fig. 4a). Plant remains are scarce (347 charred remains) and millet is absent.

For the MBA, the five sites analysed are again mainly located in Alsace and Lorraine (Fig. 4a). Charred remains are even scarcer (179) and comprise no millet, if we exclude the doubtful millet from Erstein-Parc d'activités du Pays d'Erstein (see above).

At the end of the MBA/beginning of LBA (15th-14th c. BCE), human occupation gradually increased. Four settlements were studied for this period and two of them supposedly yielded broomcorn millet. However, as explained earlier, the

Figure 4. Records of broomcorn millet (*Panicum miliaceum*) and foxtail millet (*Setaria italica*) from northeastern France: (a) from the Early Bronze Age (EBA) to the Middle Bronze Age-Late Bronze Age transition (MBA/LBA), (b) during LBA1, (c) during LBA2, (d) during LBA3.

radiocarbon dating of some of the grains of the large, charred deposits (2453 caryopses of broomcorn millet) found in Pont-sur-Seine-La Gravière (Champagne) resulted in a much later date for the millet (Fig. 3). It corresponds to the beginning of the EIA, the time of the second settlement phase at this site. At Couternon-Larrey (Burgundy), two grains of broomcorn millet were found in a silo pit dated to the MBA-beginning of LBA (Wiethold 2007).

Late Bronze Age (1350-800/750 BCE)

From the very beginning of the LBA, human settlements multiplied, and almost all of them, from east to west, cultivated broomcorn millet (Fig. 4b, Table 2, Table 3). The few sites that did not yield millet caryopses are also those that revealed very scarce remains of domestic plants in general. It is as if this cereal started to be cultivated everywhere simultaneously or at least within a very short timeframe, and was intricately linked to the expansion of human settlement, which had become visible and more permanent. There is no evidence of a slow beginning followed by a progressive increase. This could be due to the absence of data for the MBA. Maybe the first attempts at millet cultivation are invisible, much like the MBA settlements.

Cultivation of the new cereal maintains momentum throughout the LBA (Fig. 4c, d, Table 2). Both the ubiquity of broomcorn millet (Table 2) and the proportions of its caryopses in relation to those of other cereals (barley, wheat and Cerealia) during the three phases of the LBA (Fig. 5) confirm its importance in agricultural production since the very beginning of this period. Previous research has already shown that the ubiquity of millet could regularly reach 60% to 70% of the samples per site in some parts of the Upper Seine Valley or Champagne (Toulemonde 2013; Goude *et al.* 2016). Charred mass finds are not recovered before LBA3 (10th-8th c. BCE); when they appear, they include millets as well as other cereals (Figs. 4d, 5).

But what about foxtail millet (*Setaria italica*)? The first discovery comes from Sézannes, Champagne. It is comprised of a single charred caryopsis, dated to the 14th-13th c. BCE (Fig. 4b). Charred grains of broomcorn millet from a different layer within the same feature have been radiocarbon dated to 1262-1020 cal BCE (Table 1). Other records are available for the following period (12th-10th c. BCE) from all over northeastern France (Fig. 4c, d). *Setaria italica* may have been introduced as a weed of *Panicum miliaceum*, since only a few grains, associated with larger amounts of broomcorn millet, are initially present. Then, at the end of LBA3 (10th-8th c. BCE), the site of Buchères-Parc Logistique de l'Aube (Champagne) provided a cache of 200 grains of *Setaria*, mixed with 60 grains of *Panicum* and other cereals, suggesting that *Setaria* could have been a crop in its own right at that time. More *Setaria* is recorded for the Early Iron Age, confirming that its cultivation was well established, even if it seems to have never been as widespread as *Panicum miliaceum*.

Period (BCE)	Number of settlements with more than 20 charred grains of domestic plants (=X)	Number of settlements with broomcorn millet (% of X)
14th-13th c.	6	5 (83%)
14th-12th c.	6	5 (83%)
13th-11th c.	6	5 (83%)
12th-11th c.	6	6 (100%)
12th-10th c.	23	23 (100%)
11th-10th c.	6	5 (83%)
10th-9th c.	13	12 (92%)
10th-8th c.	18	15 (83%)

Table 2. Ubiquity of broomcorn millet in Late Bronze Age settlements over time.




Figure 5. Proportions (in %) of the main categories of cereals for each phase of the Late Bronze Age (LBA), calculated from the total number (nb) of charred grains in all sites of one phase. Waste assemblages (bar-charts) and mass finds (pie-charts) are analysed separately.

Conclusion on the dating of millets from northeastern France

Archaeobotanical results (Table 3) coupled with radiocarbon dates of the earliest discoveries of broomcorn millet show that *Panicum miliaceum* was not introduced in northeastern France before the 14th c. BCE. It is likely that the cultivation of broomcorn millet started between the 14th and the 13th c. BCE and spread very rapidly, becoming part of every small farm production throughout the area.

Further radiocarbon evidence concerning the cultivation of broomcorn millet in northeastern France comes from miliacin, a chemical marker strictly associated with *Panicum miliaceum*. It was isolated from a Bronze Age silo and a Late Iron Age silo at the site of Obernai, in Alsace (Courrel *et al.* 2017). The miliacin was archived in the soil filling the silos. It presumably resulted from past cultivation of millet during the Bronze Age, because the miliacin recovered from the silo of the Iron Age proved to be older than the structure itself. Four radiocarbon dates have been obtained on the miliacin extracted at different depths of the fills of the two silos. The resulting ^{14}C ages (1560-1259, 1438-996 and 2044-826 cal BCE for the Bronze Age silo; 1461-894 cal BCE for the Iron Age silo) encompass those provided by our dates on millet grains.

These observations are also consistent with the recently published chronological model of millet arrival in Europe (Filipović *et al.* 2020) based on radiocarbon dates. This model shows that by the mid-14th c. BCE, cultivation of broomcorn millet spread into central Europe, the area nearest to northeastern France, while earlier occurrences (from the mid-15th c. BCE) are recorded in the Po Basin, in northern Italy. The only millet dated to the 16th c. BCE comes from Ukraine. The authors conclude that it is very unlikely that broomcorn millet reached central Europe before the mid-14th c. BCE.

As for foxtail millet, its arrival in northeastern France is not recorded before the 13th c. BCE. It was probably introduced as a weed of broomcorn millet and was cultivated later, between the 10th and 8th c. BCE. During the LBA and EIA, it is always represented as a minor crop, restricted to some territories (Bouby *et al.* 2017).

Commune (département)	Site	Period	Cultural chronology (BCE)	Features (n)	Samples (n)	Volume (l)	Charred remains (n)	Waterlogged remains (n)	Investigator (Botany/Archaeology/Operator)
Buchères (10)	Parc logistique de l'Aube 2005-2008	EBA	Bronze A (2200-1600)	1	4	37	16		Toulemonde/Riquier/Inrap
Crévéchamps (54)	Tronc du chêne	EBA	Bronze A (2200-1600)	6	6	17	16		de Hingh/Koenig/Afan
Erstein (67)	Grasweg-PAE	EBA	Bronze A (2200-1600)	3	3	8.5	1378		Martinoli/Croutsch/Archéologie Alsace
Wahlenheim (67)	Lotissement de Wahlenheim	EBA	Bronze A (2200-1600)	1	3	30	13		Rousselet/Croutsch/Archéologie Alsace
Blénod-lès-Pont-à-Mousson (54)	Les longues Rayes	EBA	Bronze A2 (2000-1600)	2	16	119	179		Wiethold/Rachet/Inrap
Crévéchamps (54)	Tronc du chêne	EBA	Bronze A2 (2000-1600)	1	2	7.5	17		de Hingh/Koenig/Afan
Frouard (54)	Haut de Penotte	EBA	Bronze A2 (2000-1600)	17	17	53.3	68		de Hingh/Koenig/Afan
Frouard (54)	ZAC du Saule Gaillard I	EBA	Bronze A2 (2000-1600)	25	25	39.5	790		de Hing-Wiethold/Blouet/SRA
Vandières (54)	Les Grandes Corvées	EBA	Bronze A2 (2000-1600)	4	4	-	6		Zech-Matterne/Koenig/Afan-Inrap
Muttersholtz (67)	Rue des Cigognes	EBA	Bronze A2 (2000-1600)	3	3	9	6	2017	Rousselet/Pierrevelcin/Archéologie Alsace
Pagny-Les Goings (57)	Aéroport régional Lorraine	EBA/MBA	Bronze A-B (2200-1500)	4	4	29	99		de Hing-Wiethold/Blouet/SRA
Trémery (57)	Le Breuil/La Retienne	EBA/MBA	Bronze A-B (2200-1500)	1	1	10	9		Wiethold/Gérard/Inrap
Vignot (55)	Les Tranchons (2009)	EBA/MBA	Bronze A-B (2200-1500)	21	21	159	9		Wiethold/ Deffressigne/Inrap
Yutz (57)	Contournement sud-est	EBA/MBA	Bronze A-B (2200-1500)	1	1	7.5	68		de Hingh/Klag/Afan
Buchères (10)	Parc logistique de l'Aube 2012-2013	EBA/MBA	Bronze A2-B (2000-1500)	1	3	30	35		Toulemonde/Parésys/Inrap
Gougenheim (67)	Gingsheimer Feld	EBA/MBA	Bronze A2-B (2000-1500)	1	1	10	2		Durand/Thomas/Inrap
Duntzenheim (67)	Rainwasen	EBA/MBA	Bronze A-C (2200-1350)	1	1	10	127		Durand/Veber/Inrap
Erstein (67)	Parc d'Activités du Pays d'Erstein	MBA	Bronze B (1600-1500)	1	1	6		11,780	Bonnaire/Croutsch/Archéologie Alsace
Courceroy (10)	Les Dizaines	MBA	Bronze B-C (1600-1350)	1	1	15	147		Toulemonde/Lelarge/Archéopole
Duntzenheim (67)	Weingartenberg	MBA	Bronze B-C (1600-1350)	1	1	10	5		Durand/Veber/Inrap
Basse-Ham (57)	ZAC Kickelsberg	MBA	Bronze C (1500-1350)	2	2	20	7		Wiethold/Viller/Inrap
Ay-sur-Moselle (57)	Les Vélars Jacques	MBA	Bronze C (1500-1350)	1	1	20	20		Wiethold/Klag/Inrap
Blénod-lès-Pont-à-Mousson (54)	Les longues Rayes	MBA/LBA	MBA/LBA (1450-1300)	2	3	50	7		Wiethold/Rachet/Inrap
Couternon (21)	Larrey	MBA/LBA	MBA/LBA (1450-1300)	2	2	60	70		Wiethold/Labeaune/Inrap
Obernai (67)	Schulbach, Nouvel Hôpital	MBA/LBA	MBA/LBA (1450-1300)	2	4	36	29,873		Rousselet/Ferrier-Croutsch/Archéologie Alsace
Gingsheim (67)	Steinbrünnen, Aschenbuckel	LBA1	Bronze D (1350-1200)	1	1	10	150		Durand/Michler/Inrap
Balloy (77)	La Haute Borne	LBA1	Bronze D (1350-1200)	2	4	55	1254		Toulemonde/Samzung/Inrap

Table 3. The large archaeobotanical dataset from northeastern France used in this study.

Commune (département)	Site	Period	Cultural chronology (BCE)	Features (n)	Samples (n)	Volume (l)	Charred remains (n)	Waterlogged remains (n)	Investigator (Botany/Archaeology/Operator)
Erstein (67)	Parc d'Activités du Pays d'Erstein	LBA1	Bronze D (1350-1200)	1	1	9		3250	Bonnaire/Croutsch/Archéologie Alsace
Erstein (67)	Grasweg-PAE	LBA1	Bronze D (1350-1200)	1	1	6		876	Martinoli/Croutsch/Archéologie Alsace
Sézannes (51)	Maison de santé	LBA1	Bronze D (1350-1200)	1	6	52	617		Daoulas/Godard/Inrap
Hérange (57)	Gross Eichholz	LBA1	Bronze D (1350-1200)	1	12	225	370		Wiethold/Lafosse/Inrap
Jury (57)	ZAC de la Passerelle	LBA1	Bronze D (1350-1200)	1	1	10	9		Wiethold/Mangin/Inrap
Ay-sur-Moselle (57)	Les Vélars Jacques	LBA1	Bronze D (1350-1200)	3	3	32	40		Wiethold/Klag/Inrap
Meistratzheim (67)	Foegel	LBA1	Bronze D (1350-1200)	6	7	59	252		Wiethold/Veber/Inrap
Noyen-sur-Seine (77)	Carrière A2C	LBA1	Bronze D (1350-1200)	1	1	10	63		Toulemonde/Le Clézio/Évéha
Courceroy (10)	Les Dizaines	LBA1	Bronze D/Hallstatt A1 (1350-1100)	6	6	>30	38	978	Toulemonde/Lelarge/Archéopole
Réau (77)	Centre pénitencier	LBA1	Bronze D/Hallstatt A1 (1350-1100)	3	5	59	112		Toulemonde/Boulenger/Inrap
Réau (77)	Parc d'activités de l'A5	LBA1	Bronze D/Hallstatt A1 (1350-1100)	2	12	95	5212		Pradat/Froquet/Inrap
Vinneuf (89)	Le Chatelot	LBA1	Bronze D/Hallstatt A1 (1350-1100)	2	2	20	7		Toulemonde/Issenmann/Inrap
La Saulsotte (10)	Le Vieux Bouchy	LBA1	Bronze D/Hallstatt A1 (1350-1100)	3	3	30	22		Toulemonde/Peake/Inrap
Breuil (80)	Sole de l'épinette, Sole du Bois Marotte	LBA1	Bronze D/Hallstatt A1 (1350-1100)	1	1	?	24		Derreumaux/Baudry/Inrap
Entzheim et Geispolsheim (67)	Entzheim 2006	LBA1	Bronze D/Hallstatt A1 (1350-1100)	?	2	10	1383		Schaal/Landolt/Archéologie Alsace
Sainte Apollinaire (67)	Sur le Pré de Crot 1	LBA1	Bronze D/Hallstatt A1 (1350-1100)	2	2	59.6	131		Wiethold/Labeaune/Inrap
Bernolsheim/Mommenheim (67)	Plateforme Départementale d'Activités	LBA1	Bronze D/Hallstatt A1 (1350-1100)	1	1	40	14		Rousselet/Leprovost/Archéologie Alsace
Warcq (8)	Bois de Charnois	LBA1	Hallstatt A1 (1200-1100)	1	2	17	20		Toulemonde/Roseau/CG Ardennes
Buchères (10)	Parc logistique de l'Aube 2005-2008	LBA1	Hallstatt A1 (1200-1100)	3	3	28	34		Toulemonde/Riquier/Inrap
Marolles-sur-Seine (77)	Croix Saint Jacques	LBA1/LBA2	Hallstatt A1 (1200-1100)	1	1	?	546		Pradat/Peake/Inrap
Warcq (8)	Bois de Charnois	LBA1/LBA2	Hallstatt A1-A2 (1200-1000)	1	1	9	10		Toulemonde/Roseau/CG Ardennes
Courceroy (10)	Les Dizaines	LBA1/LBA2	Hallstatt A1-A2 (1200-1000)	5	7	80	196		Toulemonde/Lelarge/Archéopole
St Dizier (52)	Zone de référence 2012	LBA1/LBA2	Hallstatt A1-A2 (1200-1000)	1	1	20	8		Toulemonde/Bocquillon/Inrap
Saint Pierre du Perray (91)	TW 2008 et CSP 2010	LBA1/LBA2	Hallstatt A1-A2 (1200-1000)	6	22	153	1307		Toulemonde/Saron-Wagner/Inrap
Varennes-sur-Seine (77)	Le Volstin	LBA1/LBA2	Hallstatt A1-A2 (1200-1000)	1	1	10	27		Toulemonde/Noury/Évéha
Réau (77)	Parc d'activités de l'A5	LBA1/LBA2	Hallstatt A1-A2 (1200-1000)	10	22	228	1896		Pradat/Froquet/Inrap
St Dizier (52)	Zone de référence 2014	LBA1/LBA2	Hallstatt A (1200-1000)	6	6	38	728		Toulemonde/Bocquillon/Inrap
Berru (51)	La Maladrerie	LBA1/LBA2	Hallstatt A (1200-1000)	1	1	10	44		Toulemonde/Spies/Inrap

Commune (département)	Site	Period	Cultural chronology (BCE)	Features (n)	Samples (n)	Volume (l)	Charred remains (n)	Waterlogged remains (n)	Investigator (Botany/Archaeology/Operator)
Cernay-lès-Reims (51)	Le Bas de la Noue Saint Rémy	LBA2	Hallstatt A2 (1100-1000)	55	55	550	237		Daoulas/Rabaste/Inrap
Dieulouard (54)	Rue des Trappiers	LBA2	Hallstatt A2 (1100-1000)	1	1	2	70		Wiethold/Mangin/Inrap
Varois-et-Chaignot (21)	Le Pré du Plancher	LBA2	Hallstatt A2 (1100-1000)	4	5	180	87		Wiethold/Labeaune/Inrap
Vergigny (89)	La Grande Folie	LBA2	Hallstatt A2 (1100-1000)	1	1	20	495		Wiethold/Afan
Erstein (67)	Parc d'Activités du Pays d'Erstein	LBA2	Hallstatt A2 (1100-1000)	2	4	38	591	2660	Bonnaire/Crousch/Archéologie Alsace
Furdenheim (67)	Gruen, complexe judo-basket	LBA2	Hallstatt A2 (1100-1000)	2	4	10	424		Rousselet/Pierrevclin/Archéologie Alsace
Noyen-sur-Seine (77)	Carrière A2C	LBA2	Hallstatt A2 (1100-1000)	1	2	18	1762		Toulemonde/Le Clézio/Evéha
Vinneuf (89)	Le Chatelot	LBA2	Hallstatt A2-B1 (1100-900)	3	6	53	588		Toulemonde/Issenmann/Inrap
Courceroy (10)	Les Dizaines	LBA2	Hallstatt A2-B1 (1100-900)	8	9	79	240	1342	Toulemonde/Lelarge/Archéopole
Bréviandes (10)	Zac St Martin	LBA2	Hallstatt A2-B1 (1100-900)	1	1	20	36		Toulemonde/Laurelut/Inrap
Buchères (10)	Parc logistique de l'Aube 2005-2008	LBA2	Hallstatt A2-B1 (1100-900)	11	14	219	1511		Toulemonde/Riquier/Inrap
Pont-Sainte-Marie (10)	Rue Fernand Jaffiol	LBA2	Hallstatt A2-B1 (1100-900)	2	3	66	58		Toulemonde/ Millet/Inrap
Cormontreuil (51)	Les Grands Godets	LBA2	Hallstatt A2-B1 (1100-900)	3	5	91	72		Toulemonde/Bündgen/GrandReims
Loisy-sur-Marne (51)	Zac de la Haute Voie	LBA2	Hallstatt A2-B1 (1100-900)	6	6	49	282		Toulemonde/Toron/Evéha
St Dizier (52)	Zone de référence 2012	LBA2	Hallstatt A2-B1 (1100-900)	1	1	20	36		Toulemonde/Bocquillon/Inrap
Brie-Comte Robert (77)	Les-Prés-Le-Roi	LBA2	Hallstatt A2-B1 (1100-900)	9	9	90	260		Toulemonde/Peake/Inrap
Jaulnes (77)	Le Bas des Hauts Champs Ouest	LBA2	Hallstatt A2-B1 (1100-900)	6	6	120	955		Toulemonde/Peake/Inrap
Marolles-sur-Seine (77)	Ferme de la Muette	LBA2	Hallstatt A2-B1 (1100-900)	7	10	100	101		Toulemonde/Ferrier/Evéha
Noyen-sur-Seine (77)	Le Nord du Bois du Chêne	LBA2	Hallstatt A2-B1 (1100-900)	16	33	492	2025		Toulemonde/Nallier/Inrap
La Saulsotte (10)	Le Vieux Bouchy	LBA2	Hallstatt A2-B1 (1100-900)	3	3	43	46		Toulemonde/Peake/Inrap
Saint-Maure (10)	Culoison	LBA2	Hallstatt A2-B1 (1100-900)	3	3	28	15		Daoulas/Chauvin/Inrap
Blignicourt (10)	Les Voies de Brienne	LBA2	Hallstatt A2-B1 (1100-900)	5	6	58	318		Daoulas/Chauvin/Inrap
Les Grandes Chapelles (10)	La Taverne	LBA2	Hallstatt A2-B1 (1100-900)	3	10	38	6		Daoulas/Chauvin/Inrap
Ruvigny (10)	Proche l'Eglise	LBA2	Hallstatt A2-B1 (1100-900)	1	2	16	24		Daoulas/Roms/Inrap
Troyes (10)	Preize	LBA2	Hallstatt A2-B1 (1100-900)	1	3	26	7		Daoulas/Deborde/Inrap
Lieusaint (77)	Zac du Levant	LBA2	Hallstatt A2-B1 (1100-900)	7	19	163	1891		Pradat/Baguenier/Inrap
Basse-Ham (57)	ZAC intercommunale	LBA2	Hallstatt A2-B1 (1100-900)	7	12	165	888		Wiethold/Tikonoff/Inrap
Pévange (57)	Derrière le Pâtural	LBA2	Hallstatt A2-B1 (1100-900)	1	1	10	60		Wiethold/Viller/Inrap

Commune (département)	Site	Period	Cultural chronology (BCE)	Features (n)	Samples (n)	Volume (l)	Charred remains (n)	Waterlogged remains (n)	Investigator (Botany/Archaeology/Operator)
Coincy (57)	La réserve, Les Terres de Fer	LBA2	Hallstatt A2-B1 (1100-900)	1	1	10	1		Wiethold/Bernard/Inrap
Volstroff (57)	Les résidences de Volstroff	LBA2	Hallstatt A2-B1 (1100-900)	3	8	77	80		Wiethold/Brénon/Inrap
Guénange (57)	Carrière GSM	LBA2	Hallstatt A2-B1 (1100-900)	1	1	10	100		Wiethold/Viller/Inrap
Belleville-sur-Meuse (55)	La Pièce des vingt jours	LBA2	Hallstatt A2-B1 (1100-900)	1	1	10	212		Wiethold/Ramel/Inrap
Vignot (55)	Les Tranchons (2006)	LBA2	Hallstatt A2-B1 (1100-900)	2	2	29	9		Wiethold/Lefebvre/Inrap
Varois-et-Chaignot (21)	Le Pré du Plancher	LBA2	Hallstatt A2-B1 (1100-900)	11	12	226	1087		Wiethold/Labeaune/Inrap
Marly (57)	Le Clos des Sorbiers	LBA2	Hallstatt B1 (1000-900)	15	15	?	69		de Hingh/Klag/Afan
Yutz (57)	Contournement sud-est	LBA2	Hallstatt B1 (1000-900)	7	7	18.5			de Hingh/Klag/Afan
Art-sur-Meurthe (54)	L'Embanie – La Prairie de la Roanne	LBA2	Hallstatt B1 (1000-900)	1		7	70		Wiethold/ Deffressigne/ Inrap
Erstein (67)	Parc d'Activités du Pays d'Erstein	LBA2	Hallstatt B1 (1000-900)	3	4	38	494		Bonnaire/Croutsch/ Archéologie Alsace
Wissous (91)	Zone S/O de l'aéroport d'Orly	LBA2	Hallstatt B1 (1000-900)	3	3	23	1836		Toulemonde/Quenez/Inrap
Marly (57)	La Grange-aux-Ormes	LBA2	Hallstatt B1 (1000-900)	4	5	25	15		Wiethold/Tikonoff/Inrap
Longvic/Ouges (21)	Les Quétignières	LBA2	Hallstatt B1 (1000-900)	8	8	160	427		Wiethold/Labeaune/Inrap
Varois-et-Chaignot (21)	Le Pré du Plancher	LBA2	Hallstatt B1 (1000-900)	3	3	60	367		Wiethold/Labeaune/Inrap
Berru (51)	La Maladrerie	LBA2/ LBA3	Hallstatt B1-B2 (1000-800)	1	1	11	9		Toulemonde/Spies/Inrap
Flévy (57)	Zac de la Fontaine des Saints	LBA2/ LBA3	Hallstatt B1-B2 (1000-800)	1	1	5		132	Wiethold/Petitdidier/Inrap
Mondelange (57)	PAC de la Sente	LBA2/ LBA3	Hallstatt B1-B2/3 (1000-750)	3	3	30	5		Wiethold/Gazenbeek/Inrap
Changis-sur-Marne (77)	Les Pétreaux et La Pelle à Four	LBA3	Hallstatt B2 (900-800)	4	7	35	35		Zech-Matterne/Lafage/ Inrap
Bazancourt (51)	Sur les Petits Poissons	LBA3	Hallstatt B2/3 (900-750)	44	72	717	1867		Toulemonde/Desbrosses/ Inrap
Buchères (10)	Parc logistique de l'Aube 2012-2013	LBA3	Hallstatt B2/3 (900-750)	8	14	127	436		Toulemonde/Parésys/Inrap
Balloy (77)	La Haute Borne	LBA3	Hallstatt B2/3 (900-750)	3	6	95	112		Toulemonde/Samzung/ Inrap
Biaches (80)	Canal Seine Nord fouille 13	LBA3	Hallstatt B2/3 (900-750)	4	4	23	133		Toulemonde/Baudry/Inrap
Varennnes-sur-Seine (77)	Le Volstin	LBA3	Hallstatt B2/3 (900-750)	3	7	65	1092		Toulemonde/Noury/Evéha
Réau (77)	Centre pénitencier	LBA3	Hallstatt B2/3 (900-750)	6	6	81	174		Toulemonde/Boulenger/ Inrap
Pont-sur-Seine (10)	Gué Dehan zone 2	LBA3	Hallstatt B2/3 (900-750)	12	14	60	10,951		Schaal/Collas/Eveha
Pont-Marnay-sur-Seine (10)	La Gravière	LBA3	Hallstatt B2/3 (900-750)	8	9	111	64		Bonnaire/Fournand/Inrap
Beaumont-sur-Vesles (51)	Les Hauts de Chaillot	LBA3	Hallstatt B2/3 (900-750)	1	4	65	93		Zech-Matterne/Lamotte/ Inrap
Bussy-Lettrée (51)	Vatry Mont Lardon site 16	LBA3	Hallstatt B2/3 (900-750)	17	21	242	458		Zech-Matterne/Bailieux/ Inrap

Commune (département)	Site	Period	Cultural chronology (BCE)	Features (n)	Samples (n)	Volume (l)	Charred remains (n)	Waterlogged remains (n)	Investigator (Botany/Archaeology/ Operator)
Marly (57)	La Grange-aux-Ormes	LBA3	Hallstatt B2/3 (900-750)	4	6	56	2383		Wiethold/Tikonoff/Inrap
Trémery (57)	Le Breuil/La Retienne	LBA3	Hallstatt B2/3 (900-750)	2	21	161	178		Wiethold/Gérard/Inrap
Villeneuve-la-Guyard (89)	Les Champs Boissier	LBA3	Hallstatt B2/3 (900-750)	1	1	15	166		Wiethold/Afan
Changis-sur-Marne (77)	Les Pétreaux et La Pelle à Four	LBA3/EIA	Hallstatt B3 (900-750)	56	61	305	1849		Zech-Matterne/Lafage/ Inrap
Villiers-sur-Seine (77)	Le Gros Buisson	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	17	17	285	16,260		Toulemonde/Peake/Inrap
Plancy-l'Abbaye (10)	Saint Martin	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	6	7	67	93		Toulemonde/Moreau/Inrap
Buchères (10)	Parc logistique de l'Aube 2005-2008	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	52	66	992	4914		Toulemonde/Riquier/Inrap
Lesmont (10)	Pôle Scolaire	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	8	29	251	453		Toulemonde/Sanson/Inrap
Matignicourt-Goncourt (51)	Les Brouillards	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	2	7	113	99		Toulemonde/Richard/Inrap
Saint Léonard (51)	La Croix Chaudron	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	5	8	100	128		Toulemonde/Bündgen/ GrandReims
Saint-Martin-sur-le-Pré (51)	Rue des Castors	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	3	3	27	28		Toulemonde/Garmond/ GrandReims
Thillois (51)	La Croix Rouge	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	8	16	120	211		Zech-Matterne/Riquier/ Inrap
Réau (77)	Centre pénitencier	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	3	4	47	31		Toulemonde/Boulenger/ Inrap
Saint-Maure (10)	Culoison	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	10	10	90	450		Daoulas/Chauvin/Inrap
Vitry-sur-Seine (94)	Collège Monod	LBA3/EIA	Hallstatt B2/3-C1 (900-620)	1	14	241	21,597		Derreumaux/Maret/SA 94
Ville Saint Jacques (77)	Le Fond des Vallées/ Le Bois d'Echalas	LBA3/EIA	Hallstatt B3-C1 (900-620)	5	8	52	256		Toulemonde/Issenmann/ Inrap
Jaulnes (77)	Le Bas des Hauts Champs Ouest	LBA3/EIA	Hallstatt B3-C1 (900-620)	4	5	100	132		Toulemonde/Peake/Inrap
Florange (57)	Petit Biterfeld	LBA3/EIA	Hallstatt B3-C (900-620)	1	1	4	270		Wiethold/Franck/Inrap
Ley (57)	Le Grand Meix	LBA3/EIA	Hallstatt B3-C (900-620)	1	1	10	99		Wiethold/Mangin/Inrap
Sainte-Apollinaire (21)	Sur le petit Pré 1	LBA3/EIA	Hallstatt B3-C (900-620)	12	12	435	574		Wiethold/Labeaune/Inrap
Changis-sur-Marne (77)	Les Pétreaux et La Pelle à Four	LBA3/EIA	Hallstatt B3-C (900-620)	26	27	135	222		Zech-Matterne/Lafage/ Inrap




Figure 6. Proportions (in %) of the main categories of crops from the Early Bronze Age and Middle Bronze Age (EBA and MBA) settlements compared with the Late Bronze Age (LBA) settlements, divided by phases, calculated from the total number (nb) of charred grains in all sites of one phase. Mass finds are excluded from the calculations.

Discussion: Changes in agriculture and subsistence economy during the Late Bronze Age

Beside the introduction of millets, what else is new in the crop husbandry practices and subsistence economy of the LBA? The rare data for the EBA and MBA make comparisons difficult, but we did manage to collect convincing information that suggests more radical changes than just adding a new crop.

A flexible and resilient crop husbandry regime for an improved control of yields

The crop spectrum of the LBA includes a larger number of plants than that of the previous periods. It could feature up to nine cereals (hulled barley (*Hordeum vulgare* ssp. *vulgare*), bread wheat (*Triticum aestivum* s.l.), hard wheat/rivet wheat (*T. durum/turgidum*), emmer (*T. dicoccum*), einkorn (*T. monococcum*), spelt (*T. spelta*), Timopheev's wheat (*T. timopheevii* s.l.) – formerly known as ‘new’ glume wheat (cf. Czajkowska *et al.* 2020) – broomcorn millet (*Panicum miliaceum*) and foxtail millet (*Setaria italica*)); five pulses (grass pea (*Lathyrus cicera/sativus*), lentil (*Lens culinaris*), pea (*Pisum sativum*), bitter vetch (*Vicia ervilia*) and celtic bean (*Vicia faba* var. *minor*)); and three oil plants (gold-of-pleasure (*Camelina sativa*), flax (*Linum usitatissimum*) and opium poppy (*Papaver somniferum*)). Not only the crop spectrum, but also crop production and organisation changed. Pulses were more and more commonly cultivated and during LBA2 they became ubiquitous. For this period and the following one, they represent 16% of the charred grains in assemblages, compared with being almost completely absent from Early and Middle Bronze Age contexts (Fig. 6). At least 9 out of 10 rural settlements cultivated legumes alongside cereals. Oil plants were also much more frequent, especially gold-of-pleasure. They are recorded in one of three settlements during LBA2 and LBA3. Altogether, evidence for plant cultivation during the EBA-MBA is poor, with an average of only 2 different crop species per settlement (and a maximum of 5), and it sharply increases during the LBA, corresponding to an average of 6 crops per settlement, for a median of 5 and a maximum of 15. Most LBA farms cultivated a larger and more diversified set of crops than did farms during the earlier phases of the Bronze Age.

A larger crop spectrum meant more complexity but also more flexibility in crop management. Most of the new plants introduced and/or cultivated, such as millets, pulses and gold-of-pleasure, were summer crops. Because they are frost intolerant, they had to be sown in the spring, whereas hulled wheats were mainly sown in

winter. This allowed a better distribution of fieldwork throughout the year and it permitted inclusion of catch-up crops, especially millets, which are fast-growing plants. When winter crops failed, one could still sow millet in June and harvest it in September. Another greatly appreciated technical characteristic of millet in case of bad harvest years is that for 1 grain of millet sown, 200 are harvested (Sigaut 1995). This is 16 to 50 times higher than in wheat. This meant that almost the entire harvest of millet could be consumed; one did not need to save a substantial part of it for future sowing.

If we sum up the characteristics of the new crop husbandry regime of the LBA, we have all the elements of a resilient and sustainable system, able to maintain soil fertility and therefore yields. It includes crop diversity, with 17 available crop species and evidence that indeed up to 10 or 11 species were cultivated in a given year in one settlement (Toulemonde 2013). Hardy plants with a low demand in manure (namely hulled wheats and barley) were selected as staple crops. The focus was not on high-yield plants, such as bread wheat. What mattered was the performance in difficult times and places, in case of drought, frost or poor soils. Flexibility, with summer and winter crops and catch-up crops, was another characteristic of the LBA system.

Moreover, many of the cultivated plants complement each other agronomically, making possible (likely?) maslins (of different species of wheats), rotations (pulses/cereals), or associations of plants (pea or celtic bean and gold-of-pleasure). Gold-of-pleasure also offered the possibility of controlling weeds when cultivating pulse crops, thanks to its horizontally spreading root system. This once well-known and then forgotten characteristic is being rediscovered in modern organic farming (Rozier 1821-1823; de Carné-Carnalet 2011; Mellon and Bollot pers. comm.).

Manuring was possible in two ways. As mixed farming – crop and animal husbandry – was the rule in northeastern France (Auxiette *et al.* 2015; Auxiette and Toulemonde 2018; Toulemonde *et al.* 2018), some animal manure was available, but it may not always have been sufficient. The rotation of cereals with abundantly cultivated pulses and cruciferous plants, such as gold-of-pleasure, could have also helped maintain fertility and nutrient levels. Thanks to a symbiotic association with bacteria of the genus *Rhizobium*, pulses can fix atmospheric nitrogen in the soil, while gold-of-pleasure is rich in potassium, of which some pulses, such as celtic beans, are in high demand (de Carné-Carnalet 2011).

All these characteristics are strategies to secure production and help farming resilience. They seem to have been successfully applied, because the population size rose sharply during the LBA. Analyses of weed flora needs in terms of nutrients using a nitrogen indicator for the French territory (Julve 1998), similar to that of Ellenberg (1988) for central Europe, are available for groups of settlements from different parts of the Champagne and Île-de-France regions, dated to the LBA and EIA (Toulemonde 2013). They indicate that during the nine centuries of the LBA and EIA (1350-475 BCE), when the same farming system was in use, there was no significant decrease in soil fertility.

This multiple cropping system, with its summer and winter crops, also entailed intensive farming year-round. Millets and pulses are plants that require regular care. The cultivation of millet only succeeds through repeated hoeing and weeding, meaning a high investment in labour at different times of the cycle, and especially during field preparation, the growth period and the harvest (Sigaut 1995). This significant labour cost was even the cause of its abandonment in some French regions in the 19th century (Bonnain 1995). The intensive fieldwork required throughout the year to take care of this broad diversity of plants with different growing periods and needs could only be provided by a sedentary population. If a nomadic life was practised during the EBA and MBA, this was no longer an option with the crop husbandry regime of the LBA. Consequently, domestic and agricultural features, such as storage pits and granaries, became much more visible and land use increased and diversified, including farming of the plateaus and chalklands alongside river valleys (see above, section on archaeological background).

Gathering: A less critical resource

As crop production increased and diversified, gathering became less crucial than it had been in previous phases of the Bronze Age. During the EBA, consumption of acorns (*Quercus* sp.), a major wild source of carbohydrates, was quite important (Coquillat 1959; Vencl 1985, 1996; Karg and Haas 1996). In Lorraine, charred acorns are even considered as an index fossil for this period. There are many examples of EBA features filled with charred acorns. The settlement at Richemont-Berg VI yielded 348 acorns from a posthole dated to the end of the EBA-beginning of the MBA; at Frouard-Zac du Saule Gaillard, 722 acorns were discovered in the fills of postholes dated to the 20th-17th c. BCE (de Hingh 2000, 94); and at Yutz-Contournement sud-est, a storage pit from the beginning of the EBA yielded 50 charred acorns (de Hingh 2000, 115).

During the LBA, the gathering of acorns declined in northeastern France (Bouby *et al.* 2017). In the Upper Seine Valley, although oaks were less ubiquitous than during the EBA-MBA, they were still available near settlements, and their wood was commonly used (Leroy 2012; Chaussé *et al.* 2019). Acorn is, however, only the third most important collected wild resource, far behind hazelnut (*Corylus avellana*) and sloe (*Prunus spinosa*) (Toulemonde 2013). It seems that farming resources were at that stage sufficient to feed the population.

An overview of changes in agriculture and European convergences

The arrival of broomcorn millet in northeastern France coincides with the development of a new farming system, characterised by global strategies to secure production. This system sustained an increase in population and activities during the LBA. As glimpsed in the demographics – sluggishness then surge – changes in diet and farming practices during the LBA are proved to be a European phenomenon, which occurred at a varying pace depending on the region (*e.g.* Stika and Heiss 2013; Valamoti 2016; Bouby *et al.* 2017; Effenberger 2018; Liu *et al.* 2018; Reed 2020). It was widespread to the point of being described as ‘*the third food revolution*’ (Kneisel *et al.* 2015b). Not all references in the archaeobotanical literature can be cited here, but it is intriguing to note that in many places across Europe, from southern Italy to Bohemia, and in northeastern France, acorns are an index fossil of the EBA. They become less frequent in LBA contexts in northern Europe (Kneisel *et al.* 2015a), while gathering continues in Mediterranean regions, where it may have concerned *sempervirens* oak species, which produce fruit with lower tannin levels (Bouby 2014; Pêche-Quilichini *et al.* 2020).

More common trends in LBA farming strategies in Europe have been listed by Kneisel *et al.* (2015a). Many of them, such as the increase in crop diversity, the spread of millets, the more widespread use of pulses, the emergence of gold-of-pleasure, *etc.*, can also be observed in northeastern France. Results of the analyses of weed flora demands for nitrogen in southwestern Germany using a similar indicator to the one used for northeastern France show no significant change in soil fertility during the LBA and EIA, indicating a sustainable farming system, as observed in Champagne and Île-de-France, and suggesting the use of fertilisers (Rösch *et al.* 2015).

It is tempting to link the demographic surge of the LBA to the adoption of a new agriculture, which seemed to offer all the guarantees for a sufficient crop production and a balanced diet (Rosenstock *et al.* 2015). However, one should first understand the reasons for the apparent sluggishness of demography observed in northeastern France from the Late Neolithic to the end of the MBA. Before that period, during Early Neolithic and Middle Neolithic, other farming systems had also been successful (Bouby *et al.* 2018).

Farming, processing and uses of millet

Ethnohistory and ethnobotany teach us that, in a not-so-distant past, before being entirely replaced by maize in the 1960s, broomcorn millet was cultivated as a minor crop on every farm of the département of Vendée in western France, as well as in other areas (Hörandner 1995). Farming and processing techniques were specific to the plant due to its characteristics, which are quite different from those of other cereals (Hongrois 1995; Sigaut 1995; Lundström-Baudais *et al.* 2002). Millet was used to feed humans and poultry. People consumed it as a daily dish, but it was also on the menu for harvest festivals and wedding meals in rural areas (Hongrois 1995; Touzeau 1995). In other parts of Europe, in Portugal or Hungary for example, feasts and collective celebrations, such as carnivals, funerals, festivals of the dead, *etc.*, also featured millet in traditional dishes (Barboff 1995; Kishán 1995). Considering this documentation, what can archaeology and archaeobotany teach us about the farming and social practices of a more distant past?

Farming and processing techniques

During the LBA, broomcorn millet was cultivated on every small farmstead, regardless of the type of soil. It does not seem that poor soils were preferred, as remains are more frequent and abundant in the fertile Seine Valley than on the chalklands of Champagne (Toulemonde 2013). We have no clues as to how millets were harvested. There is no existing study on agricultural tools for the LBA and EIA in northeastern France, due to the finds being so scarce (Toulemonde *et al.* 2017). Copper alloy sickles appear at the end of the Bronze Age (14th-13th c. BCE), but it is difficult to link them to millet more than to other cereals. Moreover, these tools disappear during the EIA, a period when millet is still intensively cultivated. Evidence on storage conditions is just as poor. The first mass finds of millets date to LBA3, and more have been discovered from the EIA, but none of them are of charred stock preserved in situ. Sometimes mixed with other cereals, they come from fills of waste pits (Villiers-sur-Seine; Peake 2020) or from middens (Pont-sur-Seine-Le Gué Déhan; Collas *et al.* 2018).

The dehusking of millet seems to be a mandatory step before consumption, as their glumes are siliceous and hard to digest. It is not clear if this was systematically done in ancient times for some specific contexts (Jacquat 1989; Boenke 2003; Behre 2008), however ethnobotany, ethnohistory and the history of techniques from all over the world always refer to millet dehusking, or at least to the removal of the hulls in one way or another (Comet 1992; Hörandner 1995; Murty and Kumar 1995; ICRISAT and FAO 1996; Lundström-Baudais *et al.* 2002; Lundström-Baudais 2010; Hamon and Le Gall 2013; Moreno-Larrazabal *et al.* 2015; Bhandari *et al.* 2020). This is sometimes facilitated by the heating of the grains, a treatment also practised worldwide for broomcorn millet, although it does not seem necessary for foxtail millet (Lundström-Baudais *et al.* 2002). Charred lumps of aggregated kernels of millet in their glumes were discovered all over the settlement area of Villiers-sur-Seine, suggesting recurrent processing or food preparation (Auxiette *et al.* 2015). Experiments have shown that, if not the only explanation, heat treatment prior to dehusking is a more likely hypothesis for these aggregates than the cooking of porridge (Toulemonde 2014). Finds of aggregates of entire kernels of millet on the perforated deck of specific ovens (known as Sévrier-oven types) have also been linked to heat treatment (Coulon 2015). These discoveries were made in the Alps, but the same type of oven was also found at the settlement of Buchères, Champagne.

Daily dish, festive dish

As testified by its ubiquity in the assemblages from LBA and EIA settlements, broomcorn millet was most probably consumed as an everyday meal. More information on diet is given by stable isotope studies of carbon and nitrogen ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$). For northeastern France, these studies are just emerging. LBA funerary contexts from the site of Barbuise, Champagne, have been studied and compared with the nearby Neolithic contexts at Gurgy, Burgundy. Stable isotope analyses were performed on bone and tooth collagen and apatite from eight humans and five domestic animals dating to ca. 1300 BCE. Results indicate the consumption of ^{13}C -enriched food by humans and some animals (dog and pig) during the LBA, compared with the Neolithic. The nitrogen ratio is lower, suggesting a diet based more on plants than before (Goude *et al.* 2016). These results are in line with the archaeobotanical studies.

Millet not only contributed to the daily diet, it was also part of festive menus. The multidisciplinary study of the site of Villiers-sur-Seine has demonstrated that collective feasting was a regular feature at this fortified settlement (Auxiette *et al.* 2015; Auxiette and Peake 2020; Peake 2020). Large quantities of meat from animals slaughtered at specific times of the year were consumed during these celebrations. Abundant millet remains are associated with animal bones (Toulemonde 2020). For these festive meals, the cereal was perhaps prepared in a different way than the daily meal, symbolising this ceremony or ritual, as was the case in other societies where millet was a traditional crop (Barboff 1995; Bonnain 1995).

There is evidence that broomcorn millet, important for the living, also accompanied the dead as a funerary offering. The rich elite grave with weapons found in the Tumulus 'Géraud' at Saint-Romain de Jalionas, département of Isère, French Alps, contained an LBA3 bronze situla filled with dehusked grains of broomcorn millet (Verger and Guillaumet 1988; plant macrofossils – Wiethold unpublished data).

After the Bronze Age, at a time when archaeobotanical studies show a sharp decline in the consumption of millet (Toulemonde *et al.* 2017; Zech-Matterne *et al.* 2017), the plant still played a role in some rituals. Evidence for this comes from southern Burgundy, where dozens of ceramic vessels dated to the Late Iron Age and the Roman period (prior to the 1st c. CE) have been recovered from the Saône River. The vessels have standardised shapes and some of them contained the remains of a charred organic matter in which entire kernels of broomcorn millet were visible. Because of their location, in fords or ports, these finds have been interpreted as cereal offerings to the gods, either as porridge or as beer (Marinval and Bonnamour 2010). Other evidence comes from the settlement of Warcq-la Sauce, in the French Ardennes. A rich chariot burial dating to the 2nd c. BCE included a food-related deposit with five different pottery vessels. The biochemical analysis performed on all vessels revealed traces of miliacin, with significant concentrations in three of them, leading to interpret them as the containers for food or beverage offerings made from millet (Saurel *et al.* 2021).

These offerings of millet are surprising for the periods when the cereal seemed to be no longer in vogue. An ethnobotanical survey evokes a similar case for the 19th century in southern France, where millet, which had been replaced by maize in the human diet, was still eaten as a ceremonial meal 'to ensure good relations between the community of the dead and that of the living' (Bonnain 1995).

Overall conclusions and outlook

An overview of Bronze Age archaeobotanical data from northeastern France was combined with a series of radiocarbon dates obtained on charred caryopses of broomcorn millet. Some remains could not be dated due to the low carbon content, while others proved to be younger than expected. Results obtained so far suggest

that millet did not arrive in the area before the 14th c. BCE, which is consistent with the conclusions of the recent AMS-radiocarbon dating programme carried out to track the arrival of millet on a European level (Filipović *et al.* 2020).

The introduction of this summer crop is at the heart of a new farming system, characterised by global strategies to secure production and maintain soil fertility. The surge in demographics that followed in northeastern France and in other parts of Europe was therefore sustained by a new diet. Millets were indeed the emblematic plant of food identity at the end of the European Bronze Age. In northeastern France, they contributed to the everyday diet and were also an important part of collective celebrations. Never again have they played such a predominant role in the human diet. They started to decline during the Late Iron Age, from the 2nd c. BCE onwards, when innovations in farming tools led to the extension of cultivated areas and the development of a more specialised agriculture (Zech-Matterne and Brun 2016; Toulemonde *et al.* 2017).

Research perspectives on the role of millets in prehistoric France are numerous. They should first include a new series of radiocarbon dating to consolidate and refine results in relation to the introduction of these cereals, by tracking at a regional level their spread from the neighbouring countries (southern Germany, Switzerland, northern Italy) across the current French territory. Stable isotope studies are developing, and several have already been undertaken within the framework of a collective research project whose goal is to track mobility and changes/social differentiation in paleodiet at the transition between the MBA and LBA in the Upper Seine Valley (PCR BronzPal, led by Rebecca Peake, Inrap). Concerning the food evidence, plenty of material is available in the form of aggregated kernels of millet from various contexts, as described in this article. In the future, it could serve as the basis of a major project, aimed at combining experimentation on beer making and other food preparations with scanning electron microscope (SEM) analysis of these remains.

Acknowledgements

Radiocarbon dating of millets could be performed at the Poznań Radiocarbon Laboratory (Poznań, Poland) thanks to the financial support of the UMR 7209 AASPE MNHN-CNRS. Pictures of the millets were taken using the laboratory equipment of the UMR 7209 in Paris.

We also thank Rebecca Peake for kindly proofreading this article completely. We are grateful to Mr Mellon and Mr Bollot, organic farmers in Picardy, for sharing their experience in growing millet.

References

- Auxiette, G. and Toulemonde, F., 2018. L'intensification agricole de la fin de l'âge du Bronze et du premier âge du Fer: entre pratiques familiales et pratiques communautaires. In: N. Dohrmann and V. Riquier, eds., *Archéologie dans l'Aube: des premiers paysans au prince de Lavau, 5300 à 450 avant notre ère*. Gent: Snoeck, 177-185.
- Auxiette, G. and Peake, R., 2020. Festoyer à Villiers-sur-Seine (Seine-et-Marne) – Pratiques commensales dans un habitat aristocratique à la fin de l'âge du Bronze. In: G. Auxiette, C. Mougne, R. Peake, F. Toulemonde, eds., *Autour de la table: l'alimentation à l'âge du Bronze et au premier âge du Fer*. Actes de la Journée thématique du 3 mars 2017 à Saint-Germain-en-Laye. Bulletin de l'Association pour la Promotion des Recherches sur l'Âge du Bronze, suppl. 6, 137-154.
- Auxiette, G., Peake, R., Toulemonde, F., 2015. Food production and diet during the Late Bronze Age in the Upper Seine Valley (France). In: J. Kneisel, M. Dal Corso, W. Kirleis, H. Scholz, N. Taylor, V. Tiedtke, eds., *The Third Food Revolution? Setting the Bronze Age Table: Common Trends in Economic and Subsistence Strategies in*

- Bronze Age Europe*. Proceedings of the International Workshop ‘Socio-Environmental Dynamics over the Last 12,000 Years: The Creation of Landscapes III’ (15th-18th April 2013) in Kiel. Universitätsforschungen zur prähistorischen Archäologie 283. Bonn: Dr. Rudolf Habelt GmbH, 47-61.
- Barboff, M., 1995. Le millet au Portugal. In: E. Hörandner, ed., *Millet – Hirse – Millet*. Actes du Congrès d’Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang, 113-121.
- Bhandari, G.R., Gauchan, D., Bhandari, B., Joshi, B.K., Panta, S., 2020. Advancement, simplification and piloting of electrical proso millet de-husker (*Chino Kutak*) for the mountain farmers. In: D. Gauchan, B.K. Joshi, B. Bhandari, H.K. Manandhar, D.I. Jarvis, eds., *Traditional Crop Diversity for Mountain Food and Nutrition Security in Nepal: Tools and Research Results of the UNEP GEF Local Crop Project, Nepal*. Kathmandu, Nepal: NAGRC, LI-BIRD and the alliance of Biodiversity and CIAT, 75-81.
- Behre, K.E., 2008. Collected seeds and fruits from herbs as prehistoric food. *Vegetation History and Archaeobotany*, 17, 65-73. <https://link.springer.com/article/10.1007/s00334-007-0106-x>
- Boenke, N., 2003. Organic resources: food supply and raw materials of the miners. *The Antiquaries Journal*, 83, 147-149.
- Bonnain, R., 1995. Le pain des morts. In: E. Hörandner, ed., *Millet – Hirse – Millet*. Actes du Congrès d’Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang, 187-196.
- Bouby, L., 2014. *L’agriculture dans le Bassin du Rhône du Bronze final à l’Antiquité, agrobiodiversité, économie, cultures*. Toulouse: Archives d’Ecologie préhistorique.
- Bouby, L., Zech-Matterne, V., Bouchette, A., Cabanis, M., Derreumaux, M., Dietsch-Sellami, M.F., Durand, F., Figueiral, I., Marinval, P., Pradat, B., Rousselet, O., Rovira, N., Schaal, C., Toulemonde, F., Wiethold, J., 2017. Ressources et économie agricole en France à l’âge du Bronze et au premier âge du Fer: les données carpologiques. In: L. Carozza, C. Marcigny, M. Talon, eds., *L’habitat et l’occupation des sols à l’âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS, 299-326.
- Bouby, L., Dietsch-Sellami, M.F., Martin, L., Marinval, P., Wiethold, J., 2018. Ressources végétales et économie de subsistance au Néolithique en France (6000-2000 av. J.-C.). In: J. Guilaine and D. Garcia, eds., *La Protohistoire de la France*. Paris: Hermann, 141-152.
- Bronk Ramsey, C., 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon*, 51, 337-360. <https://doi.org/10.1017/S0033822200033865>
- Bronk Ramsey, C., 2017. Methods for summarizing radiocarbon datasets. *Radiocarbon*, 59, 1809-1833. <https://doi.org/10.1017/RDC.2017.108>
- Buchez, N., Henton A., Le Guen, P., Talon, M., 2017. L’habitat et l’occupation du sol en Nord Picardie. In: L. Carozza, C. Marcigny, M. Talon, eds., *L’habitat et l’occupation des sols à l’âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS, 33-52.
- Capuzzo, G., Zanon, M., Dal Corso, M., Kirleis, W., Barcelo, J.A., 2018. Highly diverse Bronze Age population dynamics in central-southern Europe and their response to regional climatic patterns. *PLoS ONE*, 13 (8), e0200709. <https://doi.org/10.1371/journal.pone.0200709>
- de Carné-Carnavalet, C., 2011. *Agriculture biologique, une approche scientifique*. Las-say-les-Châteaux: France Agricole.
- Carozza, L., Marcigny, C., Talon, M., eds., 2017a. *L’habitat et l’occupation des sols à l’âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS.
- Carozza, L., Marcigny, C., Talon, M., with the collaboration of F. Audouit and C.A. Perdu, 2017b. Introduction. In: L. Carozza, C. Marcigny, M. Talon, eds., *L’habitat et l’occupation des sols à l’âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS, 13-31.

- Chaussé, C., Leroyer, C., Allenet de Ribemont, G., Perrière, J., Coubray, S., 2019. Une montille bordée d'eau. In: R. Peake, ed., *Villiers-sur-Seine, un habitat aristocratique du IXe siècle avant notre ère*. Recherches Archéologiques 18. Paris: Inrap/CNRS, 19-78.
- Collas, R., Brenot, J., Donnart, K., Fronteau, G., Lachiche, C., Mauduit, C., Noury, A.C., Paradis-Grenouillet, S., Pescher, B., Poudroux, S., Roscio, M., Schaal, C., Siquin, J.-B., 2018. *Pont-sur-Seine (10), Le Gué Dehan, zone 2*. Limoges: Evéha, SRA Grand Est.
- Comet, G., 1992. *Le paysan et son outil: essai d'histoire technique des céréales (France VIIIe-XVe siècles)*. Collection de l'Ecole Française de Rome 165. Rome: Ecole Française de Rome.
- Coquillat, M., 1959. Les glands du chêne dans l'alimentation des hommes préhistoriques. *Bulletin Mensuel de la Société Linnéenne de Lyon*, 28 (1), 28-32.
- Coulon, J., 2015. Les fours dit de 'potier' de type Sévrier, Haute-Savoie, France: les indices d'une hypothèse fonctionnelle alternative. In: S. Boulud-Gazo and T. Nicolas, eds., *Artisanats et productions à l'âge du Bronze*. Actes de la Journée de la Société Préhistorique Française, Nantes, 8 octobre 2011. Séances de la Société Préhistorique Française 4. Dijon: Société Préhistorique Française, 109-121.
- Courel, B., Schaeffer, P., Adam, P., Motsch, E., Ebert, Q., Moser, E., Féliu, C., Bernasconi, S.M., Hajdas, I., Ertlen, D., Schwartz, D., 2017. Molecular, isotopic and radiocarbon evidence for broomcorn millet cropping in northeast France since the Bronze Age. *Organic Geochemistry*, 110, 13-24. <https://doi.org/10.1016/j.orggeochem.2017.03.002>.
- Croutsch, C., Bonnaire, E., Kucharski, M., Leprovost, C., Pascutto, E., Pélissier, A., Philippe, M., Putelat, O., Rault, E., Tegel, W., Véber, C., Wuscher, P., 2016a. *Erstein, Bas-Rhin, Parc d'Activités du Pays d'Erstein, tranche 2: Evolution d'un terroir entre la fin du IIIe et le début du Ier millénaire avant notre ère*. Sélestat: Pôle d'Archéologie Interdépartemental Rhénan.
- Croutsch, C., Bonnaire, E., Kucharski, M., Leprovost, C., Pascutto, E., Pélissier, A., Philippe, M., Putelat, O., Rault, E., Tegel, W., Véber, C., Wuscher, P., 2016b. *Erstein, Bas-Rhin, Parc d'Activités du Pays d'Erstein, tranche 2: Evolution d'un terroir entre la fin du IIIe et le début du Ier millénaire avant notre ère*. Sélestat: Pôle d'Archéologie Interdépartemental Rhénan.
- Czajkowska, B.I., Bogaard, A., Charles, M., Jones, G., Kohler-Schneider, M., Mueller-Bieniek, A., Brown, T.A., 2020. Ancient DNA typing indicates that the 'new' glume wheat of early Eurasian agriculture is a cultivated member of the *Triticum timopheevii* group. *Journal of Archaeological Science*, 1231, 105258. <https://doi.org/10.1016/j.jas.2020.105258>
- Deffressigne, S., Tikonoff, N., Boulanger-Bouchet, K., Chaussée, K., Tesnier-Hermetey, C., 2002. Les gisements d'habitat de la fin du premier âge du Fer à Gondreville Fontenoy-sur-Moselle (54): le stockage intensif et ses conséquences économiques et sociales. *Archaeologia Mosellana*, 4, 81-184.
- Desbrosse, V. and Riquier, V., 2012. Les établissements ruraux palissadés hallstattiens en Champagne. In: M. Schönfelder and S. Sievers, eds., *L'âge du Fer entre la Champagne et la vallée du Rhin*. 34^e colloque international de l'Association Française pour l'Étude de l'âge du Fer, Aschaffenburg 13-16 Mai 2010. RGZM-Tagungen 14. Mainz: Römisch-Germanisches Zentralmuseum, 3-27.
- Ducreux, F., 2017. L'occupation des sols et l'habitat en Bourgogne: les départements de la Côte-d'Or, de la Saône-et-Loire, la Nièvre et l'Yonne. In: L. Carozza, C. Marcigny, M. Talon, eds., *L'habitat et l'occupation des sols à l'âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS, 275-298.
- Dupéré, B., Verrier, G., Allard, P., Avival, F., Bandelli, A., Bardel, D., Dubreucq, E., Duda, D., Fechner, K., Filipiak, B., Frouin, M., Gauvain, F., Loiseau, S., Meunier, K., Monchablon, C., Nicolas, T., Sauvin, C.E., Souffi, B., Thiol, S., Toulemonde, F., Van Es, M., 2013. *Pont-sur-Seine, Aube, 'La Gravière': une occupation mésolithique, des*

- fosses néolithiques et des installations agricoles de l'âge du Bronze et du début du Second âge du Fer: Rapport Final d'Opérations*. Châlons-en-Champagne: Inrap/SRA Champagne-Ardenne.
- Effenberger, H., 2018. The plant economy of the northern European Bronze Age – more diversity through increased trade with southern regions. *Vegetation History and Archaeobotany*, 27, 65-74. <https://link.springer.com/article/10.1007/s00334-017-0621-3>
- Ellenberg, H., 1988. *Vegetation Ecology of Central Europe*, translated by G.K. Strutt. Cambridge: Cambridge University Press.
- Filipović, D., Meadows, J., Wiethold, J., Jahns, S., Bittmann, F., Kirleis, W., 2018. Before and after: millet cultivation and the transformation of prehistoric crop production in northern Germany. *Antiquity*, 92 (365). <https://doi.org/10.15184/aqy.2018.225>
- Filipović, D., Meadows, J., Corso, M.D., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Klooss, S., Kolher-Schneider, M., Kroll, H., Makarowics, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A.M., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, L., Przybyla, M., Reed, K., Rennwanz, J., Stika, H.P., Stobbe, A., Tolar, T., Wasylkowa, K., Wiethold, J., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Goude, G., Rey, L., Toulemonde, F., Cervel, M., Rottier, S., 2016. Dietary changes and millet consumption in northern France at the end of prehistory: evidence from archaeobotanical and stable isotope data. *Environmental Archaeology*, 22 (3), 268-282. <https://doi.org/10.1080/14614103.2016.1215799>
- Guilaine, J. and Garcia, D., eds., 2018. *La Protohistoire de la France*. Paris: Hermann.
- Hamon, C. and Le Gall, V., 2013. Millet and sauce: the uses and functions of querns among the Minyanka (Mali). *Journal of Anthropological Archaeology*, 32, 109-121. <https://www.sciencedirect.com/science/article/abs/pii/S0278416512000578>
- Herrscher, E., André, G., Bodet, C., Chataigner, C., Decaix, A., Goude, G., Hamon, C., Le Mort, F., Lyonnet, B., Martin, L., Messenger, E., Oberlin, C., Ollivier, V., Poulmarc'h, M., Sermet, C., Vila, E., 2018. The origins of millet cultivation in the Caucasus: archaeological and archaeometric approaches *Préhistoires Méditerranéennes*, 6. <http://journals.openedition.org/pm/1367>
- de Hingh, A.E., 2000. *Food Production and Food Procurement in the Bronze Age and Early Iron Age (2000-500 BC): the organisation of a diversified and intensified agrarian system in the Meuse-Demer-Scheldt region (The Netherlands and Belgium) and the region of the river Moselle (Luxemburg and France)*. Archaeological Studies Leiden University 7. Leiden: Leiden University.
- Hongrois, M., 1995. Do mell en Vendée ou culture et consommation du mil en Vendée, dans le canton de la Chataigneraie. In: E. Hörandner, ed., *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang, 71-86.
- Hörandner, E., ed., 1995. *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang.
- Hunt, H.V., Vander Linden, M., Liu, X., Motuzaite-Matuzeviciute, G., Colledge, S., Jones, M.K., 2008. Millets across Eurasia: chronology and context of early records of the genera *Panicum* and *Setaria* from archaeological sites in the Old World. *Vegetation History and Archaeobotany*, 17, 5-18. <https://doi.org/10.1007/s00334-008-0187-1>
- Hunt, H.V., Badakshi, F., Romanova, O., Howe, C.J., Jones, M.K., Heslop-Harrison, J.S.P., 2014. Reticulate evolution in *Panicum* (Poaceae): the origin of tetraploid broomcorn millet, *P. miliaceum*. *Journal of Experimental Botany*, 65, 3165-3175. <https://doi.org/10.1093/jxb/eru161>

- Hunt, H.V., Rudzinski, A., Jiang, H., Wang, R., Thomas, M.G., Jones, M.K., 2018. Genetic evidence for a western Chinese origin of broomcorn millet (*Panicum miliaceum*). *The Holocene*, 28 (12), 1968-1978. <https://doi.org/10.1177%2F0959683618798116>
- ICRISAT (International Crops Research Institute for the Semi-Arid Tropics) and FAO (Food and Agriculture Organization of the United Nations), 1996. *The World Sorghum and Millet Economies: Facts, Trends and Outlook*. <http://www.fao.org/docrep/W1808E/w1808e00.HTM>.
- Issenmann, R., Ameye, N., Auxiette, G., Bardel, D., Bedault, L., Bertin, P., Bertrand, I., Bostyn, F., Delattre, V., Meunier, K., Peake, R., Pilon, F., Praud, I., Séguier, J.M., Sidera, I., Toulemonde, F., Wiethold, J., 2009. *Ville-Saint-Jacques 'Le Fond des Vallées' / 'Le Bois d'Échalas' (Seine-et-Marne, Île-de-France), Occupations du Néolithique ancien, du Bronze final, du Hallstatt moyen, du Hallstatt final, de La Tène ancienne et de l'époque gallo-romaine*. Paris: Inrap/SRA Île-de-France.
- Jacquat, C., 1989. Hauterive-Champréveyres 2: les plantes de l'âge du Bronze: contribution à l'histoire de l'environnement et de l'alimentation. *Archéologie Neuchâteloise* 8. Saint Blaise: Edition du Ruau.
- Jones, M.K., 2004. Between fertile crescents: minor grain crops and agricultural origins. In: M.K. Jones, ed., *Traces of Ancestry: Studies in Honour of Colin Renfrew*. McDonald Institute Monographs. Cambridge: McDonald Institute for Archaeological Research, 127-135.
- Julve, P., 1998. Baseflor: index botanique, écologique et chorologique de la flore de France. Version: '21 Juillet 2011'. <http://perso.wanadoo.fr/philippe.julve/catminat.htm>
- Karg, S. and Haas, J.N., 1996. Indizien für den Gebrauch von mitteleuropäischen Eicheln als prähistorische Nahrungsressource. In: I. Comper, J. Hahn, M. Uerpman, eds., *Spuren der Jagd – Die Jagd nach Spuren: Festschrift für Hans-Jürgen Müller-Beck*. Tübinger Monographien für Urgeschichte 11. Tübingen: Mo Vince Verlag, 429-435.
- Kisbán, E., 1995. Millet in the hierarchy of meals in Hungary. In: E. Hörandner, ed., *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang, 123-131.
- Kneisel, J., Dal Corso, M., Kirleis, W., Taylor, N., Tiedtke, V., 2015a. The third food revolution? Common trends in economic and subsistence strategies in Bronze Age Europe. In: J. Kneisel, M. Dal Corso, W. Kirleis, H. Scholz, N. Taylor, V. Tiedtke, eds., *The Third Food Revolution? Setting the Bronze Age Table: Common Trends in Economic and Subsistence Strategies in Bronze Age Europe*. Proceedings of the International Workshop 'Socio-Environmental Dynamics over the Last 12,000 Years: The Creation of Landscapes III' (15th-18th April 2013) in Kiel. Universitätsforschungen zur prähistorischen Archäologie 283. Bonn: Dr. Rudolf Habelt GmbH, 275-286.
- Kneisel, J., Dal Corso, M., Kirleis, W., Scholz, H., Taylor, N., Tiedtke, V., eds., 2015b. *The Third Food Revolution? Setting the Bronze Age Table: Common Trends in Economic and Subsistence Strategies in Bronze Age Europe*. Proceedings of the International Workshop 'Socio-Environmental Dynamics over the Last 12,000 Years: The Creation of Landscapes III' (15th-18th April 2013) in Kiel. Universitätsforschungen zur prähistorischen Archäologie 283. Bonn: Dr. Rudolf Habelt GmbH.
- Koenig, M.P. and Klag, T., 2017. Habitats, contextes funéraires et occupation du sol au Bronze final en Lorraine. In: L. Carozza, C. Marcigny, M. Talon, eds., *L'habitat et l'occupation des sols à l'âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS, 241-259.
- Lachenal, T., Mordant, C., Nicolas, T., Véber, C., eds., 2017. *Le Bronze moyen et l'origine du Bronze final en Europe occidentale (XVIIe-XIIIe siècle av. J.-C.)*. Colloque international de l'APRAB, Strasbourg, 17 au 20 juin 2014. Monographies d'Archéologie du Grand Est 1. Strasbourg: Association pour la Valorisation de l'Archéologie du Grand Est.

- Leroyer, C., David, R., Mazier, F., Allenet, G., Lanos, P., Dufresne, P., 2012. Environnement et anthropisation du milieu durant l'âge du Bronze dans le Bassin parisien: l'apport des données polliniques et de la modélisation du couvert végétal. In: M. Mélin and C. Mougne, eds., *L'Homme, ses ressources et son environnement, dans le nord-ouest de la France à l'âge du Bronze: actualités de la recherche*. Actes du Séminaire Archéologique de l'Ouest du 22 mars 2012. Mémoires de Géosciences hors-série 8. Rennes: Géosciences, 7-26.
- Lightfoot, E., Liu, X., Jones, M.K., 2013. Why move starchy cereals? A review of the isotopic evidence for prehistoric millet consumption across Eurasia. *World Archaeology*, 45 (4), 574-623. <https://doi.org/10.1080/00438243.2013.852070>
- Liu, X., Motuzaitė Matuzevičiūtė, G., Hunt, H.V., 2018. From a fertile idea to a fertile arc: the origins of broomcorn millet 15 years on. In: X. Liu, E. Lightfoot, H. Hunt, eds., *Far from the Hearth: Essays in Honour of Martin K. Jones*. McDonald Institute Conversations. Cambridge: McDonald Institute for Archaeological Research, 155-164. <https://doi.org/10.17863/CAM.38312>
- Lundström-Baudais, K., 2010. *Ciurā* et *muri*: le traitement en formes de flocons et de gruaux de quatre céréales dans le nord-ouest du Népal. In: H. Franconie, M. Chastanet, F. Sigaut, eds., *Couscous, boulgour et polenta: transformer et consommer les céréales dans le monde*. Paris: Karthala, 261-315.
- Lundström-Baudais, K., Rachoud-Schneider, A.-M., Baudais, D., Poissonnier, B., 2002. Le broyage dans la chaîne de transformation du millet (*Panicum miliaceum*): outils, gestes et écofacts. In: H. Procopiou and R. Treuil, eds., *Moudre et Broyer, I: Méthodes*. Paris: CTHS, 181-209.
- Magny, M., Bossuet, G., Gauthier, É., Richard, H., Vannière, B., Billaud, Y., Marguet, A., Mouthon, J., 2007. Variations du climat pendant l'Âge du Bronze au centre-ouest de l'Europe: vers l'établissement d'une chronologie à haute résolution. In: H. Richard, M. Magny, C. Mordant, eds., *Environnements et cultures à l'âge du Bronze en Europe occidentale*. Paris: CTHS, 13-28.
- Marinval, P., 1992. Archaeobotanical data on millets (*Panicum miliaceum* and *Setaria italica*) in France. *Review of Palaeobotany and Palynology*, 73, 259-270.
- Marinval, P., 1995. Données carpologiques françaises sur les millets (*Panicum miliaceum* L. et *Setaria italica* (L.) Beauv.) de la protohistoire au Moyen Âge. In: E. Hörterndner, ed., *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang, 31-61.
- Marinval, P. and Bonnamour, L., 2010. Bouillies de millet gallo-romaines découvertes dans la moyenne vallée de la Saône (France). In: H. Franconie, M. Chastanet, F. Sigaut, eds., *Couscous, boulgour et polenta: transformer et consommer les céréales dans le monde*. Paris: Karthala, 233-244.
- Michler, M., Schneikert, F., Véber C., 2017. L'Alsace et l'âge du Bronze: bilan et perspectives. In: L. Carozza, C. Marcigny, M. Talon, eds., *L'habitat et l'occupation des sols à l'âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS, 261-273.
- Miller, N.F., Spengler, R.N., Frachetti, M., 2016. Millet cultivation across Eurasia: origins, spread and the influence of seasonal climate. *The Holocene*, 26 (1), 1566-1575. <https://doi.org/10.1177%2F0959683616641742>
- Mordant, C., 2013. The Bronze Age in France. In: H. Fokkens and A. Harding, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: University Press, 571-593.
- Moreno-Larrazabal, A., Teira-Brión, A., Sopelana-Salcedo, I., Arranz-Otaegu, A., Zapata, L., 2015. Ethnobotany of millet cultivation in the north of the Iberian Peninsula. *Vegetation History and Archaeobotany*, 24, 541-554. <https://doi.org/10.1007/s00334-015-0518-y>

- Motuzaitė-Matuzevičiūtė, G., Staff, R.A., Hunt, H.V., Liu, X., Jones, M.K., 2013. The early chronology of broomcorn millet (*Panicum miliaceum*) in Europe. *Antiquity*, 87 (338), 1073-1085. <https://doi.org/10.1017/S0003598X00049875>
- Murty, D.S. and Kumar, K.A., 1995. Traditional uses of sorghum and millets. In: D.A.V. Dendy, ed., *Sorghum and Millets: Chemistry and Technology*. St. Paul, Minnesota: American Association of Cereal Chemists, 185-221.
- Peake, R., ed., 2020. *Villiers-sur-Seine, un habitat aristocratique du IXe siècle avant notre ère*. Recherches archéologiques 18. Paris: Inrap/CNRS.
- Peake, R., Brunet, P., Delattre, V., Néré, É., Boulenger, L., Gouge, P., 2017a. Bilan des connaissances de l'habitat et l'occupation du sol à l'âge du Bronze et du premier âge du Fer en Île-de-France. In: L. Carozza, C. Marcigny, M. Talon, eds., *L'habitat et l'occupation des sols à l'âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS, 187-212.
- Peake, R., Roscio, M., Issenmann, R., Riquier, V., Simonin, D., 2017b. Le Bronze moyen et le début du Bronze final du sud du Bassin parisien. In: T. Lachenal, C. Mordant, T. Nicolas, C. Véber, eds., *Le Bronze moyen et l'origine du Bronze final en Europe occidentale (XVIIe-XIIIe siècle av. J.-C.)*. Colloque international de l'APRAB, Strasbourg, 17 au 20 juin 2014. Monographies d'Archéologie du Grand Est 1. Strasbourg: Association pour la Valorisation de l'Archéologie du Grand Est, 195-216.
- Pêche-Quilichini, K., Paolini-Saez, H., Martin, L., Drieu, L., Arobba, D., 2020. Des balanophages en Corse entre Bronze moyen et premier âge du Fer? Une protohistoire du gland en contexte insulaire. In: G. Auxiette, C. Mougne, R. Peake, F. Toulemonde, eds., *Autour de la table: l'alimentation à l'âge du Bronze et au premier âge du Fer*. Actes de la Journée thématique du 3 mars 2017 à Saint-Germain-en-Laye. *Bulletin de l'Association pour la Promotion des Recherches sur l'Âge du Bronze*, suppl. 6, 77-91.
- Reed, K., 2020. Farmers in transition: crop choice during the Bronze and Early Iron Age in Croatia (2400-100 cal BC). In: G. Auxiette, C. Mougne, R. Peake, F. Toulemonde, eds., *Autour de la table: l'alimentation à l'âge du Bronze et au premier âge du Fer*. Actes de la Journée thématique du 3 mars 2017 à Saint-Germain-en-Laye. *Bulletin de l'Association pour la Promotion des Recherches sur l'Âge du Bronze*, suppl. 6, 67-76.
- Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, Ch., van der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S.M., Fogtman-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S., 2020. The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0-55 cal kBP). *Radiocarbon*, 62, 725-757. <https://doi.org/10.1017/RDC.2020.41>
- Ren, X., Lemoine, X., Mo, D., Kidder, T.R., Guo, Y., Qin, Z., Liu, X., 2016. Foothills and intermountain basins: does China's fertile arc have 'hilly flanks'? *Quaternary International*, 426, 86-96. <https://doi.org/10.1016/j.quaint.2016.04.001>
- Riquier, V., Auxiette, G., Fechner, K., Grisard, J., Loicq, S., Nicolas, T., Sehier, E., Toulemonde, F., Zipper, K., 2012. Oscillations et évolutions de l'habitat et des systèmes agraires en Champagne méridionale (2200-450 av. J.-C.): le terroir du 'Parc logistique de l'Aube'. In: M. Honegger and C. Mordant, eds., *L'Homme au bord de l'eau: Archéologie des zones littorales du Néolithique à la Protohistoire*. Actes du 135^e Congrès national des sociétés historiques et scientifiques du CTHS 'Paysages', Neuchâtel, 6-11 avril 2010, Session de pré- et protohistoire. Cahiers d'archéologie romande 132 (Documents préhistoriques 30). Lausanne and Paris: CTHS, 65-87.

- Riquier, V., Auxiette, G., Fechner, K., Loicq, S., Toulemonde, F., 2015. Éléments de géographie humaine et économique à l'âge du Bronze et à l'âge du Fer dans la Plaine de Troyes. *Bulletin de la Société Préhistorique Française*, 112 (2), 339-367.
- Riquier, V., Grisard, J., Le Goff, I., 2017a. L'âge du Bronze et le premier âge du Fer en Champagne-Ardenne: l'occupation du sol vue sous l'angle de l'archéologie préventive. In: L. Carozza, C. Marcigny, M. Talon, eds., *L'habitat et l'occupation des sols à l'âge du Bronze et au début du premier âge du Fer*. Recherches Archéologiques 12. Paris: Inrap/CNRS, 213-326.
- Riquier, V., Le Goff, I., Nicolas, T., 2017b. Le Bronze moyen (et l'origine du Bronze final) en Champagne à la lumière de l'archéologie préventive. In: T. Lachenal, C. Mordant, T. Nicolas, C. Véber, eds., *Le Bronze moyen et l'origine du Bronze final en Europe occidentale (XVIIe-XIIIe siècle av. J.-C.)*. Colloque international de l'APRAB, Strasbourg, 17 au 20 juin 2014. Monographies d'Archéologie du Grand Est 1. Strasbourg: Association pour la Valorisation de l'Archéologie du Grand Est, 135-155.
- Rösch, M., Fisher, E., Lechterbeck, J., Kleinmann, A., Sillmann, M., Tserendorj, G., Wick, L., 2015. Bronze Age land use and food production in southwest Germany in light of botanical on-site and off-site data. In: J. Kneisel, M. Dal Corso, W. Kirleis, H. Scholz, N. Taylor, V. Tiedtke, eds., *The Third Food Revolution? Setting the Bronze Age Table: Common Trends in Economic and Subsistence Strategies in Bronze Age Europe*. Proceedings of the International Workshop 'Socio-Environmental Dynamics over the Last 12,000 Years: The Creation of Landscapes III' (15th-18th April 2013) in Kiel. Universitätsforschungen zur prähistorischen Archäologie 283. Bonn: Dr. Rudolf Habelt GmbH, 201-219.
- Rosenstock, E., Groß, M., Hujić, A., Scheibner, A., 2015. Back to good shape: biological standard of living in the Copper and Bronze Ages and the possible role of food. In: J. Kneisel, M. Dal Corso, W. Kirleis, H. Scholz, N. Taylor, V. Tiedtke, eds., *The Third Food Revolution? Setting the Bronze Age Table: Common Trends in Economic and Subsistence Strategies in Bronze Age Europe*. Proceedings of the International Workshop 'Socio-Environmental Dynamics over the Last 12,000 Years: The Creation of Landscapes III' (15th-18th April 2013) in Kiel. Universitätsforschungen zur prähistorischen Archäologie 283. Bonn: Dr. Rudolf Habelt GmbH, 121-151.
- Rozier, F., 1821-23. Caméline. In: F. Rozier. *Dictionnaire raisonné et universel d'agriculture*. Vol. 3. Paris: Deterville, 342-344.
- Saurel, M., Fronteau, G., Garnier, N., Roseau, B., Verdin, P., Auxiette, G., Toulemonde, F., 2021. Du contenu et de l'apparence: l'analyse des céramiques de la tombe à char de Warcq 'La Sauce' dans les Ardennes (II^e siècle av. J.-C.). In: D. Frère et al., eds., *Manger, boire, se parfumer pour l'éternité : rituels alimentaires et odorants en Italie et en Gaule du IX^e siècle avant au I^{er} siècle après J.-C.* Naples, Centre Jean Bérard, 53, 393-332.
- Sherrat, A., 2006. The trans-Eurasian exchange: the prehistory of Chinese relations with the West. In: V. Mair, ed., *Contact and Exchange in the Ancient World*. Honolulu: Hawaii University, 30-61.
- Sigaut, F., 1995. Les millets en Eurasie, d'une fête populaire à des questions pour les chercheurs. In: E. Hörandner, ed., *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang, 133-153.
- Stika, H.P. and Heiss, A.G., 2013. Plant cultivation in the Bronze Age. In: H. Fokkens and A. Harding, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 340-361.
- Toulemonde, F., 2013. *Économie végétale et pratiques agricoles au Bronze final et au premier âge du Fer, de la côte de l'Île-de-France à la côte de Champagne*. Doctoral dissertation, Université de Paris 1.
- Toulemonde, F., 2014. Les blocs carbonisés de millet commun de l'habitat protohistorique de Villiers-sur-Seine 'Le Gros Buisson' (Seine-et-Marne): peut-on parler

- de préparation alimentaire? *ArchéoSciences, Revue d'Archéométrie*, 38, 109-118. <https://doi.org/10.4000/archeosciences.4192>
- Toulemonde, F., 2020. La consommation végétale, du banal au singulier. In: R. Peake, ed., *Villiers-sur-Seine, un habitat aristocratique du IXe siècle avant notre ère*. Recherches archéologiques 18. Paris: Inrap/CNRS, 183-212.
- Toulemonde, F., Zech-Matterne, V., Wiethold, J., Brun, C., Malrain, F., Riquier, V., 2017. Reconstitution des pratiques agricoles du Ier millénaire a.C. en France orientale, d'après le croisement des données carpologiques et archéologiques. In: G. Bataille, J. Kaurin, S. Deffressigne, S. Marion S., eds., *Production et Proto-industrialisation aux Âges du Fer: Perspectives sociales et environnementales*. Actes du 39^e colloque international de l'AFEAF, Nancy 14-17 Mai 2015. Mémoires 47. Bordeaux: Ausonius, 29-50.
- Toulemonde, F., Auxiette, G., Bouby, L., Goude, G., Peake, R., with the collaboration of V. Forrest, 2018. L'alimentation à l'âge du Bronze en France. In: J. Guilaine and D. Garcia, eds., *La Protohistoire en France*. Paris: Hermann, 297-309.
- Touzeau, R., 1995. La fête du millet à Aizenay. In: E. Hörandner, ed., *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang, 19-29.
- Valamoti, S.M., 2016. Millet, the late comer: on the tracks of *Panicum miliaceum* in prehistoric Greece. *Archaeological and Anthropological Sciences*, 8, 51-63. <https://doi.org/10.1007/s12520-013-0152-5>
- Vencl, S., 1985. Zaludy jako potravina: K poznání významu sběru pro výživu v pravěku [Acorns as food: assessing the significance of food-gathering for prehistoric dietary habits]. *Archeologické rozhledy*, 37, 516-565.
- Vencl, S., 1996. Acorns as food: again. *Památky archeologické*, 87, 95-111.
- Verger, S. and Guillaumet, J.P., 1988. Les tumulus de Saint-Romain-de-Jalionas (Isère). In: J.P. Mohen, A. Duval, C. Eluère, eds., *Les Princes Celtes et la Méditerranée*. Rencontres de l'École du Louvre, 25-27 novembre 1987, Paris. Paris: La Documentation Française, 231-240.
- Wiethold, J., 2007. Restes végétaux carbonisés provenant des sites de l'Âge du Bronze (St. Apollinaire, 'Le pré de Crot 1'; Couternon, 'Larrey', Varois-et-Chaignot, 'Le pré du Plancher') [Rapport carpologique 2005]. In: R. Labeaune, F. Ducreux, P. Barral, J. Wiethold, C. Bémilli, eds., *Liaison Routière Dijon / Arc-sur-Tille: les occupations protohistoriques: Vol. 5 Synthèses*. Dijon: AFAN, Inrap, SRA de Bourgogne, Conseil Général de la Côte d'Or, 58-63.
- Zech-Matterne, V. and Brun, C., 2016. Vers une agriculture extensive? Étude diachronique des productions végétales et des flores adventices associées, au cours de la période laténienne, en France septentrionale. In: F. Malrain and G. Blancquaert, eds., *Évolution des sociétés gauloises du Second âge du Fer, entre mutations internes et influences externes*. Actes du 38^e colloque international de l'AFEAF, Amiens, 29 mai-1 juin 2014. *Revue Archéologique de Picardie*, suppl. 30, 623-638.
- Zech-Matterne, V., Bonnaire, E., Daoulas, G., Derreumaux, M., Durand, F., Rousselet O., Schaal, C., Toulemonde, F., Wiethold, J., 2017. Diversité et évolution des productions céréalières et fruitières dans le quart nord-est de la France d'après les données carpologiques (II^e s. av. J.-C.-V^e s. ap. J.-C.). In: S. Lepetz and V. Zech-Matterne, eds., *Productions agro-pastorales, pratiques culturelles et élevage dans le nord de la Gaule du deuxième siècle avant J.-C. à la fin de la période romaine*. Actes de la Table Ronde Internationale des 8 et 9 mars 2016 à l'Institut National d'Histoire de l'Art (Paris). *Archéologie des Plantes et des Animaux* 5. Quint-Fonsegrives: Mergoïl, 43-62.
- Zohary, D., Hopf, M., Weiss, E., 2012. *Domestication of Plants in the Old World: the Origin and Spread of Domesticated Plants in Southwest Asia, Europe, and the Mediterranean Basin*. (4th edition). Oxford: Oxford University Press.

On-site to off-site: A multidisciplinary and multiscale consideration of the 13th to 11th century BCE transformation in northern Germany

*Ingo Feeser, Stefanie Schaefer-Di Maida, Stefan Dreibrodt,
Jutta Kneisel, Dragana Filipović*

Abstract

Older and more recent excavations in the Mang de Barga micro-region of northern Germany have revealed traces of local human activities, starting from the Late Neolithic (ca. 2200 BCE) and continuing, through the Bronze Age and Early Iron Age, to the Roman Iron Age (1st century BCE). This paper focuses on the Late Neolithic and Bronze Age occupation, when the site was used mainly for funerary activities. Near- and on-site palaeo-environmental investigations revealed that the local environment was changing during this period. The fundamental shift in burial rites during the 13th to 11th centuries BCE coincided with a fundamental change in land-use practices. This probably included an intensification of use of open, agricultural land associated with a shift from woodland to grassland pasturage. A number of crops typical of Late Neolithic and Bronze Age agriculture in northern Germany were in use at Mang de Barga. A comparison of the developments at the local scale (based on both on- and near-site data) with the archaeological, archaeobotanical and palaeo-environmental evidence from a regional (Schleswig-Holstein, Germany) and a supra-regional (northern Germany) scale (using off-site data) reveals similar and coinciding changes in material culture, burial rites and land-use patterns. The changing land-use practices in northern Germany during the last centuries of the 2nd millennium BCE are also reflected in the diversification of the crop spectrum, including the beginning of cultivation of broomcorn millet (*Panicum miliaceum*). This supports the idea that the 13th to 11th centuries BCE were the time of far-reaching societal transformation. Substantial changes occurred in many forms of material expression (settlement architecture, layout and size; utilitarian objects), as well as in ideology (mortuary practice), technology (raw materials and techniques), trade and exchange (intensified circulation of goods), and the food economy (new crops and domesticates).

The 13th to 11th century BCE transformation in northern Germany

The period covering the 13th to 11th centuries BCE was an exceptionally dynamic one in the cultural history of northern Germany. On the basis of the recently produced absolute dates (Schaefer-Di Maida 2020), it is considered as the transition from the Older (Period Ib-III, 1700-1100 BCE) to the Younger Bronze Age (Period IV-VI, 1100-530/500 BCE), which starts in the second half of Period III (around 1200 BCE) in the regional archaeological periodisation (cf. Fig. 4). The transition brought a number of changes and innovations in the location and size of settlements, treatment of the dead, access to and choice of raw materials, tool and weapon type preference, and the size of the population (Feesser *et al.* 2019; Kneisel *et al.* 2019). The changes were substantial and far-reaching; the period is, therefore, regarded as a time of societal transformation evident in multiple aspects of life.

The archaeological evidence indicating these large-scale developments at around 1200 BCE can be summarised as a series of up- or downward trends in different archaeological datasets reflecting some major areas of human activity: social, technological, ideological and economic. Through systematisation and chronological alignment of these numerous on-site and off-site datasets from the Bronze Age in northern Germany, Kneisel *et al.* (2019) were able to identify several important general trends from about 1200 BCE onwards, relative to the previous periods. These are:

- a decrease in house size;
- an increase in the number of houses, settlements and settlement pits;
- the emergence of so-called storage-pit fields;
- already from 1500 BCE, increased human pressure on the landscape resulting from population growth;
- lower amounts of amber; and
- a sharp drop in the number of metal axes and sickles and, somewhat later, of swords and daggers.

In the 13th or 12th century BCE, a new cereal crop arrived in the region, probably from the south or southeast (Filipović *et al.* 2020a) – broomcorn millet (*Panicum miliaceum*). The earliest occurrences include large deposits, such as those at Rullstorf (site No. 5, county of Lüneburg, Lower Saxony; Kirleis 2003) and Badegow (site No. 19, county of Ludwigslust-Parchim, Mecklenburg-Vorpommern; Saalow *et al.* 2014), as well as many contemporaneous cases where smaller amounts of grains were discovered (Filipović *et al.* 2020a, Supp. Dataset). Radiocarbon dates produced directly on broomcorn millet grains suggests that this innovation spread very quickly across northern Germany. The start of millet cultivation was, potentially, a major agricultural change at the transition from the Older to the Younger Bronze Age. In northern Germany, this period is also marked by the beginning of the use, or increase in the use, of some other crops, such as pulses and oil/fibre plants.

Approaching the research questions

At least some of these synchronous and spatially overlapping trends were intertwined, or one caused the other. For instance, the intensified trade and exchange network, usually understood as distributing metal, metal objects and cultural influences, probably also served as a mechanism for the diffusion of crops. The new cultivar (broomcorn millet) may have led to changes in the ways the land was used for agricultural and other purposes. There may have been new tools and techniques employed in food production.




Figure 1. Location of Mang de Bergen and additional sites and geographic locations mentioned in the text. A: Map of Schleswig-Holstein showing location of the site of Mang de Bergen and other archaeobotanically analysed sites in the wider Mang de Bergen region mentioned in the text, and indicating the area shown in Fig. 1B (red rectangle). B: Map of the area around the site of Mang de Bergen, showing burial mounds (red dots) and clusters of burials (black lines); the cluster Mang de Bergen is shown in detail in Fig. 1C. C: Map showing the distribution of burial locations at Mang de Bergen (the numbers refer to the LA-numbers mentioned in the text).

Despite the good archaeological knowledge of this period, some key aspects remain elusive. For instance, how was the increasing human pressure on the environment exerted? Was a greater area needed to accommodate cultivation of the new crop? Did this increase the value of the land suitable for cropping and, possibly, create tensions in society? The demand for wood fuel presumably increased due to the new burial ritual; would this have been reflected in the landscape?

These and similar questions are important, as their answers can clarify the interconnections between changes in the environment and society and thus elucidate their knock-on effect. They cannot be addressed using a single line of evidence; they are also unlikely to be clarified to the same level of detail at the local to regional to supra-regional scales. Therefore, we take a multidisciplinary approach and look at the datasets produced by archaeological, palaeo-environmental and archaeobotanical analyses. We consider them separately, and then we combine them at different scales.

The local scale is here represented by the site of Mang de Barga, in Schleswig-Holstein, Germany (Fig. 1). The results of the archaeological, palaeo-environmental and archaeobotanical analyses that have recently been carried out there have now been published (Feaser *et al.* 2020; Filipović *et al.* 2020b; Schaefer-Di Maida 2020; Kneisel *et al.* in prep.), meaning that we can use the final data and that the archaeological context and dating of the different materials have been fully reconstructed. We take the area of Schleswig-Holstein, and in particular the eastern younger morainic region, for which generally similar and synchronous cultural developments can be anticipated during the Bronze Age (cf. Schaefer-Di Maida 2020), as representing the regional scale of investigations. It includes Lake Belau, from which one of the palaeo-environmental archives discussed here derives (see below), as well as a number of other archaeological sites that are of similar age as Mang de Barga (Fig. 1). Northern Germany is our supra-regional scale, and it serves as the wider geographical context for our observations made at lower scales. In parallel, we use some observations valid for all of northern Germany to inform our understanding of the developments at the regional and local levels.

This multidisciplinary and multi-scalar approach has three advantages. First, it allows us to compensate for the low availability or resolution of some of the data (*e.g.* archaeobotanical), which is an inevitable consequence of the nature of the studied archives and their differential rate of preservation. As will be shown below, not all of the selected evidential strands offer the same amount of detail, be it qualitative or quantitative, within and between the spatial scales. Second, it enables us to articulate the observations made from the different forms of data and to devise hypotheses on the cause-and-effect relationships between the changes seen in different aspects of the environment and human behaviour. Third, we can move between the scales, that is, use the site-based information to inform the supra-regional picture, and *vice versa*.

Mang de Barga: Location and history of research

The site of Mang de Barga (also called Mank den Barga) is located about 2 km east of the town of Bornhöved, in the county of Segeberg, in the state of Schleswig-Holstein. It is situated between the Grimmelsberg end-morainic ridge (ca. 83 m asl) in the southeast and the Bornhöved lake chain in the northwest (Fig. 1B). The northwestern slope of the Grimmelsberg is composed of several morainic tops dating from the last Ice Age. The end-morainic landscape is covered by a variety of soil types, such as luvisols, cambisols and brown soil, underlain by a glacier-accumulated substrate made up of sand, gravel and clay lenses. The base of slopes is covered in colluvium. Along the lakes and ponds, one finds gleys and peat (Piotrowski 1991; Dreibrodt *et al.* 2009).

A number of burial mounds were constructed on the slopes of the Grimmelsberg ridge and along its foot during the Late Neolithic and the Bronze Age (Fig. 1B), most of them grouped in clusters of 3–30. Mang de Barga itself, which represents a cluster of about 20 burial mounds (with 76 burials), has the highest concentration of these features. In the pre-Roman Iron Age and succeeding periods, an Urnfield cemetery (LA 115) located adjacent to the barrow area developed, and it preserved remains of 201 cremations. A little farther to the northwest, ca. 750 m away from the burial

zone, are a settlement from the Iron Age (LA 97) and a small Urnfield cemetery from the Younger Bronze Age (LA 78), as yet unexcavated but known from surveys.

The Mang de Barga mounds (Fig. 1C) were recorded as early as the 1960s, within a regional archaeological survey (Schwerin von Krosigk 1976). In 2004-2005 and 2014, the Archaeological Heritage Department of Schleswig-Holstein carried out rescue excavations prior to the expansion of a local gravel pit. With the exception of one mound (LA 57), all burial mounds visible on the surface, as well as the pre-Roman Iron Age urnfield, were investigated. Some of the burials were heavily disturbed, and only traces of these were discovered. Others were almost entirely destroyed and offered no evidence of former barrows or graves.

Parallel to the archaeological fieldwork in 2005, the Ecology Centre of CAU Kiel conducted investigations of the soil cover around the burial mounds in order to determine the degree of vegetation openness based on colluvial formations (Dreibrodt *et al.* 2009, 481-491). Previous work on the reconstruction of the vegetation and occupation history of the Bornhöved lake chain had taken place in the 1980s and 1990s within two research projects (the Bundesministerium für Bildung und Forschung-funded project 'Ökosystemforschung Bornhöveder Seenkette' and the Deutsche Forschungsgemeinschaft-funded project 'Neolithisierung in Schleswig-Holstein'). As part of these programmes, a high-resolution pollen diagram was produced from Lake Belau, offering insight into the regional human impact on the landscape (Wiethold 1998; Dörfler *et al.* 2012; Dreibrodt and Wiethold 2015).

Mound LA 57 was excavated in 2017, by the Collaborative Research Centre (CRC) 1266 (subproject D3), Kiel University (Schaefer-Di Maida 2020; Kneisel *et al.* in prep.). The magnetic prospection of the cemetery was performed by the Institute of Geosciences in Kiel. Additional magnetic prospections and test excavations were carried out in the wider surroundings of the cemetery (Kneisel *et al.* 2017). Also in the context of the CRC 1266, the history of the Mang de Barga palaeo-environment was reconstructed, based on the analysis of sediment samples taken on-site – from the archaeological contexts – and the study of a short peat sequence from the area 300 m north of the site (subproject F2). The latter represents a so-called near-site archive, with local and extra-local pollen as the dominant components (*sensu* Moore *et al.* 1991). The analysis of the macro-botanical remains was also part of the research done within the CRC 1266 (subproject F3).

Prehistoric burial activity at Mang de Barga

The relative chronological attribution of the artefacts and the absolute dating of cremated bones and charcoal show that the burial activity at Mang de Barga started in the Late Neolithic, at ca. 2200 BCE, and lasted until the mid-1st century BCE. From the Late Neolithic until the start of Bronze Age Period III (ca. 1300 BCE), the burial activity involved the formation of grave mounds (barrows), under and within which the deceased were inhumated. Each mound covered a single primary burial in its centre, but some also enclosed secondary burials placed in the upper layers or in the periphery of the mound. Only a small, selected ('privileged') part of the community was buried in the barrows (*e.g.* 10% according to Kristiansen 2018, 110). The rest of the community may have received flat graves; these are, however, difficult to trace archaeologically. In the central grave, the deceased was laid in an oak-log coffin placed upon a stone-built base or surrounded by a row of stones. This construction was then covered with one or more heaps or layers of earth, creating the core and the mantle of the mound.

Only a few burials investigated at Mang de Barga derived from the Late Neolithic phase of its use (Fig. 2), such as those at LA 63 and LA 64 (Schaefer-Di Maida 2020). Almost no burials were recorded for the period of transition from the Neolithic to the Bronze Age, that is, Bronze Age Period I (1750-1500 BCE). Either the degree of use




Figure 2. Chronological sequence and nature of the cultural developments at Mang de Bergen, reconstructed from the archaeological evidence.

of the site was much lower during this phase, or there was a gap in its occupation altogether. Bronze Age Period II (1500-1300 BCE) witnessed a considerable rise in the burial activity at Mang de Bergen and was the main phase of barrow construction.

The final barrow was built around 1300 BCE, marking the end of this funerary practice. This last barrow also contained the earliest cremation burial at the site, thus clearly documenting the shift from inhumation to cremation, which happened all over Schleswig-Holstein at about this time. Subsequently, from the middle of Bronze Age Period III (around 1200 BCE) onwards, the pre-existing barrows and their immediate surroundings at Mang de Bergen were re-used for secondary burials. These were exclusively cremations.

The burial activity that incorporated the new ritual was relatively low up until 1200 BCE. From this point on, it increased rapidly, and the number of burials peaked at 950/900 BCE (Fig. 2). Afterwards, the number of graves decreased, and it remained very low during Bronze Age Period VI (700-530/500 BCE). Another surge in burials began shortly before the transition to the pre-Roman Iron Age (530/500 BCE). This was mostly due to the start of the use of the urn cemetery (LA 115) located next to the grave mound area of the site, from ca. 530/500-480 BCE.

The cultural transformation of the 13th to 11th centuries BCE at Mang de Bergen

The archaeological evidence indicates a gradual change in burial rites from 1300 BCE onwards. Whereas at around 1300 BCE the treatment of the dead changed, *i.e.* from inhumation to cremation, the method of grave construction, *i.e.* the erection of grave mounds, continued unchanged. The repertoire of grave goods did not change either;

it continued to consist of tools, weapons and pieces of jewellery. There were also some new, non-funerary elements introduced at this time, though, and these are the so-called cooking pits. They were found in the immediate vicinity of Mang de Barga, arranged in circles and dated to approximately 1300 BCE (Schaefer-Di Maida 2020).

From 1200 BCE (start of the Younger Bronze Age), the grave construction at Mang de Barga changed fundamentally. The deceased were now exclusively cremated. Their remains were buried in urns placed within stone circles or were laid directly on a bed of stone. Burial took place within the pre-existing mounds or along their margins. More people were buried here in the last centuries of the 2nd and early centuries of the 1st millennium BCE than in any of the previous phases of use of the site (Fig. 2). For example, up to 35 secondary burials have been detected around a single mound (LA 57). From a practical point of view, the more intensive use of the site could have simply been because urn or directly buried cremation graves required less space than a single-grave barrow, and therefore more graves could be placed in the area. Also, the amount of resources needed, such as construction materials (wood, soil) and human labour (for the mound building, but also for the transport of earth and vegetation sods; cf. Schunke 2018), may have been higher for the creation of barrows than for the execution of cremation funerals. On the other hand, considerable resources and labour were also required for the cremation burials, *i.e.* for the collection of wood, construction of the funeral pyre, and transport of the stones used for lining/encircling the graves. Falkenstein suggests that the amount of labour invested in cremation burials would have been similar to that invested in building tumuli (Falkenstein 2012, 336). He goes on to suggest that the time required to collect (enough) wood for the pyre would have been comparable to the time needed to raise a mound 7 m in diameter (Falkenstein 2017, 80-81). The diameter of the burial mounds at Mang de Barga generally exceeds 20 m, so constructing these mounds would, following these calculations, have taken much more resources and energy expenditure than cremation funerals.

The 12th century BCE transformation in the burial ritual at Mang de Barga entailed a new repertoire of grave goods. Tools, weapons and jewellery were replaced by dress ornaments and pottery. Overall, grave good inventories appear less diverse, almost standardised. This, together with the simple, uniform burials perhaps suggests that the members of the community now received more equal funerary treatment. Indeed, the period of burial mound construction has been described as the time of chiefs, warrior elites and pronounced social stratification (Jensen 1982; Kristiansen 1987, 1991; Earle 1991, 2002; Vandkilde 1996, 2006). The dramatic change in the burial ritual and other aspects of life in the Younger Bronze Age may point to a major societal change, a transformation of the social structure, apparently from non-egalitarian to (more) egalitarian.

Plant use at Mang de Barga: The archaeobotanical perspective

In order to retrieve evidence of prehistoric plant use at Mang de Barga, archaeobotanical sampling was employed at 11 burial locations, and at one location (LA 116) that comprised remains of different Younger Bronze Age features (pits, ovens, hearths) reflecting activities whose character – whether residential or funerary – could not be clearly distinguished. An assortment of archaeological contexts of interest to the researchers were sampled at these locations: graves, pits, cooking pits, hearths and an oven. The number of samples taken at each of the locations depended on the presence and number of such clearly defined contexts. The samples, measuring, on average, 8-9 litre in volume, were floated manually using buckets and a 0.3 mm geological sieve. All plant remains encountered are in

			Site	LA 63	LA 64	LA 23	LA 69	LA 115	LA 57	LA 116	
			Type of site	Burial	Burial	Burial	Burial	Burial	Burial	Settlement	
			Period	Late Neolithic		BA Period I-II			BA Period II-III	BA Period V-VI	
			Number of analysed features	2	1	2	2	1	21	8	
			Volume of analysed samples (l)	16	18	14	25	6	348	38	
TAXON	Weight of wood charcoal (g)		2.5	1.3	20.2	4.2	75.7	73.7	25.1		
Crops	plant part	common name									
<i>Hordeum vulgare</i>	grain	barley	3	1					8	8	
<i>Triticum</i> cf. <i>monococcum</i>	grain	einkorn							1		
<i>Triticum monococcum/dicoccum</i>	grain	einkorn/emmer							1		
<i>Triticum dicoccum</i>	grain	emmer	1						9		
<i>Triticum dicoccum</i>	glume base	emmer							5		
<i>Triticum aestivum/durum/turgidum</i>	grain	free-threshing wheat		1					1		
<i>Triticum spelta</i>	glume base	wheat							1		
<i>Triticum</i> sp.	grain	wheat			2					4	
<i>Cerealia indeterminata</i>	grain	non-millet cereal	1	1					21	5	
cf. <i>Cerealia indeterminata</i>	grain	cereal								1	
<i>Panicum miliaceum</i>	grain	broomcorn millet								2	
Wild plants	plant part	common name									
<i>Alnus</i> sp.	catkin fragment	alder							1		
<i>Bromus</i> sp.	fruit	brome grass				1					
<i>Calluna vulgaris</i>	fragment of flower	heather								2	
<i>Carex</i> sp., <i>tricarpetate</i>	nutlet	sedge								3	
<i>Carpinus betulus</i>	seed	common hornbeam							1	1	
cf. <i>Mentha</i> sp.	seed	mint					1				1
<i>Chenopodiaceae</i>	seed	goosefoot family								2	
<i>Chenopodium album</i>	seed	fat hen								13	
<i>Chenopodium</i> sp.	seed	goosefoot								1	
<i>Coenococcum geophilum</i>	sclerotium	soil fungi								40	
<i>Corylus avellana</i>	nutshell fragment	hazelnut								3	
<i>Fabaceae</i>	seed	legumes								2	
<i>Hypericum</i> cf. <i>perforatum</i>	seed	St. John's wort								1	
<i>Persicaria lapathifolia</i> agg.	nutlet	pale persicaria									1
<i>Persicaria lapathifolia/maculosa</i>	nutlet				1		1			1	
<i>Persicaria maculosa</i>	nutlet	lady's thumb								1	2
<i>Phleum pratense</i> s.str.	fruit	timothy grass	1								

Table 1 (continued on opposite page). Results of the archaeobotanical analysis of different locations and contexts at Mang de Bergen.




Figure 3. Selected archaeobotanical remains from Mang de Bergen (photo: Dragana Filipović, Tanja Reiser): 1. Free-threshing wheat or spelt grain (LA 57); 2. Free-threshing wheat or emmer grain (LA 57); 3. Emmer grain (LA 57); 4. Barley grain (LA 57); 5. Spelt glume base (LA 57); 6. Emmer glume base (LA 57); 7. Broomcorn millet grain (LA 116); 8. Acorn cotyledon (LA 57); 9. *Alnus* sp. catkin (LA 57). 10. *Carpinus betulus* seed (LA 115); 11. Rhizomes, possibly *Arrhenatherum* (LA 57); 12. *Rubus* seed fragment (LA 57); 13. *Phleum* fruit (LA 63); 14. *Hypericum* cf. *perforatum* seed (LA 57); 15. *Viola* seed (LA 57); 16. *Brassicaceae* seed (LA 69); 17. *Chenopodium album* seed (LA 57); 18. *Fallopia convolvulus* seed and endosperm (LA 57); 19. *Plantago lanceolata* seed (LA 57); 20. *Vicia* seed (LA 57); 21. *Scirpus mucronatus* seed (LA 57); 22. *Carex* seed (LA 57).

Crop taxa	Late Neolithic	Bronze Age Period I-II	Bronze Age Period II-III	Bronze Age Period V-VI
Barley	X		X	X
Emmer	X		X	
Free-threshing wheat	X		X	
Spelt			X	
Einkorn			X	
Broomcorn millet				X

Table 2. Presence/absence of crop taxa in the different phases of occupation of Mang de Barga.

a charred state, and the majority of them represent fragments of wood charcoal. Detailed archaeobotanical data are available for LA 23, LA 57, LA 63, LA 64, LA 69, LA 115 and LA 116 and are given in Table 1 (Filipović *et al.* 2020b).

Despite the large number of samples processed, only a very small non-wood plant assemblage was recovered, composed of poorly preserved parts of crops and wild plants (Table 1; Fig. 3). The assemblage is, nonetheless, quite diverse. Crops are represented by grains of barley (*Hordeum vulgare*), grains and chaff of emmer (*Triticum dicoccum*), grains of free-threshing wheat (*T. aestivum/durum*) and einkorn (*T. monococcum*), grains and chaff of spelt (*T. spelta*) and grains of broomcorn millet. Finds of barley and emmer are the most frequent. The seeds identified as *Vicia* sp. may belong to potentially cultivated common vetch (*Vicia sativa*). The assemblage of wild plants from Mang de Barga is also diverse and consists of seeds and fragments of nuts, nutshell, culms, rhizomes and tubers. Some of the remains belong to edible parts of trees and bushes, such as oak (*Quercus*), common hazel (*Corylus avellana*) and blackberry (*Rubus idaeus*), and they may represent traces of food. Seeds of fat hen (*Chenopodium album*), knotgrass (*Polygonum aviculare*), pale persicaria (*P. lapathifolium*), sheep's sorrel (*Rumex acetosella*), black nightshade (*Solanum nigrum*) and others document the presence of arable weeds in the assemblage. The vegetative plant parts (stems, culm bases, stolons/rhizomes) may represent traces of plants used as kindling for cremation or domestic fires. Some of these remains perhaps come from false oat-grass (*Arrhenatherum elatius* ssp. *bulbosum*), which was documented at some other sites in this region as well as throughout northwestern and north-central Europe (e.g. Bornhöved LA 117 [Kneisel *et al.* in prep.] and Gönnebek 2013-215 [Effenberger 2018a]; see Roehrs *et al.* 2013).

Plant remains were found in both burial and non-burial contexts, suggesting that the plants were used and/or became charred in different locations across the site. Another possibility is that there was general mixing of the material from different contexts due to recurrent digging and redepositing of soil to cover the graves. Thus, for instance, crop-processing waste (cereal chaff, arable weed seeds) was found in both settlement and burial pits, but cleaning of the grain before use may have taken place elsewhere – perhaps away from both residential and funerary areas. Due to the long-term and apparently intensive use of the site through later prehistory, it was expected that plant materials dating from different periods of occupation would have become mixed together in the archaeological layers. Therefore, a selection of plant remains were submitted for radiocarbon dating (Schaefer-Di Maida 2020), allowing their approximate age to be established. Using a combination of absolute dates and contextual or stratigraphic information, the plant remains are here broadly attributed to individual phases of occupation/use of Mang de Barga. This provides an overview of the presence/absence of different crop taxa through time (Table 2).

The largest diversity of crop species was noted at LA 57, which is where the number of sampled contexts was the highest (22, compared with <10 at other locations; see Table 1). Interestingly, just three Late Neolithic contexts yielded traces

of three crop taxa, whereas the five Bronze Age Period I-II contexts did not contain any crop remains. Based on this, the impression is that there was no or little crop cultivation or consumption at or around Mang de Barga during Bronze Age Periods I and II. This is very much in line with the absence of human activity at the site during Bronze Age Period I, as inferred from the archaeological evidence (see above).

Some crop species were in use at Mang de Barga in both the Late Neolithic and the later phases of the Bronze Age, namely barley, emmer and free-threshing wheat. Although wheat remains were not confirmed for the final phases of the Bronze Age, they may be present in the category ‘*Cerealia indeterminata*’ (indeterminate cereal grain) (Table 1). Broomcorn millet finds from Mang de Barga derived from the Younger Bronze Age. Only two grains were found, both at LA 116, in a pit dated to the 8-6th century BCE. Broomcorn millet arrived in and spread across northern Germany in the 13th-12th centuries BCE (Effenberger 2018a; Filipović *et al.* 2020a) and may have been available to the Mang de Barga occupants already from this time onwards.

For the purpose of radiocarbon dating, a small collection of wood charcoal fragments, extracted from inhumations or cooking pits from locations LA 23, LA 57, LA 64 and LA 116, has been identified. A fragment of birch (*Betula*) wood from LA 64 was dated to the second half of the 3rd millennium BCE (the Late Neolithic). Several fragments of oak (*Quercus*) and hazel (*Corylus*) wood from LA 23 and LA 57 derived from the second half or the end of the 2nd millennium BCE (Schaefer-Di Maida 2020).

Trends in local land use through time: The palaeo-environmental evidence

On-site investigations

Pedological and sedimentological analyses on samples taken just east of barrow LA 57 revealed a fossil soil covered by three colluvial layers (Fig. 4). Based on the optically stimulated luminescence (OSL)-dates, the fossil soil horizon (fAh) dates to the Late Neolithic (Kneisel *et al.* in prep.), and the colluvial material has been accumulating on top of it from the Iron Age onwards. Palynological analysis has been carried out for the fAh horizon and the earliest colluvial layer (M1); the results are summarised in Table 3. The predominance of subfossil pollen grains of herbaceous land plants (72% excluding fern spores) along with the presence of cereal-type pollen, partly in clumps, indicates that the site was used for agriculture probably already during the Late Neolithic. The findings of fused birch pollen could point to the existence of fallow land and the beginning of regeneration of woody vegetation near LA 57. Despite the possibility of taphonomic enrichment of fern spores due to selective preservation (Haviga 1984), the comparatively high proportion of spores of spotted fern (*Polypodium vulgare*) and common viper’s tongue (*Ophioglossum*), both of which prefer a low-nutrient environment, could indicate nutrient-impoverished soil conditions (cf. Feeser and Dörfler 2016, 2019), which is not unlikely given the generally sandy soils in the area.




Figure 4. Pedological, sedimentological and palynological analyses of a soil profile from the archaeological layers at LA 57. Results of the analysis of pollen samples Pollen 1 and 2 are given in Table 3.

Site	LA 57	LA 57	Site	LA 57	LA 57
Context	N-Profil	N-Profil	<i>Ranunculus acris</i> -type	-	1
Sample	Pollen 1	Pollen 2	<i>Sinapis</i> -type	-	4
Labnr.	22152	22153	<i>Succisa</i>	-	2
Volume (ml)	7	8	Cultivars		
Weight (g)	6.05	10.11	<i>Avena</i> -type	-	1
Added markers	19223	19223	<i>Cerealia</i> -type indet.	5	21/4/-
Counted markers	12	597	<i>Hordeum</i> -type	-	4
Trees and shrub taxa			Unidentified		
<i>Alnus</i>	-	2	Corroded indet.	11	122
<i>Betula</i>	-	13/6/-	Varia indet.	-	1
<i>Corylus</i>	-	5	Spores		
<i>Pinus</i>	-	5	<i>Botrychium</i>	-	4
<i>Quercus</i>	-	3	<i>Ophioglossum</i>	-	30
<i>Tilia cordata</i> -type	1	1	monoletic fern spore (-perine)	-	19
Dwarf shrub taxa			<i>Polypodium vulgare</i>	1	5/-/1
<i>Calluna</i>	1	14	<i>Pteridium</i>	2	26
Non-arboreal pollen taxa			<i>Sphagnum</i>	-	2
<i>Asteraceae</i> p.p.	1	4	Non-pollen palynomorphs		
<i>Bidens</i> -type	1	-	<i>Botryococcus</i>	-	1
<i>Cerastium</i> -type	-	1	<i>Pseudoschizaea</i>	1	51
<i>Geranium</i>	1	1	HdV-201: <i>Xylomyces</i>	-	1
<i>Wild grass</i> -type	7	18/8/2	HdV-214: <i>Debarya</i> -type	-	1
<i>Hornungia</i> -type	-	2	HdV-461	-	62
Liguliflorae	5	48	Microcharcoal		
<i>Plantago lanceolata</i> -type	-	1/8/-	Particles (>37 µm)	50	6495

Table 3. Results of palynological analysis of soil samples from Mang de Barga LA 57. Indicated are the counts for the different micro-fossil taxa. Where multiple numbers are given, they refer to number of subfossil grains/number of subfossil pollen grains in clumps/number of well-preserved, probably modern, grains.

Near-site investigations

In order to find suitable deposits for palaeo-ecological investigations in the immediate vicinity of Mang de Barga, test gouge coring in a small, wet, seasonally flooded depression ca. 300 m northwest of Mang de Barga was carried out in 2017. The core, labelled MDB1, was extracted using an 80 mm Usinger piston corer. It revealed the presence of peat deposits, lying under a 1.2 m thick modern sand layer. At 2.15 m below the surface, the piston hit an oak log, which impeded further coring.

The peat section of profile MDB1 was continuously subsampled in 1 cm thick slices for palynological and geochemical analyses, as well as for loss-on-ignition examination. The dating of the core is based on three radiocarbon dates on terrestrial plant material and the detection of two cryptotephra layers (Hekla 3 and 4) with their corresponding age estimates from the nearby Lake Belau (cf. Dörfler *et al.* 2012). The chronology of the investigated sequence covers the interval between ca. 3200 BCE and 650 CE, with an average resolution of ca. 40 years per sample.

Given the small size of the depression, and its location in the immediate vicinity of the site, the majority of the pollen grains are understood to have derived from the local and 'extra-local' vegetation, *i.e.* from a radius of a few hundred metres around the archaeological site. This near-site pollen archive enabled a detailed reconstruction of the local land-use history for the entire period of occupation of Mang de Barga.




Figure 5. Selection of palynological and sedimentological taxa and parameters from the near-site profile (MDB 1). Middle Neolithic (MN), Younger Neolithic (YN), Late Neolithic (LN), Older Bronze Age (OBA), Younger Bronze Age (YBA), Iron Age (IA).

Figure 5 shows the percentages of selected palynological and sedimentological proxies recorded in the MDB1 profile. Birch (*Betula*) and hazel (*Corylus*) are light-demanding trees, and their presence can reflect woodland openness. Birch, as a pioneer tree, is generally indicative of woodland regeneration, pointing to the existence of relatively open, early stage secondary woodlands. Birch grows quickly on land left fallow, and this land can be used for forest pasturing. The growth of hazel, on the other hand, is generally promoted by opening of the woodland and can therefore point to woodland disturbance. Further evidence of woodland disturbance at Mang de Barga is provided by the presence of *Pteridium* spores. Bracken (*Pteridium aquilinum*) is a fern favoured by opening of the canopy and is often regarded as an indicator of clearance through burning or woodland pasturing. The former can also be inferred based on the high quantities of micro-charcoal particles in the sediment.

Among the non-arboreal pollen (NAP) types at Mang de Barga, there are standard anthropogenic indicators (cf. Behre 1981), including *Plantago lanceolata*-type and *Rumex acetosa*-type; both of these were also registered in the charred seed assemblage from this site (Table 1). They are understood as reflecting anthropogenic open landscape and agricultural areas, respectively. Ribwort plantain (*Plantago lanceolata*), for example, is an indicator not only of pastoral activities or grassland, but also of arable activities and associated fallow land (Behre 1981; Brinkkemper and van Wijngaarden-Bakker 2005; Kirleis 2019, 76-77). The rising values for *Calluna* possibly signal increasing soil depletion in the anthropogenic parts of the landscape and the associated spread of heather (*Calluna vulgaris*). The amounts of cereal-type pollen can be used to infer the presence and importance of arable farming. Animal grazing/browsing is documented by non-pollen palynomorphs relating to fungi growing on animal dung – so-called coprophilous fungal spores – such as *Sporormiella*-type (HdV-113) and *Podospora*-type (HdV-368) (cf. Feeser and O'Connell 2009; Baker *et al.* 2013).

Another proxy for local land-use activity causing soil erosion is the minerogenic content of the peat, reconstructed based on the results of the loss-on-ignition at 550°C (LOI550), where low values indicate lower organic content of the sediment, and on the concentration of silt particles in the pollen samples.

Changing land-use and cultural practices at Mang de Barga: A multidisciplinary perspective

Here we integrate the results of different analyses (palaeo-environmental, archaeobotanical, archaeological) and use them to chart the history of local land use at Mang de Barga. The fluctuation over time in the quantities of different taxa in the pollen record demonstrates changes in the vegetation forms and/or their extent around the site as a result of low vs. high degree of human impact. Beyond this quantitative aspect of prehistoric human activity at Mang de Barga, the pollen and sedimentological archives allow us to also infer a qualitative aspect – the *nature* of the activity – and changes in it over time.

The near-site pollen diagram (Fig. 5) opens at ca. 3300 BCE and shows a period of increased human activity up until ca. 2900 BCE, based on the quantitative representation of the indicators described above. In culture-historical terms, this time relates to the Middle Neolithic period. The surroundings of Mang de Barga may have been used at this time, but this left no conclusive on-site archaeological evidence.

A further phase of increased land use can be palynologically inferred for the period ca. 2200-1650 BCE, overlapping with the regional Late Neolithic. There are indeed archaeological traces of human activity at Mang de Barga during

this period, in the form of burial mounds containing primary graves (see above). The archaeobotanical evidence indicates cultivation of emmer and barley and perhaps gathering of wild resources (*e.g.* acorn). That agricultural activity took place locally is demonstrated by the palynological analysis and OSL-dating of the fossil soil horizon at LA 57. These results are further supported by the presence of ard marks under at least four burial mounds (LA 18, 57, 60 and 63, cf. Fig. 1C; Schaefer-Di Maida 2020), indicating possible agricultural activities before the erection of the barrows. The finds of spores of coprophilous fungi in the near-site profile MDB1 show that perhaps domestic animals were also kept in the site's surroundings.

There seems to have been a lull in human activity during the second quarter of the 2nd millennium BCE, as indicated by the declining values for anthropogenic indicators in profile MDB1 in the period after ca. 1750 BCE, continuing to their minimum at around 1500 BCE. The archaeological layers broadly associated with this phase did not contain evidence of crop cultivation or consumption. This is in agreement with the archaeological data from Mang de Bergen, which show a decrease or gap in burial activity from around 1700 BCE until about 1500 BCE (the period of Bronze Age Period I in Schleswig-Holstein).

At ca. 1500 BCE, barrow construction activity recommenced at the site, probably reaching maximum levels in a very short time, after which it gradually decreased until 1300 BCE, when it came to an end (Fig. 2). Based on the palynological record, this is associated with a phase of increased grazing activity (a peak in *Sporomiella*-type (HdV-113) and *Podospora*-type (HdV-368)) and low agricultural activity (only sparse records of cereal-type pollen; no archaeobotanical evidence of crops) between ca. 1500 and 1300 BCE (BA Periods I-II).

The time around 1300 BCE is when the new burial rites – cremation and, later on, placement of cremation urns into flat graves – entirely replaced inhumations and barrows at Mang de Bergen and when a sharp rise in burial activity began. The higher number of post-1200 cal BCE burials at Mang de Bergen may imply that more people lived in the area from this time onwards. However, more graves need not reflect demographic growth. In light of the large-scale social change around 1200 BCE, understood as a shift from elite-centred, highly stratified communities to egalitarian communities, it is possible that what used to be a dedicated barrow area in earlier times was now a burial place (also) open to groups previously not entitled to a grave here due to their social status (Schaefer-Di Maida 2020). Either way, the demand for wood may have surged.

The palynological data shows an increasing share of NAP from ca. 1300 BCE onwards, suggesting an increasing opening of the landscape. Local cereal cultivation regained its importance, as indicated by the increasing quantities of cereal-type pollen in the MDB1 core. Furthermore, the change in the relative proportion of woody taxa, *i.e.* a distinct decline in *Corylus* and an associated increase in most arboreal taxa, indicates a change in woodland composition and structure. Further distinct changes in the palaeo-environmental record include increased evidence for soil erosion (*i.e.* high silt particle concentration and lower LOI550) and lower fire activity (*i.e.* low micro-charcoal concentrations).

The latter seems to contradict the evidence for increasing importance of cremations in the area. But, as outlined below, the earlier higher micro-charcoal values are interpreted to primarily reflect the burning of woodland undershrub, which would produce a lot of charcoal in the catchment. Burning of wood in pyres, instead, is much more localised and probably also reaches higher temperatures and more complete combustion, which in turn results in comparatively little charcoal output (cf. Patterson *et al.* 1987; Veal *et al.* 2011).

The palynological evidence for the increasing importance of cereal cultivation is corroborated by the presence of cereal remains in the coeval archaeological

layers, now coming from a greater range of species (Table 2) and possibly including broomcorn millet. This in fact could support the idea of an increase in population during the Younger Bronze Age, reflected not only by an increasing number of graves, but also by a growing demand for agricultural products.

The presented results therefore refine the picture of agricultural history at Mang de Barga presented by Dreibrodt *et al.* (2009), who found no evidence for local Bronze Age agricultural activities. Based on the presence of colluvial layers, Dreibrodt *et al.* (2009) reconstructed main phases of soil erosion, and therefore local agricultural activities, for the Late Neolithic (around 2400/2200 BCE) and, in particular, the Iron Age (c. 250 to 400 CE). Although these two phases are in good agreement with the results from MDB1, it is conceivable that local agriculture and also soil erosion rates had already increased in the Younger Bronze Age, before reaching a maximum in the Iron Age.

At the same time in the woodlands, the decreasing importance of hazel and the associated generally lower representation of bracken points towards less disturbance. Taken together, these trends point to more intensive use of the open land at the same time when the woodlands were used more extensively. This is regarded as signalling a different form of land use from 1200 BCE onwards compared with the earlier phases recorded for the area of Mang de Barga.

A possible explanation is that until 1300 BCE, woodland pasture was an important component. Grazing of animals and periodic cutting of trees for fuel or raw materials would have resulted in the opening of the woodland canopy and affected its structure. This form of land use would have favoured the light-demanding hazel, which constituted woodland undergrowth, and also the proliferation of the disturbance-tolerant bracken. Fire was probably used to improve the accessibility and quality of the grazed vegetation, *e.g.* by encouraging resprouting of plants and thus increasing the amount of forage (cf. Mellars 1976), and fire could have also been applied to clear the land for cultivation of crops.

After 1200 BCE, the record of coprophilous fungal spores suggests pastoral activities, but without clear evidence for woodland disturbance. This could be explained by a shift in pastoral activities to open land. Increased representation of non-arboreal pollen supports this impression; consequently, open grassland may have become the preferred, or the only available, grazing zone. A possible consequence of this intensification of pastoral activities in the open lands around Mang de Barga is the spread of heather due to increasing soil depletion. The distinct increase in coprophilous fungal spores after 1200 BCE in profile MDB1 is associated with an increase in heather. This process seems to have reached a maximum in the two centuries before and after BCE and is in agreement with pedological investigations in the region, which found the first evidence for podsolisation during the Iron Age (Dreibrodt and Wiethold 2015).

To sum up, our results indicate that, although from an archaeological point of view Mang de Barga was, throughout much of its history, principally the site of burial activity, parts of the land locally or in the surrounding region were used for animal grazing and crop cultivation. It seems that, with the complete shift to cremation at 1200 BCE, animal grazing was relocated from one landscape zone (forested) to another (cleared), either in response to modified agricultural practices (*e.g.* new crops and cultivation practices, a change in animal husbandry, and size of herds, a focus on secondary products) or newly emerged human needs (*e.g.* securing wood for funeral and domestic fires, construction or metallurgy), or both.

Culture and land-use changes with a special focus on the 13th to 11th centuries BCE across spatial scales

In this chapter, we place the evidence from Mang de Barga, *i.e.* the local scale, into the broader history of land use and cultural developments by looking at the regional and supra-regional palaeo-environmental, archaeobotanical and archaeological evidence.

We take the well-dated, high-resolution off-site pollen record from Lake Belau, ca. 3.5 km north of Mang de Barga (Wiethold 1998; Dörfler *et al.* 2012; Dreibrodt and Wiethold 2015; Fig. 1B), as representative of the regional palaeo-environmental developments (Fig. 6).

The supra-regional view of the human impact on the environment through time is provided by a combination of pollen, pedo-sedimentological and archaeological radiocarbon data (see details in Feaser *et al.* 2019). The palynological data derive from the annually laminated sediments of Lake Belau (Wiethold 1998; Dörfler *et al.* 2012; Dreibrodt and Wiethold 2015) and Lake Woserin (in Westmecklenburg; Feaser *et al.* 2016). The pedo-sedimentological data consist of records of colluvial deposits in Schleswig-Holstein and Mecklenburg-Vorpommern. The radiocarbon dates come from dozens of single- and multi-period sites in northern Germany (Dreibrodt *et al.* 2010; Hinz *et al.* 2012; Kneisel *et al.* 2013; Fig. 7). These datasets have each been amalgamated into single temporal trajectories, *i.e.* proxy curves for human environmental impact, covering much of the later prehistory of northern Germany. The synchronous trends in the three curves are taken as reflecting changes in the supra-regional population size, where the upward trend signifies an increase in the number of people living in the region, resulting in greater/more intensive use of the landscape and thus changes in the vegetation cover and an increase in soil erosion rates (Fig. 7). These trends have been termed ‘boom phases’ and the intervals between them ‘bust phases’, regarded to reflect growing and declining trends in population numbers, respectively (Feaser *et al.* 2019).

Similar to the data from the local scales, the regional and supra-regional palaeo-environmental data reveal increasing human activity in the last centuries of the 3rd millennium BCE and the first centuries of the 2nd millennium BCE (the Late Neolithic). In both the near-site pollen archive from Mang de Barga and the pollen diagram for Lake Belau (Fig. 6), this is associated with increased evidence for cereal cultivation, clearly discernible at around 2200 BCE (see curves for cereal-type pollen), *i.e.* with the start of the Late Neolithic. Although we know from the local record MDB1 that pastoral activities increased at the same time, it seems likely that clearance of woodlands for agricultural fields was the primary contributor to the changes in the vegetation composition locally, as it was across the wider region (Feaser *et al.* 2019). Given the regular evidence of former arable activities in the soils under the burial mounds, it is unlikely that, even on a local scale, clearing in preparation for burial mound constructions played an important role in opening up the landscape. Rather, it seems that, as in other regions and previous periods, the graves were erected on former farmland, which was thus given a cultural value in addition to an economic one (Doorenbosch 2013; Feaser and Dörfler 2016, 2019).

The archaeobotanical data confirm that agriculture played an important role. Barley, emmer and, to some extent, free-threshing wheat were by this time being grown throughout northern Germany and southern Scandinavia (Kirleis *et al.* 2012; Kirleis and Fischer 2014). In the Late Neolithic, spelt wheat was also taken into cultivation. Spelt remains were discovered inside house structures at the Late Neolithic-Younger Bronze Age site of Todesfelde LA 31, located south of Mang de Barga (Fig. 1A). Spelt grain from one of the houses, from a pit that contained a large concentration of mostly emmer grain, was dated to 2σ 2298–2135 cal BCE (KIA-49888, 3785±30 BP; Effenberger 2018a, Table 3). This is so far the earliest directly dated




Figure 6. Comparison of selected palynological taxa (% total terrestrial pollen) from the regional profile from Lake Belau and the near-site profile (MDB1), Archaeological periods as in Figure 5.




Figure 7. Evidence of supra-regional population dynamics in northern Germany and southern Jutland, Denmark, as derived from multi-proxy investigation (Feaser *et al.* 2019). Vertical bars indicate inferred phases of population growth (green) and decline (red). Archaeological periods as in Figure 5. The map shows the locations of the sites sampled for palynological data (stars), colluvial deposits (red dots) and archaeological radiocarbon dates (blue dots).

find of spelt in northern Germany. There are several reports of spelt wheat from Early Neolithic sites, but they have not been directly dated¹. At Mang de Bergen, a large spelt deposit was encountered at burial location LA 18 (Alsleben pers. comm.). One grain and one spikelet fork from this deposit have been radiocarbon dated by the CRC 1266. Both returned dates falling in the period between the late 15th and early 13th century BCE – the late phases of the Older Bronze Age (grain: KIA-54920, 3116±28 BP, 2σ 1443-1296 cal BCE; spikelet fork: KIA-54921, 3085±29 BP, 2σ 1421-1269 cal BCE). At another settlement site in the region – Bark 2014-264 (Fig. 1A) – a fire pit was discovered that contained cereal remains dated to the end of the 18th or the 17th century BCE (MAMS22199, 3400±21, 2σ 1746-1635 cal BCE). The plant assemblage from the pit comprised mainly emmer and spelt glume bases (Effenberger 2018a, Tables 4, 34). Overall, the archaeobotanical evidence from northern Germany suggests that emmer and spelt were grown as a mixed (maslin) crop (Effenberger 2018a).

From the 17th century BCE onwards, the regional palaeo-environmental data, in agreement with the picture of the local cultural and environmental development, indicate a decline in human activity, reaching a minimum at around 1500 BCE (Figs. 2, 5, 6). Reduced human presence can also be inferred in the wider region from the low number of burials (Aner and Kersten 1978, 1979, 1991, 1993; Aner *et al.* 2005, 2011, 2017) or settlements associated with this period (Bronze Age Period I-II). Generally, data on Older Bronze Age settlements are very limited (cf. Meier 2013) due to insufficient research or difficulties in locating the remains in archaeological surveys. The groups of burial mounds adjacent to Mang de Bergen have been excavated only partially or not at all; moreover, some of the excavations took place in the 19th century and do not offer data of comparable resolution (e.g. the 'Königsbarg' burial mound, with 12 burials, was excavated in 1883 and 1888; Aner *et al.* 2011, 18).

Traces of several Late Neolithic and/or Bronze Age Period I houses have been found at the site of Schmalfeld LA 29 south of Mang de Bergen (Effenberger 2018a); however, they may be older than the 17th century BCE, since only relative chronological determination is provided. Curiously, none of the 250 archaeobotanical samples from this site contain crop remains; only fragments of hazelnut shell and acorn and blackberry seeds were found (Effenberger 2018a, 27). The fire pit at Bark 2014-264 perhaps derives from this period, judging from the date on the cereal remains from it (see above). At another recently excavated burial location in the

¹ Spelt wheat chaff from an Older Bronze Age house at the site of Rothenkirchen, on the island of Rügen, gave a date of 1984-1876 cal BC (KIA-37002) (Alsleben 2019).

same region – Bornhöved LA 117 – remains of barley, emmer, free-threshing wheat, einkorn, lentil (*Lens culinaris*) and broomcorn millet were recovered, but they could be of later date. This barrow was erected in the Late Neolithic and then rebuilt or modified several times, up until the Younger Bronze Age, when cooking pits and urn graves were created on it; settlement pits were also detected here (Kneisel *et al.* in prep.). Two other archaeobotanically analysed sites in the region, Gönnebek 2013-215 and Todesfelde LA 31, yielded only few traces of domestic activity from the Older Bronze Age (Effenberger 2018a, 26-41; Kneisel *et al.* in prep.). Besides residential structures, the latter site also comprised a burial mound from the Older Bronze Age (Lütjens 2012, cited in Effenberger 2018a). Archaeobotanical samples were taken from the mound, the layer of burning at its base, the ancient ploughing horizon underlying the mound, and different layers of the ditch encircling the mound (Effenberger 2018a, 31). Small amounts of barley, emmer and spelt were found in these contexts (Effenberger 2018a, Table 21), but their age is unclear, and they may derive from different phases of use of this multi-period site. In sum, the archaeobotanical evidence from the early period of activity at Mang de Barga fits well into the regional picture, which shows that, if production of crops was practised in the region during the Older Bronze Age, it was on a very low level, leaving only a small imprint on the vegetation and soil cover.

The locally and regionally observed decline in human activity is also reflected at the supra-regional scale, with a ‘bust’ phase from around 1700 BCE to the middle of the 2nd millennium BCE (Feaser *et al.* 2019). A coinciding lull in human environmental impact, generally centred at around 1600/1500 BCE, is also discernible across northern and central Germany (e.g. Hellmund *et al.* 2011; Jahns and Kirleis 2013; Kneisel 2013; Jahns *et al.* 2018). This period of low human environmental impact coincided with the archaeologically documented decrease in human presence/activity throughout central Europe (16th-15th centuries BCE) inferred from the decreased number of settlements from this period (Kneisel *et al.* 2012) or the ‘break in archaeological cultures’ seen in the radiocarbon dates (Kneisel *et al.* 2013; Risch and Meller 2015, 240). The time around 1600-1500 BCE is when the Únětice culture collapsed. This was a prominent bronze production and trade network and a major political power structure of the first half of the 2nd millennium BCE in eastern central Europe. Its demise had a knock-on effect down the line and resulted in dissolution of some local socio-economic entities, including those in the marginal zones of the trade network. Triggered by the social upheaval, a general decentralisation of economic production and political power ensued in Europe after ca. 1500 BCE (Müller and Czebreszuk 2010; Kneisel 2012, 2013; Müller 2013). In some regions, this new political vacuum was instantly filled up by emerging powers, such as the Mycenaean city-states in the Aegean (Risch and Meller 2015, 256). In northern Germany, there is no evidence of anything akin to the dissolution of social structures at this time. Despite the contemporaneous palaeo-environmental evidence of agricultural decline and population decrease around the 17th and 16th centuries BCE, the apparent ‘bust’ was immediately followed by a gradual increase in human presence/activity – a ‘boom’ from ca. 1500 BCE onwards encompassing profound socio-economic changes that culminated in the 12th century BCE.

Between 1500 and 1300 BCE, a strong increase in bronze use, bronze hoarding and associated trade networks can be observed. This further coincides with a rise in the construction of grave mounds. This increase in monumentality and economic expenditure (raw materials, labour distribution and expense) is visible at the local and regional scales. In addition, changes in the spectrum of grave goods are recognisable for this phase, which reveal differentiation in terms of the grave furnishings. On the one hand, a clear difference can be observed between the flat graves without grave goods and the in part richly furnished grave mounds. On the other hand, differences among the burial mounds are also visible, too. This concerns

not only the number of grave goods, which can vary between none and more than nine grave goods, but also the repertoire (Schaefer-Di Maida 2020). Although isolated grave goods involving weaponry are already known from Period I, the frequency and diversity of such grave goods increases distinctly. This is argued to reflect the highlighting of individual personal identity beyond death. These deceased, often termed warrior elites (Vandkilde 1996, 2006; Earle 2002, 363), are indicative of a strong social stratification, which is generally regarded to reflect the emergence of so-called chiefs based on kinship structures (Jensen 1982; Kristiansen 1987, 1991; Earle 1991). Another important aspect during this phase is the emergence and establishment of the three-aisle house structure and thus the enlargement of houses and the use of byre-dwellings. With this change in house construction, a change in the way of living and the household can be assumed.

Despite sporadic first evidence for cremation burials in Schleswig-Holstein from ca. 1500 BCE onwards, it is only around 1300 BCE that a widespread regional change in the treatment of the dead, from inhumation to cremation, can be observed, including at Mang de Bergen (Schaefer-Di Maida 2020). The changed burial ritual can be understood as a local reflection of the general supra-regional ideological shift in mortuary behaviour that took place with the spread of the Urnfield phenomenon across Europe in the last centuries of the 2nd millennium BCE (e.g. Falkenstein 2012). A similar sequence of changes, including the shift to new burial rites at the regional scale, can be inferred from the barrows, although this is reliant on relative, not absolute chronology (Aner and Kersten 1978, 1979, 1991, 1993; Aner *et al.* 2005, 2011, 2017). In contrast to the steady amounts of prestige goods and the continuous construction of burial mounds or secondary burials, the amount of other goods in the graves was decreasing, which may further indicate a divergence in the social structure. Only with the full dominance of urn graves, from Bronze Age Period IV (1100-900 BCE) onwards, can a general decline in the quantities of grave goods be observed, indicating a shift towards a more egalitarian social organisation (Kneisel *et al.* 2019).

Changes also took place in other aspects of life, such as house construction. Starting from the 13th century, the house size was generally smaller, and single-, two- and three-aisled structures were built, perhaps with differing and multiple functions. Schaefer-Di Maida (2020) suggests that the likely smaller number of people now needed for grave construction, on the one hand, and the smaller house sizes in Schleswig-Holstein, on the other, may together indicate the development of smaller household communities around 1200 BCE. Hereby it is conceivable that a reorganisation on the household level may reflect or have brought about changes in subsistence practices.

A recent archaeological survey of the area within a 50 km radius of Lake Woserin confirmed that, following the apparent bust in the Older Bronze Age, human presence/activity increased again around the 12th century BCE, as demonstrated by the reappearance of graves and finds of material culture. Coincidentally, intensified land use is evident from the higher rate of soil erosion at Lake Woserin (Fig. 7), causing a distinct increase in minerogenic input in the lake sediments from ca. 1200 cal BCE onwards (Feeser *et al.* unpubl. data). This goes hand in hand with the evidence for cultivation of a remarkably wide spectrum of crops in northern Germany in the final centuries of the 2nd millennium BCE and early in the 1st millennium BCE (Effenberger 2018a, 2018b; Filipović *et al.* 2019). In addition to the cultivars that had been grown since the Neolithic – emmer, barley, free-threshing wheat, spelt wheat – the cereal spectrum of the Younger Bronze Age included broomcorn millet and possibly oat (*Avena sativa*) and rye (*Secale cereale*). Leguminous and oil/fibre crops seem to have become much more prominent in this period. Common pea (*Pisum sativum*), lentil, flax/linseed (*Linum usitatissimum*) and poppy (*Papaver somniferum*) may have been grown from the Older Bronze Age or the Late Neolithic, whereas

faba bean (*Vicia faba*) and gold-of-pleasure (*Camelina sativa*) were introduced here at the end of the 2nd millennium BCE. The use of wild plants seems to have increased towards and after the turn of the millennium (Effenberger 2018a, Fig. 35). It focused on some previously unused or less-used resources, such as acorns, and perhaps also large-grained grasses growing as crop weeds, for instance rye brome (*Bromus secalinus*) and common wild oat (*Avena fatua*), both of which are relatively abundant in the archaeobotanical assemblages from this period (e.g. Behre 2008; Effenberger 2018a; Filipović *et al.* 2019).

On a supra-regional scale, the archaeobotanical data therefore show that, after the apparent regional break or decline in the use/cultivation of crops in the centuries preceding the 12th century BCE, the degree of agricultural activity increased, relying on the suite of crops known previously, but now with the addition of broomcorn millet (Table 2; Effenberger 2018a, 26-32, 66-69, 2018b; Filipović *et al.* 2020a). The diversification of the crop spectrum seems to have been associated with a change in land-use practices, as suggested by the local and regional palaeo-environmental data. Around 1200 cal BCE and after at Lake Belau, a decline in *Corylus* is observed, along with a gradual rise in the representation of *Calluna*, which is almost identical to the local signal from Mang de Barga (Fig. 6). Thus the changes in land use as reconstructed for Mang de Barga probably represent a local manifestation of the region-wide developments in agropastoral activities and woodland exploitation around the beginning of the Younger Bronze Age – as has already been proposed by Overbeck (1975, 468) for all of northwestern Germany.

Summary and conclusions

Previous and more recent excavations in the Mang de Barga micro-region of northern Germany revealed a series of archaeological features, mostly burials, from the Late Neolithic, the Bronze Age and the Early Iron Age. Near- and on-site palaeo-environmental investigations revealed that the local environment was changing during the course of the site's occupation. Archaeobotanical analysis identified the use of a number of crops typical of Late Neolithic and Bronze Age agriculture in northern Germany, reflecting the general process of diversification of the plant resource base in the late 2nd millennium BCE. The findings offer on-site and off-site perspectives on the local cultural and associated environmental dynamics. Whereas the burials document long-term ritual use of the site, on-site domestic activity has been archaeologically confirmed only for the very late phase of the Bronze Age occupation or later, based on the traces of the Early Iron Age settlement. However, the presence of crops and associated weeds in both early and late periods of site use suggests that processing and perhaps also consumption of crops took place on-site. That the arable fields may have been located in the area is further indicated by the palynological evidence from on- and near-site palaeo-environmental records. Animal herding was also practised in the site's surroundings from very early on, as attested in the near-site palaeo-environmental record.

The shift in the funerary custom in the 13th-11th centuries BCE and the subsequent 'boom' in burial activity at Mang de Barga coincide with a number of other changes in human lifeways including subsistence and land-use practices. Not only did the observed changes in the ritual and economic spheres of life probably take place at about the same time, they were also intertwined or causally related. Coinciding evidence for social restructuring and associated changes in landscape exploitation on the regional and supra-regional scale suggest that the local developments reflect and parallel developments on a much larger spatial scale.

This supports our idea that the multidisciplinary reconstruction of environmental and cultural change at Mang de Barga can be used to improve our understanding

of the wider socio-environmental transformation in the period between the 13th and 11th centuries BCE in northern Germany. We therefore suggest that the specific activities identified as drivers of changes in land use locally are pertinent to the reconstruction of the regional and supra-regional developments.

The quantification of on-site (archaeological) and near-site anthropogenic indicators at Mang de Barga documents high-intensity use of the site and the surrounding area from around 1200 BCE onwards. The qualitative evidence points to specific human actions and their possible consequences. Cremation burials are a testament to the cutting and collection of wood for the purpose of building and fuelling funeral pyres. A changing demand for wood – not just for funeral and domestic fires, but also for construction or metallurgy – could have been one reason for changes in woodland management or usage. The proliferation of open land in the surroundings was a result of intensified land use, at least partly resulting from the shift from woodland to grassland pasture. Further, despite their poor preservation and low quantity, the archaeobotanical remains document processing and consumption of a wide range of crops in the last centuries of the 2nd millennium BCE. At least some were cultivated or processed near the site, as suggested by the presence of cereal-type pollen in the near-site core. Indeed, broomcorn millet cultivation, which started in northern Germany around 1200 BCE, would have allowed expansion onto previously uncultivated soils (e.g. those that would have been waterlogged in cold seasons), thus increasing human impact on the landscape. Expanding and more intensive use of areas under arable farming could have driven soil erosion and depletion, especially if the same land was used for both animal keeping and crop growing – for instance in a system where fallowing was practised and animals were grazed on fallow fields or where farming of the same fields extended throughout the year, for example, with wheat in winter and broomcorn millet in summer, as occurs in multi-cropping regimes described ethnographically by, for instance, Buttler (1999).

The societal transformation of the 13th–11th centuries BCE in northern Germany entailed fundamental changes in cultural practices, such as the shift in burial rites, changes in the number and layout of settlements and houses, innovations and new developments in agropastoral production, and a significant increase in the number of people residing in this part of Europe from this period onwards (Feaser *et al.* 2019; Kneisel *et al.* 2019). The transformation represented an interplay of different, interconnected social and economic factors. The synchronous patterns we see in the palaeo-environmental records from a range of spatial scales capture the imprint on the landscape left by this transformation. This process is best discernible at the local scale. At Mang de Barga, we have been able to directly link changes in the on-site and off-site cultural practices through the reconstruction of the potential impact on the environment of the shift in burial customs and intensified food production. The present study, therefore, highlights the potential of combining complementary data from different spatial scales and disciplines to improve our understanding of socio-environmental transformations.

Acknowledgements

This research was conducted and financed in the context of the Collaborative Research Centre (CRC) 1266 ‘Scales of Transformation – Human-Environmental Interaction in Prehistoric and Archaic Societies’ of the German Research Foundation (DFG, project number 2901391021 – SFB 1266). Coring assistance for the MDB1 profile was provided by W. Dörfler, C. Floors, S. Ibens and A. Oflaz. C. Floors, S. Ibens and T. Reiser assisted with sample processing and laboratory analyses. Y. Dannath identified wood charcoal fragments selected for ¹⁴C dating. The authors thank J. Wiethold from the Institut national de recherches archéologiques préventives (Inrap), Metz, France, for providing pollen data for Lake Belau. Further thanks

go to the Archäologisches Landesamt Schleswig-Holstein for providing access to the data and materials from Mang de Bergen, and to A. Alsleben, Akademie der Wissenschaften und der Literatur in Mainz, for preliminary information on the archaeobotanical remains from two locations at Mang de Bergen and for permitting the radiocarbon dating of spelt remains from LA 18.

References

- Alsleben, A., 2019. The fossil plant remains from the Early Bronze Age site of Roth-enkirchen on Rügen: spatial distribution patterns as a reflection of household activities. In: M. Dal Corso, W. Kirleis, J. Kneisel, N. Taylor, M. Wieckowska-Lüth, M. Zanon, eds., *How's Life? Living Conditions in the 2nd and 1st Millennia BCE*. Leiden: Sidestone Press, 85-101. <https://www.sidestone.com/books/how-s-life>
- Aner, E. and Kersten, K., 1978. *Südschleswig-Ost: Die Kreise Schleswig-Flensburg und Rendsburg-Eckernförde (nördlich des Nord-Ostsee-Kanals)*. Die Funde der älteren Bronzezeit des nordischen Kreises in Dänemark, Schleswig-Holstein und Niedersachsen 4. Neumünster: Wachholtz.
- Aner, E. and Kersten, K., 1979. *Südschleswig-West: Nordfriesland*. Die Funde der älteren Bronzezeit des nordischen Kreises in Dänemark, Schleswig-Holstein und Niedersachsen 5. Neumünster: Wachholtz.
- Aner, E., and Kersten, K., 1991. *Dithmarschen*. Die Funde der älteren Bronzezeit des nordischen Kreises in Dänemark, Schleswig-Holstein und Niedersachsen 17. Neumünster: Wachholtz.
- Aner, E. and Kersten, K., 1993. *Kreis Steinburg*. Die Funde der älteren Bronzezeit des nordischen Kreises in Dänemark, Schleswig-Holstein und Niedersachsen 18. Neumünster: Wachholtz.
- Aner, E., Kersten, K., Willroth, K.-H., 2005. *Kreis Rendsburg-Eckernförde (südlich des Nord-Ostsee-Kanals) und die kreisfreien Städte Kiel und Neumünster*. Die Funde der älteren Bronzezeit des nordischen Kreises in Dänemark, Schleswig-Holstein und Niedersachsen 19. Neumünster: Wachholtz.
- Aner, E., Kersten, K., Willroth, K.-H., 2011. *Kreis Segeberg*. Die Funde der älteren Bronzezeit des nordischen Kreises in Dänemark, Schleswig-Holstein und Niedersachsen 20. Neumünster: Wachholtz.
- Aner, E., Kersten, K., Willroth, K.-H., 2017. *Kreise Plön und Ostholstein und die kreisfreie Hansestadt Lübeck*. Die Funde der älteren Bronzezeit des nordischen Kreises in Dänemark, Schleswig-Holstein und Niedersachsen 21. Neumünster: Wachholtz.
- Baker, A.G., Bhagwat, S.A., Willis, K.J., 2013. Do dung fungal spores make a good proxy for past distribution of large herbivores? *Quaternary Science Reviews*, 62, 21-31. <https://doi.org/10.1016/j.quascirev.2012.11.018>
- Behre, K.-E., 1981. The interpretation of anthropogenic indicators in pollen diagrams. *Pollen et Spores*, 23, 225-245.
- Behre, K.-E., 2008. Collected seeds and fruits from herbs as prehistoric food. *Vegetation History and Archaeobotany*, 17, 65-73. <https://doi.org/10.1007/s00334-007-0106-x>
- Brinkkemper, O. and Van Wijngaarden-Bakker, L., 2005. All-round farming: food production in the Bronze Age and the Iron Age. In: L.P. Louwe Kooijmans, P.W. van der Broeke, H. Fokkens, A.L. van Gijn, eds., *The Prehistory of the Netherlands: Volume 2*. Amsterdam: Amsterdam University Press, 491-512.
- Buttler, A., 1999. Traditional seed cropping systems in the temperate Old World: models for antiquity. In: C. Gosden and J. Hather, eds., *The Prehistory of Food: Appetites for Change*. London: Routledge, 452-467.
- Doorenbosch, M., 2013. *Ancestral Heaths: Reconstructing the Barrow Landscape in the Central and Southern Netherlands*. Leiden: Sidestone Press. <https://www.sidestone.com/books/ancestral-heaths>

- Dörfler, W., Feeser, I., Van den Bogaard, C., Dreibrodt, S., Erlenkeuser, H., Kleinmann, A., Merkt, J., Wiethold, J., 2012. A high-quality annually laminated sequence from Lake Belau, northern Germany: revised chronology and its implications for palynological and tephrochronological studies. *The Holocene*, 22 (12), 1413-1426. <https://doi.org/10.1177%2F0959683612449756>
- Dreibrodt, S. and Wiethold, J., 2015. Lake Belau and its catchment (northern Germany): a key archive of environmental history in northern central Europe since the onset of agriculture. *The Holocene*, 25 (2), 296-322. <https://doi.org/10.1177%2F0959683614558648>
- Dreibrodt, S., Nelle, O., Lütjens, I., Mitusov, A., Clausen, I., Bork, H.-R., 2009. Investigations on buried soils and colluvial layers around Bronze Age burial mounds at Bornhöved (northern Germany) – An approach to test the hypothesis of 'landscape openness' by the incidence of colluviation. *The Holocene*, 19 (3), 481-491. <https://doi.org/10.1177%2F0959683608101397>
- Dreibrodt, S., Lubos, C., Terhorst, B., Damm, B., Bork, H.-R., 2010. Historical soil erosion by water in Germany: scales and archives, chronology, research perspectives. *Quaternary International*, 222 (1-2), 80-95. <https://doi.org/10.1016/j.quaint.2009.06.014>
- Earle, T., 1991. The evolution of chiefdoms. In: T. Earle, ed., *Chiefdoms, Power, Economy, Ideology*. Cambridge: Cambridge University Press, 1-15.
- Earle, T., 2002. *Bronze Age Economics: The Beginning of Political Economies*. Boulder: Westview Press.
- Effenberger, H., 2018a. *Pflanzennutzung und Ausbreitungswege von Innovationen im Pflanzenbau der Nordischen Bronzezeit und angrenzender Regionen*. Kiel: Wachholtz.
- Effenberger, H. 2018b. The plant economy of the northern European Bronze Age – More diversity through increased trade with southern regions. *Vegetation History and Archaeobotany*, 27, 6574. <https://doi.org/10.1007/s00334-017-0621-3>
- Falkenstein, F., 2012. The development of burial rites from the Tumulus to the Urnfield culture in southern central Europe. In: E. Borgna and S. Müller Celka, eds., *Ancestral Landscapes: Burial Mounds in the Copper and Bronze Ages (Central and Eastern Europe – Balkans – Adriatic – Aegean, 4th-2nd Millennium B.C.)*. Proceedings of the international conference held in Udine, May 15th-18th 2008. Travaux de la Maison de l'Orient et de la Méditerranée 58. Lyon: Maison de l'Orient et de la Méditerranée Jean Pouilloux, 329-340.
- Falkenstein, F., 2017. Zum Wandel der Bestattungssitten von der Hügelgräber- zur Urnenfelderkultur in Süddeutschland. In: D. Brandherm and B. Nessel, eds., *Phasenübergänge und Umbrüche im bronzezeitlichen Europa*. Universitätsforschungen zur prähistorischen Archäologie 297. Bonn: Dr. Rudolf Habelt GmbH, 77-96.
- Feeser, I. and O'Connell, M., 2009. Fresh insights into long-term changes in flora, vegetation, land use and soil erosion in the karstic environment of the Burren, western Ireland. *Journal of Ecology*, 97 (5), 1083-1100. <https://doi.org/10.1111/j.1365-2745.2009.01533.x>
- Feeser, I. and Dörfler, W., 2016. Landschaftsentwicklung und Landnutzung. In: H. Dibbern, ed., *Das trichterbecherzeitliche Westholstein: eine Studie zur neolithischen Entwicklung von Landschaft und Gesellschaft*. Frühe Monumentalität und soziale Differenzierung 8. Bonn: Dr. Rudolf Habelt GmbH, 17-24.
- Feeser, I., Dörfler, W., Czymzik, M., Dreibrodt, S., 2016. A mid-Holocene annually laminated sediment sequence from Lake Woserin: The role of climate and environmental change for cultural development during the Neolithic in Northern Germany. *The Holocene*, 26 (6), 947-963. <https://doi.org/10.1177/0959683615622550>
- Feeser, I. and Dörfler, W., 2019. Palynologische Untersuchungen zum Bestattungsort Wangels LA 69. *Journal of Neolithic Archaeology*, 21, 89-102. <https://doi.org/10.12766/jna.2019.5>

- Feeser, I., Dörfler, W., Kneisel, J., Hinz, M., Dreibrodt, S., 2019. Human impact and population dynamics in the Neolithic and Bronze Age: multi-proxy evidence from north-western central Europe. *The Holocene*, 29 (10), 1596-1606. <https://doi.org/10.1177%2F0959683619857223>
- Feeser, I., Van den Bogaard, C., Dörfler, W., 2020. Palaeoecological investigations at the archaeological site Mang de Barga, Bornhöved, Kreis Segeberg: new insights into local to over-regional land-use changes during the Bronze Age. In: S. Schaefer-Di Maida, *Bronzezeitliche Transformationsprozesse in Schleswig-Holstein am Beispiel vom Fundplatz von Mang de Barga, Bornhöved (Kr. Segeberg)*. Doctoral dissertation, University of Kiel, Appendix VII.3.
- Filipović, D., Mewis, F., Saalow, L., Schmidt, J.P., Kirleis, W., 2019. Waste disposal in the Bronze Age: plants in pits at Wismar-Wendorf, northern Germany. In: M. Dal Corso, W. Kirleis, J. Kneisel, N. Taylor, M. Wieckowska-Lüth, M. Zanon, eds., *How's Life? Living Conditions in the 2nd and 1st Millennia BCE*. Leiden: Sidestone Press, 103-130.
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Kloß, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H., Stobbe, A., Tolar, T., Wasylukowa, K., Wiethold, J., Zerl, T., 2020a. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Filipović, D., Reiser, T., Dannath, Y., 2020b. Archaeobotanical remains from selected Mang de Barga sites, northern Germany. In: S. Schaefer-Di Maida, *Bronzezeitliche Transformationsprozesse in Schleswig-Holstein am Beispiel vom Fundplatz von Mang de Barga, Bornhöved (Kr. Segeberg)*. Doctoral dissertation, University of Kiel, Appendix VII.2.
- Havinga, A.J., 1984. A 20-year experimental investigation into the differential corrosion susceptibility of pollen and spores in various soil types. *Pollen et Spores*, 26, 541-558.
- Hellmund, M., Wennrich, V., Becher, H., Krichel, A., Brühlheide, H., Melles, M., 2011. Zur Vegetationsgeschichte im Umfeld des Süßen Sees, Lkr. Mansfeld-Südharz – Ergebnisse von Pollen- und Elementanalysen. In: H.-R. Bork, H. Meller, R. Gerlach, eds., *Umweltarchäologie – Naturkatastrophen und Umweltwandel im archäologischen Befund*. 3. Mitteldeutscher Archäologentag vom 07. bis 09. Oktober 2010 in Halle (Saale). Halle (Saale): Landesamt für Denkmalpflege und Archäologie in Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 111-127.
- Hinz, M., Furholt, M., Müller, J., Raetzl-Fabian, D., Rinne, C., Sjögren, K.-G., Wotzka, H.-P., 2012. RADON – Radiocarbon Dates Online 2012: Central European and Scandinavian Database of 14C Dates for the Neolithic and Early Bronze Age. *Journal of Neolithic Archaeology*. <https://doi.org/10.12766/jna.2012.65>
- Jahns, S. and Kirleis, W., 2013. Die bronzezeitliche Besiedlung in Pollendiagrammen aus Brandenburg. In: K.-H. Willroth, ed., *Siedlungen der älteren Bronzezeit: Beiträge zur Siedlungsarchäologie und Paläoökologie des zweiten vorchristlichen Jahrtausends in Südsandinavien, Norddeutschland und den Niederlanden*. Workshop vom 7.-9.4.2011 in Sankelmark. Studien zur nordeuropäischen Bronzezeit 1. Neumünster: Wachholtz, 239-246.
- Jahns, S., Alsleben, A., Bittmann, F., Brande, A., Christiansen, J., Dannath, Y., Effenberger, H., Giesecke, T., Jäger, K.-D., Kirleis, W., Kloß, S., Kloss, K., Kroll, H., Lange, E., Medović, A., Neef, R., Stika, H.-P., Sudhaus, D., Wiethold, J., Wolters, S., 2018. Zur Geschichte der nacheiszeitlichen Umwelt und der Kulturpflanzen im Land Brandenburg. In: S. Flohr and P. Morgenstern, eds., *Beiträge zur Archäozoologie und Prähistorischen Anthropologie: Band XI*. Langenweißbach: Beier & Beran, 9-35.

- Jensen, J., 1982. *The Prehistory of Denmark*. London: Methuen.
- Kirleis, W., 2003. Vegetationsgeschichtliche und archäobotanische Untersuchungen zur Landwirtschaft und Umwelt im Bereich der prähistorischen Siedlungen bei Rullstorf, Ldkr. Lüneburg. *Probleme der Küstenforschung im südlichen Nordseegebiet*, 28, 65-132.
- Kirleis, W., 2019. *Atlas of Neolithic Plant Remains from Northern Central Europe*. Groningen: Barkhuis Publishing.
- Kirleis, W. and Fischer, E., 2014. Neolithic cultivation of tetraploid free threshing wheat in Denmark and northern Germany: implications for crop diversity and societal dynamics of the Funnel Beaker culture. *Vegetation History and Archaeobotany*, 23, 81-96. <https://doi.org/10.1007/s00334-011-0328-9>
- Kirleis, W., Kloof, S., Kroll, H., Müller, J., 2012. Crop growing and gathering in the northern German Neolithic: a review supplemented by new results. *Vegetation History and Archaeobotany*, 21, 221-242. <https://doi.org/10.1007/s00334-011-0328-9>
- Kneisel, J., 2012. The problem of the Middle Bronze Age inception in northeast Europe or: did the Únětice society collapse? In: J. Kneisel, W. Kirleis, M. Dal Corso, N. Taylor, V. Tiedtke, eds., *Collapse or Continuity? Environment and Development of Bronze Age Human Landscapes*. Proceedings of the International Workshop Socio-Environmental Dynamics over the Last 12,000 Years: The Creation of Landscapes II (14th-18th March 2011) in Kiel. Universitätsforschungen zur prähistorischen Archäologie 205. Bonn: Dr. Rudolf Habelt GmbH, 209-233.
- Kneisel, J., 2013. Der Übergang von der Frühbronzezeit zur Mittelbronzezeit im nordöstlichen Mitteleuropa: Lücke oder Forschungsdesiderat? In: J. Kneisel, H.J. Behnke, A. Schopper, eds., *Frühbronzezeit-Mittelbronzezeit: Neue Erkenntnisse zur Besiedlung zwischen Elbe und Warthe und angrenzender Regionen (2000-1400 v. Chr.)*. Symposium vom 24.-25. Sept. 2011 in Welzow, Brandenburg. Studien zur Archäologie in Ostmitteleuropa 10. Bonn: Dr. Rudolf Habelt GmbH, 95-120.
- Kneisel, J., Kirleis, W., Dal Corso, M., Taylor N., 2012. Collapse or Continuity? Concluding remarks on the environment and development of Bronze Age human landscapes. In: J. Kneisel, W. Kirleis, M. Dal Corso, N. Taylor, V. Tiedtke, eds., *Collapse or Continuity? Environment and Development of Bronze Age Human Landscapes*. Proceedings of the International Workshop Socio-Environmental Dynamics over the Last 12,000 Years: The Creation of Landscapes II, 14th-18th March 2011 in Kiel. Universitätsforschungen zur prähistorischen Archäologie 205. Bonn: Dr. Rudolf Habelt GmbH, 267-279.
- Kneisel, J., Hinz, M., Rinne, C., 2013. *RADON-B – Radiocarbon Dates Online (version 2014): Database for European Dates for the Bronze and Early Iron Age*. Webproject at Kiel University. <http://radon-b.ufg.uni-kiel.de>
- Kneisel, J., Brinkmann, J., Corradini, E., Erkul, E., Feeser, I., Panning, D., Pickartz, N., Rabbal, W., Schaefer, S., Stümpel, H., 2017. Mang de Barga bei Bornhöved – Kleine Region, große Fragen. *Archäologische Nachrichten aus Schleswig-Holstein*, 23, 48-55.
- Kneisel, J., Dörfler, W., Dreibrodt, S., Schaefer-Di Maida, S., Feeser, I., 2019. Cultural change and population dynamics during the Bronze Age: Integrating archaeological and palaeoenvironmental evidence for Schleswig-Holstein, Northern Germany. *The Holocene*, 29 (10), 1607-1621. <https://doi.org/10.1177/0959683619857237>
- Kneisel, J., Schaefer-Di Maida, S., Feeser, I., Dreibrodt, S., Filipović, D. in prep. Leben und Ansichten von LA 117 und LA 57, Grabhügel: Vorbericht über die Ausgrabungen bronzezeitlicher Grabhügel bei Bornhöved, Kr. Segeberg.
- Kristiansen, K., 1987. From stone to bronze: the evolution of social complexity in northern Europe, 2300-1200 BC. In: E.M. Brumfield and T. Earle, eds., *Specialization, Exchange and Complex Societies*. Cambridge: Cambridge University Press, 30-51.
- Kristiansen, K., 1991. Chieftdoms, states and systems of social evolution. In: T. Earle, ed., *Chieftdoms: Power, Economy and Ideology*. Cambridge: Cambridge University Press, 16-43.

- Kristiansen, K., 2018. The rise and fall of the Bronze Age societies in Thy, northwest Jutland. In: J.-H. Bech, B.V. Eriksen, K. Kristiansen, eds., *Bronze Age Settlement and Land-Use in Thy, Northwest Denmark: Volume 1*. Højbjerg: Jutland Archaeological Society, 107-132.
- Lütjens, I., 2012. Todesfelde, Kreis Segeberg, LA 31: Siedlung des Spätneolithikums und/oder frühen Bronzezeit, Grabhügel der älteren Bronzezeit, Siedlungen der späten Bronze-/frühen Eisenzeit und Brandgräberfeld der älteren Vorrömischen Eisenzeit: Bericht über die archäologische Hauptuntersuchung 2012. Excavation report, Archäologisches Landesmuseum Schleswig in der Stiftung Schleswig-Holsteinische Landesmuseen, Neumünster.
- Meier, D., 2013. Eine Siedlung der älteren Bronzezeit in Schleswig-Holstein. In: K.-H. Willroth, ed., *Siedlungen der älteren Bronzezeit: Beiträge zur Siedlungsarchäologie und Paläoökologie des zweiten vorchristlichen Jahrtausends in Südschweden, Norddeutschland und den Niederlanden*. Workshop vom 7. bis 9. April 2011 in Sankelmark. Studien zur nordeuropäischen Bronzezeit 1. Neumünster: Wachholtz Verlag, 91-118.
- Mellars, P., 1976. Fire ecology, animal populations and man: a study of some ecological relationships in prehistory. *Proceedings of the Prehistoric Society*, 42, 15-45.
- Moore, P.D., Webb, J.A., Collinson, M.E., 1991. *An Illustrated Guide to Pollen Analysis*. Oxford: Blackwell Scientific Publications.
- Müller, J., 2013. 1600 B.C. – social topographies and the development of Early Bronze Age societies in central Europe. In: H. Meller, F. Bertemes, H.-R. Bork, R. Risch, eds., *1600 – Kultureller Umbruch im Schatten des Thera-Ausbruchs*. 4. Mitteldeutscher Archäologentag vom 14. bis 16. Oktober 2011 in Halle (Saale). Tagungen des Landesmuseums für Vorgeschichte Halle (Saale) 9. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 527-538.
- Müller, J. and Czebreszuk, J., 2010. Bruszczewo und Łęki Małe – ein frühbronzezeitliches Machtzentrum in Großpolen. In: H. Meller and F. Bertemes, eds., *Der Griff nach den Sternen*. Internationales Symposium in Halle 16.-21.2. 2005. Tagungen des Landesmuseums für Vorgeschichte Halle (Saale) 5/1. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 381-396.
- Overbeck, F., 1975. *Botanisch-geologische Moorkunde*. Neumünster: Wachholtz Verlag.
- Patterson, W.A., Edwards, K.J., Maguire, D.J., 1987. Microscopic charcoal as a fossil indicator of fire. *Quaternary Science Reviews*, 6, 3-23. [https://doi.org/10.1016/0277-3791\(87\)90012-6](https://doi.org/10.1016/0277-3791(87)90012-6)
- Piotrowski, J., 1991. *Quartär- und hydrogeologische Untersuchungen im Bereich der Bornhöveder Seenkette, Schleswig-Holstein*. Berichte / Geologisch-Paläontologisches Institut und Museum, 43. Kiel: Geologisch-Paläontologisches Institut und Museum, Christian-Albrechts-Universität.
- Risch, R. and Meller, H., 2015. Change and continuity in Europe and the Mediterranean around 1600 BC. *Proceedings of the Prehistoric Society*, 81, 239-264. <https://doi.org/10.1017/ppr.2015.10>
- Roehrs, H., Kloos, S., Kirleis, W., 2013. Evaluating prehistoric finds of *Arrhenatherum elatius* var. *bulbosum* in north-western and central Europe with an emphasis on the first Neolithic finds in northern Germany. *Archaeological and Anthropological Sciences*, 5, 1-15. <https://doi.org/10.1007/s12520-012-0109-0>
- Saalow, L., Schmidt, J.-P., Selent, A., 2014. 340 km Ausgrabung – Planung, Durchführung und Ergebnisse der Untersuchung im Trassenverlauf. In: A. Dworzak, U.M. Meier, L. Saalow, H. Gräfin von Schmettow, J.-P. Schmidt, A. Selent, eds., *Pipeline Archäologie: Ausgrabungen auf den großen Ferngastrassen in Meck-*

- lenburg-Vorpommern. Schwerin: Landesamt für Kultur und Denkmalpflege Archäologie und Denkmalpflege, 23-38.
- Schaefer-Di Maida, S., 2020. *Bronzezeitliche Transformationsprozesse in Schleswig-Holstein am Beispiel vom Fundplatz von Mang de Bergen, Bornhöved (Kr. Segeberg)*. Doctoral dissertation, University of Kiel.
- Schmidt, J.-P., 2010. Neue Erkenntnisse zu bronzezeitlichen Siedlungen in Nordostdeutschland. In: M. Meyer, ed., *Haus – Gehöft – Weiler – Dorf: Siedlungen der Vorrömischen Eisenzeit im nördlichen Mitteleuropa*. Internationale Tagung an der Freien Universität Berlin vom 20.-22. März 2009. Berliner archäologische Forschungen 8. Rahden: Leidorf, 85-99.
- Schunke, T., 2018. Fahren oder Tragen? Methoden des Antransports von Baumaterial für bronzezeitliche Grabhügel. In: B. Nessel, D. Neumann, M. Bartelheim, eds., *Bronzezeitlicher Transport: Akteure, Mittel und Wege*. Tübingen: Tübingen University Press, 85-115.
- Schwerin von Krosigk, H.G., 1976. *Untersuchungen zum vor- und frühgeschichtlichen Siedlungsablauf am Fundbild der Gemarkungen Bornhöved, Gönnebek, Groß-Kummerfeld, Schmalensee, Kreis Segeberg/Holstein*. Schleswig: Verein zur Förderung des Schleswig-Holsteinischen Landesmuseums für Vor- und Frühgeschichte.
- Vandkilde, H., 1996. *From Stone to Bronze: the Metalwork of the Late Neolithic and Earliest Bronze Age in Denmark*. Aarhus: Aarhus University Press.
- Vandkilde, H., 2006. Archaeology and war: presentations of warriors and peasants in archaeological interpretations. In: T. Otto, H. Thrane, H. Vandkilde, eds., *Warfare and Society: Archaeological and Social Anthropological Perspectives*. Aarhus: Aarhus University Press, 57-73.
- Veal, R., O'Donnell, L., McParland, L., 2011. Measuring burn temperatures from charcoal using the reflectance method, first results from an Irish Bronze Age cremation site. *Saguntum: Papeles del Laboratorio de Arqueología de Valencia-Extra*, 11, 21-22.
- Wiethold, J., 1998. *Studien zur jüngeren postglazialen Vegetations- und Siedlungsgeschichte im östlichen Schleswig-Holstein*. Bonn: Dr. Rudolf Habelt GmbH.

**SECTION 2: APPROACHING MILLET
CULTIVATION AND CONSUMPTION
THROUGH HIGH-END MICROSCOPY,
CHEMISTRY AND ETHNOGRAPHY**

Putting millet into a culinary context: Organic residue analysis and the identification of *Panicum miliaceum* in pottery vessels

Edward A. Standall, Oliver E. Craig, Carl Heron

Abstract

Broomcorn millet (*Panicum miliaceum*) is a cereal of global significance. From domestication in East Asia, to translocation across Eurasia, this species significantly influenced subsistence strategies and agricultural systems, presenting a new way of life for many communities. Previous research has focused on the investigation of this species in terms of subsistence, yet few studies have explored the cultural implications of its translocation and adoption in prehistory. By investigating the incorporation of broomcorn millet into cuisine, archaeologists are able to examine nuanced cultural practices and infer the significance of this species to past peoples and cultures. To explore subtle and complex uses of this cereal, in culinary contexts, researchers are applying organic residue analysis (ORA) and the biomarker approach to detect miliacin, a discrete compound concentrated in *P. miliaceum* caryopses. This paper summarises the motivations, methods, and benefits of ORA, in the investigation of *P. miliaceum*, and presents a review of previously published studies that attempt to elucidate past cultural activities incorporating broomcorn millet. In doing so, this paper outlines current limitations in the interpretation of ‘millet residues’ in pottery vessels and details areas of future research that are necessary to better explore the uses of broomcorn millet in archaeological contexts.

Introduction

Broomcorn millet (*Panicum miliaceum*) is thought to have been domesticated in northeastern China around 6000 BCE (Zhao 2011), although the chronology and nature of the domestication process remain topics of significant debate (Miller *et al.* 2016; Leipe *et al.* 2019; Stevens *et al.* 2020). Broomcorn millet is among the earliest domesticated cereals globally and represents an important developmental driver

of early societies in China. It spread relatively slowly across China, but expanded rapidly westwards, across Eurasia, in the third and second millennium BCE (Frachetti *et al.* 2010; Motuzaite-Matuzeviciute 2013; Spengler *et al.* 2018; Leipe *et al.* 2019; Filipović *et al.* 2020). Agricultural communities, of eastern and western traditions, likely converged around the northeastern terminus of the 'Inner Asian Mountain Corridor' (IAMC) at the beginning of the third millennium BCE, with probable evidence for *P. miliaceum* in Kazakhstan from ca. 2700 cal BCE (Hermes *et al.* 2019) and direct evidence of wheat (*Triticum* spp.) in China from ca. 2600 cal BCE (Long *et al.* 2018). The adoption of a new cereal is neither passive nor mundane and may occur concurrently with cultural, social and economic change. Therefore, the means and motives of cereal translocation have caused considerable debate, from which one dominant question arises: 'Why move starchy cereals?' (Lightfoot *et al.* 2013).

Archaeobotanical evidence indicates that broomcorn millet was incorporated into various cropping regimes in Europe (Kapcia and Mueller-Bieniek 2019; Mueller-Bieniek *et al.* 2019) and ecological opportunism, for either economic gain or food security, represents a compelling motive for its adoption (Jones *et al.* 2011). Furthermore, environmental limitations potentially provide an explanation as to why some communities did not adopt this species (Miller 2015; Miller *et al.* 2016). However, neither provides a universal explanation for the adoption and rejection of broomcorn millet across Eurasia, as is evident from archaeobotanical (Miller *et al.* 2016; Valamoti 2016) and isotopic (Lightfoot *et al.* 2013; Wang *et al.* 2017; Hermes *et al.* 2019) investigations.

Cultural and social motivations for the introduction of new cereals are emphasised by Boivin *et al.* (2012). Millet would have enriched culinary practices and cuisine, providing a different taste and texture from western cereals. These characteristics may correspond with ideas of exotica, prestige, and conspicuous consumption. Conversely, new species may be perceived negatively and either shunned or proscribed within and between communities. Indeed, attitudes may differ on either a localised social scale or more broadly on a cultural scale. The exotic nature of introduced products may result in their incorporation into either new or existing cultural frameworks that are influenced by either internal or external factors. Furthermore, the perception and uses of a new foodstuff are subject to change over time. However, it is important to consider how cultural ideas and practices are represented in the archaeological record and whether our investigative techniques are sufficient to elucidate them.

Reconstructing *P. miliaceum* use in cuisine

One method of understanding the role and perception of foodstuffs in prehistory is by investigating the culinary context of their use. The selection, preparation and consumption of different foods is culturally mediated and deeply rooted in individual and group identity (Hastorf 2017). Separation and combination of different products, in daily meals and celebratory events, may demonstrate their significance and meaning. However, the intricate scale of analysis required to study culinary activity necessitates consideration of the material culture employed in the manipulation, processing, cooking, and consumption of foodstuffs. Indeed, understanding the role of pottery containers, where archives of food preparation and consumption are preserved microscopically and at the molecular level, is integral to enhancing knowledge of cultural practices.

A direct association between millet and material culture, in this case pottery vessels, is challenging to establish. Impressions in pottery, thought to be made by millet caryopses and interpreted as evidence for cultivation of *P. miliaceum*, are often erroneously identified (An *et al.* 2019). There are very occasional caches of millet caryopses (*e.g.* Reed and Drnić 2016) and processed millet foods, such as

noodles (Lu *et al.* 2005; Gong *et al.* 2011), that are preserved in either intact or near-intact pottery vessels. Microscopy has been used to identify millet grains in charred surface deposits adhering to pottery vessels, with observations supported by isotope analysis (Miyata *et al.* 2007). Additionally, starch grains and phytoliths, recovered from the surface of pottery vessels, as well as other objects such as grinding stones, have been used to identify millet residues (Yang *et al.* 2012a, b; Dal Corso *et al.* 2017; Liu *et al.* 2020). However, arguably the most promising development, in the identification of processing *P. miliaceum* in ceramic vessels, is the application of combined molecular and isotopic analysis (Heron *et al.* 2016).


The aim of this paper is to review and evaluate the role of organic residue analysis (ORA) in detecting *P. miliaceum* processing in pottery vessels. Emphasis is placed on the molecular characterisation of lipid residues in ceramic-absorbed and visible surface deposits (e.g. charred ‘foodcrusts’). We review and critically assess recent findings of *P. miliaceum* residues and evaluate some of the current limitations of ORA in the identification of broomcorn millet. Finally, we explore areas of new research that are needed to take these enquiries forward.

Organic residue analysis

The analysis of organic residues, extracted from ceramic fabrics and charred surface deposits, is a well-established analytical approach, contributing to a range of enquiries including manufacturing technology and typology, subsistence, cuisine, population movement, cultural interaction, and past environments (Roffet-Salque *et al.* 2017). Organic residues are characterised through the application of several analytical techniques, including gas chromatography-mass spectrometry (GC-MS), elemental analysis – isotope ratio mass spectrometry (EA-IRMS), and gas chromatography-combustion-isotope ratio mass spectrometry (GC-C-IRMS). These three techniques enable the identification of individual compounds absorbed in ceramics and preserved in surface deposits, bulk carbon and nitrogen isotope analysis of surface deposits, and compound-specific carbon isotope analysis of individual molecules absorbed in ceramic and surface deposit residues. These techniques have enabled the characterisation of a wide range of plant and animal products (Roffet-Salque *et al.* 2017). However, the detection of cereals has proven difficult, compared to animal and resinous plant products, as they contain a relatively low abundance of lipids and produce few diagnostic ‘biomarker’ compounds (Colonese *et al.* 2017; Hammann and Cramp 2018). Notable exceptions include two widely cultivated C_4 plants, maize (*Zea mays*) and broomcorn millet, that have been identified in archaeological ceramics, following analysis of authentic and experimental samples (Reber and Evershed 2004; Reber *et al.* 2004; Heron *et al.* 2016).

The effectiveness of isotopic techniques in the identification of *P. miliaceum* processing varies according to geography. Western Eurasia is dominated by C_3 plants, therefore, as a C_4 plant, broomcorn millet is isotopically distinct from most native carbon sources (Lightfoot *et al.* 2013; Wang *et al.* 2017). However, it is not possible to distinguish between isotopic enrichment derived from either broomcorn millet, other C_4 plants, marine products, or terrestrial animals that have consumed ^{13}C -enriched resources (Lee-Thorp 2008). Combined archaeobotanical and zooarchaeological analysis, in addition to isotopic analysis of human and other animal remains, may exclude potential sources of ^{13}C enrichment (e.g. Chakraborty *et al.* 2020), yet there are limitations to this approach. Therefore, in many scenarios, molecular characterisation by ORA is imperative.

Identification of *P. miliaceum* processing in archaeological ceramics is based on the detection of a discrete organic molecule, miliacin (Figure 1), together with evidence of ^{13}C enrichment (Heron *et al.* 2016). This approach is based on the



Miliacin

Figure 1. Structure of miliacin (olean-18-en-3β-ol).

detection of miliacin in sedimentary samples, particularly pollen cores and pit fills, applied to detect areas of past cultivation and storage of broomcorn millet (Jacob *et al.* 2008; Motuzaite-Matuzeviciute *et al.* 2016; Courel *et al.* 2017). Miliacin (olean-18-en-3β-ol) is a pentacyclic triterpene methyl ether (PTME) concentrated in the caryopses of *P. miliaceum*. Species in the genera *Panicum*, *Pennisetum*, *Paspalum*, *Digitaria*, *Chionocloa*, *Eragrostis*, *Glyceria*, and *Microstegium* are also reported to produce miliacin, in varying abundance, in addition to other PTMEs (Jacob *et al.* 2005; Bossard *et al.* 2013). Therefore, while miliacin dominates (c. 90%) the PTME fraction of broomcorn millet, when examining archaeological samples, it is important to note that the compound is not exclusive to this species (Bossard *et al.* 2013). In prehistoric European contexts, known sources of miliacin are limited to *Digitaria sanguinalis* (Ohmoto *et al.* 1970), a native species commonly known as hairy crabgrass, and *P. miliaceum* (Jacob *et al.* 2008; Bossard *et al.* 2013; Heron *et al.* 2016). There is little evidence to suggest that *D. sanguinalis* caryopses were ever gathered, processed, and consumed in large quantities, although it may feature as a weed. Therefore, in culinary contexts, it is most likely that *P. miliaceum* processing is the source of miliacin in pottery residues. Beyond Europe, a better understanding of the presence and abundance of miliacin in plants may be required. *P. miliaceum* is distinguishable from foxtail millet (*Setaria italica*), as it does not produce miliacin (Bossard *et al.* 2013). The expansion of foxtail millet cultivation into Europe is less clear, due to difficulties in distinguishing it from wild varieties and its presence as a potential weed in broomcorn millet agriculture, although some abundant finds have been reported (Reed and Drnić 2016).

Experiments have shown that miliacin is mobilised in water and absorbed into ceramic matrices when *P. miliaceum* caryopses are cooked in pottery vessels (Heron *et al.* 2016). However, the abundance of absorbed miliacin is a fraction of that present in caryopses (Bossard *et al.* 2013). Further experiments are necessary to explore the mobilisation, absorption, and retention of miliacin, during *P. miliaceum* processing, in addition to studies focused on the impact of concurrent and sequential processing with other foods, as has been investigated for other cereals (Hamman and Cramp 2018; Miller *et al.* 2020). The effect of different processing methods, *e.g.* boiling and fermentation, on the transfer and recovery of miliacin should also be investigated. Finally, the survival potential of miliacin in archaeological materials should be explored. In lieu of further experimental and archaeological research, it is not possible to compare the frequency and abundance of miliacin to other compounds identified in organic residues and it is premature to interpret an absence of miliacin as an absence of broomcorn millet processing.

Identification of broomcorn millet by organic residue analysis of pottery vessels: A brief review

Heron *et al.* (2016) investigated millet processing in Bronze Age and Iron Age ceramics from Bruszczevo (Poland) and Bronze Age ceramics from Majeon-ri (Korean Peninsula). At Bruszczevo, bulk isotope analysis of Early Bronze Age (EBA) and Late Bronze Age/Early Iron Age (LBA/EIA) charred surface deposits identified ^{13}C enrichment, in the latter period, indicative of C_4 plant input. Comparable ^{13}C enrichment was observed in compound-specific carbon isotope analysis of a single LBA/EIA sample examined by GC-C-IRMS. Furthermore, the lipid profile of this residue was consistent with a plant origin and a trace of miliacin was observed. In combination, these data securely identify *P. miliaceum* processing in pottery vessels at Bruszczevo and support the chronology of its introduction and adoption at the site (Kroll 2010). Miliacin was also identified in absorbed ceramic residues from Majeon-ri, although no surface deposits were available for analysis. The results indicate that *P. miliaceum* caryopses were processed in ceramic vessels at both sites, forming charred surface deposits at Bruszczevo and absorbed ceramic residues at Majeon-ri. Further analysis is currently underway at Bruszczevo to investigate absorbed ceramic residues. The presence of broomcorn millet residues confirms direct consumption by humans, opening new avenues into detecting culinary practices incorporating this cereal. Furthermore, this study demonstrates the ability of ORA methods in understanding the prevalence of *P. miliaceum* processing and temporally tracking its adoption in Europe.

The initial report by Heron *et al.* (2016) has prompted several subsequent investigations of potential millet residues, albeit applying slightly different methodologies. Common to most of these investigations is the use of GC-MS to detect miliacin in absorbed ceramic residues (Fig. 2). Bulk carbon and nitrogen isotope analysis is infrequently undertaken due to the relative scarcity of charred surface deposits. Additionally, compound-specific carbon isotope analysis has yet to be applied routinely to detect C_4 plant influence.

Isaksson and Nilsson (2018) report GC-MS analysis of 12 sherds from the LBA cemetery at Maciejowice (Poland). Miliacin was identified in four samples and is attributed to *P. miliaceum* processing. This interpretation is supported by the local chronology of this species (Kapcia and Mueller-Bieniek 2019; Filipović *et al.* 2020) and scorch marks on vessel exteriors, that indicate cooking. The presence of broomcorn millet residues in burial contexts potentially demonstrates the inclusion of this species in cultural activities as either a staple food or product with greater significance. Further analysis of residues, from cemeteries and settlements, in the surrounding region may elucidate the role of this species.

Ganzarolli *et al.* (2018) identified miliacin in 26 of 45 Late Antique and Early Medieval Age (6th-10th century) ceramic sherds from Padova (Italy). Miliacin was either the only, or the dominant, PTME observed in these 26 samples and its abundance was low, relative to other lipid constituents, in all but one sample. This is typical of residues produced during *P. miliaceum* processing (Figure 3 shows an example from Bruszczevo). However, one sample contained a higher than usual abundance of miliacin, potentially demonstrating either different processing or preservation conditions. This enabled determination of the compound-specific carbon isotope value of the compound (-20.5‰), confirming a ^{13}C -enriched, *i.e.* C_4 plant, origin.

Kučera *et al.* (2019) detected miliacin in the soil content of a complete vessel, placed in a Corded Ware culture cremation grave (ca. 2707-2571 cal BCE), from Držovice (Czech Republic). Miliacin was not observed in either the ceramic fabric of this vessel or the soil contents of nearby vessels. Therefore, Kučera *et al.* (2019) suggest that broomcorn millet was stored in the jug and deposited in the grave,



Figure 2. European sites wherein miliacin has been identified in ceramics (filled circles) and soils/sediments (open circles). 1. Vix-Mont Lassois (Rageot *et al.* 2019a) 2. Lake Paladru (Simonneau *et al.* 2013) 3. Lake le Bourget (Jacob *et al.* 2008) 4. Obernai (Courel *et al.* 2017) 5. Heuneburg (Rageot *et al.* 2019b) 6. Padova (Ganzarolli *et al.* 2018) 7. Bruszczewo (Heron *et al.* 2016) 8. Držovice (Kučera *et al.* 2019) 9. Maciejowice (Isaksson and Nilsson 2018) 10. Zanolovskoe (Motuzaite-Matuzeviciute *et al.* 2016).

perhaps as an offering. However, this interpretation is not supported by the current chronology of *P. miliaceum* translocation across Eurasia (Frachetti *et al.* 2010; Spengler *et al.* 2018; Hermes *et al.* 2019; Leipe *et al.* 2019). Indeed, Direct AMS-dating of over 100 European millet caryopses has produced no evidence for the presence of this species in Europe prior to the 2nd millennium BCE (Motuzaite-Matuzeviciute *et al.* 2013; Filipović *et al.* 2020). Therefore, either the source or date of the miliacin observed in this study requires reconsideration. The analysis of soil fills of pottery vessels is unlikely to be a reliable approach without supporting evidence from either charred surface deposit/absorbed ceramic residues or archaeobotanical analysis.

Rageot *et al.* (2019a, b) report the observation of miliacin in their analysis of over 200 vessels from the Early Celtic sites at Vix-Mont Lassois (France) and Heuneburg (Germany) respectively. Miliacin was not observed in any imported wares, likely suggesting that broomcorn millet was neither imported nor processed in these vessels. Miliacin is observed in locally produced vessels, yet it is less prevalent in high-status areas and in later periods, at both sites, where Mediterranean influence is more pronounced. Furthermore, it is completely absent in late occupation high-status areas at Heuneburg (Rageot *et al.* 2019b). The data indicates a shift in the perception of *P. miliaceum* by the Celtic elite, at both sites, but persistence of subsistence practices, that include broomcorn millet, among lower status individuals (Rageot *et al.* 2019a, b). The observed decline of *P. miliaceum* among elite groups correlates with a higher abundance of Mediterranean vessels, associated with wine consumption (Rageot *et al.* 2019a). Miliacin is observed in both domestic and fine wares at Vix-Mont Lassois and Heuneburg, indicating varied processing and consumption contexts (Rageot *et al.* 2019a, b). However, at Vix-Mont Lassois, miliacin is most commonly observed in drinking type vessels, potentially indicating the consumption of a millet-based beverage. The co-occurrence of miliacin and beeswax in eight fine handmade bowls, in addition to the presence of bacterial hopanoids in half of these vessels, led the authors to propose the consumption of a fermented beverage (beer) containing *P. miliaceum* (Rageot *et al.* 2019a). Distinguishing cooked and fermented millet products is highly desirable. However, further experimental and archaeological research is necessary to prove such distinction is possible.




Figure 3. Partial total ion chromatogram of a charred 'foodcrust' from Bruszczewo that contains a low abundance of miliacin, relative to dominant fatty acids, $C_{16:0}$ (palmitic) and $C_{18:0}$ (stearic). ☆ Internal C34 and C36 alkane standards.

Junno *et al.* (2020) present ORA data, from Hamanaka 2 (Rebun Island, Japan), that is characteristic of marine product processing, including ^{13}C enriched palmitic ($C_{16:0}$) and stearic ($C_{18:0}$) acids and abundant aquatic biomarkers. The ^{13}C enrichment from marine products is likely to mask a contribution of C_4 plant lipids to residues, therefore, this study is reliant on miliacin to detect *P. miliaceum* processing. A single sample analysed contained miliacin and was interpreted as evidence for the earliest identification of *P. miliaceum* in the region. Cultivation of this species is known among Okhotsk communities on the island of Hokkaido, yet there is presently no archaeobotanical evidence for its presence in the region of Hamanaka 2 (Leipe *et al.* 2017). This data may demonstrate early processing of *P. miliaceum*, and the integration of Hamanaka 2 into long-distance exchange networks (Junno *et al.* 2020). However, to confirm this theory, greater sampling of local flora and vessels from Hamanaka 2 and other Okhotsk sites is necessary to track and trace the appearance of miliacin and *P. miliaceum* throughout the region.

Chakraborty *et al.* (2020) present absorbed ceramic residue data from Kotada Bhadli (Gujarat, India). Samples demonstrate varying degrees of lipid ^{13}C enrichment, that may derive from the processing of either C_4 plants or C_4 -fed animal products. Miliacin is not observed in any of the residues analysed, despite *P. miliaceum* cultivation in the region at this time. Therefore, the authors contend that broomcorn millet was not directly processed in the vessels (Chakraborty *et al.* 2020). However, it is debatable whether an absence of miliacin in pottery residues can be used to rule out the processing of *P. miliaceum*, as aspects of miliacin release, transfer, absorption, accumulation, and preservation are uncertain.

Next steps in the study of millet processing in archaeological ceramics

The detection of residues from processing *P. miliaceum*, in association with material culture, provides a rare opportunity to investigate the introduction, adoption, uses and role of a single crop. The criteria underpinning the detection of broomcorn millet processing, proposed by Heron *et al.* (2016) at Bruszczewo and Majeon-ri are as follows:

- Elevated bulk carbon isotope values in charred surface deposits (foodcrusts), consistent with a C₄ plant influence.
- Identification of miliacin (olean-18-en-3 β -ol methyl ether) in either charred surface deposits or ceramic-absorbed residues.
- Elevated carbon isotope values in *n*-alkanoic (fatty) acids and miliacin, if available in sufficient quantities, consistent with C₄ plant influence.

These three criteria have not been met in most applications reviewed above since the preservation of charred surface deposits is rare in many archaeological contexts, hence bulk isotope analysis is not possible. Further complications arise and may underestimate the occurrence of broomcorn millet in pottery vessels. Firstly, miliacin may be present at levels that are below the detection limit of GC-MS, especially in cases where broomcorn millet was processed as a minor constituent in a vessel. Furthermore, our current understanding of the uptake and preservation of this compound, in archaeological ceramics, is limited. Secondly, the mixing of isotopically diverse foods may either mask or diminish the C₄ isotope signal imparted during the processing of *P. miliaceum*, complicating the interpretation of bulk and compound-specific isotope data.

Further experimental research is necessary to investigate and understand these issues. The transfer of miliacin, from seed to archaeological material, under different processing conditions, *e.g.* boiling and fermentation, in addition to its preservation, during multiple cooking events and deposition under different burial conditions, require detailed experimental studies. Furthermore, while the dominance and overall abundance of miliacin in broomcorn millet caryopses provides a degree of confidence in assigning the compound to *P. miliaceum*, reliance on the detection of miliacin is likely to remain the subject of debate, due to its occurrence in other plant species. Therefore, extensive sampling of reference materials is also strongly advised. Finally, an examination of the impact that *P. miliaceum* processing has on carbon isotope values of surface deposits and absorbed ceramic residues, by EA-IRMS and GC-C-IRMS, would help understand isotopically complex datasets.

The detection of miliacin may prove useful in the identification of *P. miliaceum* processing in other materials, *e.g.* grinding stones. The compound has also been detected in coprolites, providing evidence of direct consumption (Zhang *et al.* 2020). Therefore, exciting opportunities exist to expand the detection of this cereal to new fields.

There is much to be gained by pursuing the study of culinary activities, from single cooking events to entire cuisines. Large-scale studies, assessing substantial numbers of specific vessel types, are necessary to examine the contribution of *P. miliaceum* to food processing and consumption patterns and move our understanding beyond simple presence/absence observations of the crop in antiquity.

Acknowledgements

We thank Antony Simpson (British Museum) for producing the figures. We also thank the organisers of the conference for providing the opportunity to present our research and contribute to this publication. Thanks also to the constructive comments of the referees. This work is supported by the Wellcome Trust (097365/Z/11/Z).

References

- An, T., Pashkevich, G., Jones, M., 2019. Re-examining millet impressions in Usatovo clay materials from NW Black Sea region, Ukraine. *Archaeological and Anthropological Sciences*, 11 (7), 3201-3211. <https://doi.org/10.1007/s12520-018-0718-3>
- Boivin, N., Fuller, D.Q., Crowther, A., 2012. Old World globalization and the Columbian exchange: comparison and contrast. *World Archaeology*, 44 (3), 452-469. <https://doi.org/10.1080/00438243.2012.729404>
- Bossard, N., Jacob, J., Le Milbeau, C., Sauze, J., Terwilliger, V., Poissonnier, B., Vergès, E., 2013. Distribution of miliacin (olean-18-en-3 β -ol methyl ether) and related compounds in broomcorn millet (*Panicum miliaceum*) and other reputed sources: implications for the use of sedimentary miliacin as a tracer of millet. *Organic Geochemistry*, 63, 48-55. <https://doi.org/10.1016/j.orggeochem.2013.07.012>
- Chakraborty, K.S., Slater, G.F., Miller, H.M.L., Shirvalkar, P., Rawat, Y., 2020. Compound specific isotope analysis of lipid residues provides the earliest direct evidence of dairy product processing in South Asia. *Scientific Reports*, 10 (1), 16095. <https://doi.org/10.1038/s41598-020-72963-y>
- Colanese, A.C., Hendy, J., Lucquin, A., Speller, C.F., Collins, M.J., Carrer, F., Gubler, R., Kühn, M., Fischer, R., Craig, O.E., 2017. New criteria for the molecular identification of cereal grains associated with archaeological artefacts. *Scientific Reports*, 7 (1), 6633. <https://doi.org/10.1038/s41598-017-06390-x>
- Courel, B., Schaeffer, P., Adam, P., Motsch, E., Ebert, Q., Moser, E., Féliu, C., Bernasconi, S.M., Hajdas, I., Ertlen, D., Schwartz, D., 2017. Molecular, isotopic and radiocarbon evidence for broomcorn millet cropping in northeast France since the Bronze Age. *Organic Geochemistry*, 110, 13-24. <https://doi.org/10.1016/j.orggeochem.2017.03.002>
- Dal Corso, M., Nicosia, C., Balista, C., Cupitò, M., Dalla Longa, E., Leonardi, G., Kirleis, W., 2017. Bronze Age crop processing evidence in the phytolith assemblages from the ditch and fen around Fondo Paviani, northern Italy. *Vegetation History and Archaeobotany*, 26 (1), 5-24. <https://doi.org/10.1007/s00334-016-0573-z>
- Filipović, D., Meadows, J., Corso, M.D., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Kloof, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A.M., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H.-P., Stobbe, A., Tolar, T., Wasylkowa, K., Wiethold, J., Zerl, T., 2020. New AMS ^{14}C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Frachetti, M.D., Spengler, R.N., Fritz, G.J., Mar'yashev, A.N., 2010. Earliest direct evidence for broomcorn millet and wheat in the central Eurasian steppe region. *Antiquity*, 84 (326), 993-1010. <https://doi.org/10.1017/S0003598X0006703X>
- Ganzarolli, G., Alexander, M., Arnau, A.C., Craig, O.E., 2018. Direct evidence from lipid residue analysis for the routine consumption of millet in early medieval Italy. *Journal of Archaeological Science*, 96, 124-130.
- Gong, Y., Yang, Y., Ferguson, D.K., Tao, D., Li, W., Wang, C., Lü, E. and Jiang, H., 2011. Investigation of ancient noodles, cakes, and millet at the Subeixi Site, Xinjiang, China. *Journal of Archaeological Science* 38, 470-479. <https://doi.org/10.1016/j.jas.2010.10.006>

- Hammann, S. and Cramp, L.J.E., 2018. Towards the detection of dietary cereal processing through absorbed lipid biomarkers in archaeological pottery. *Journal of Archaeological Science*, 93, 74-81. <https://doi.org/10.1016/j.jas.2018.02.017>
- Hastorf, C.A., 2017. *The Social Archaeology of Food: Thinking about Eating from Prehistory to the Present*. Cambridge: Cambridge University Press.
- Hermes, T.R., Frachetti, M.D., Doumani Dupuy, P.N., Mar'yashev, A., Nebel, A., Makarewicz, C.A., 2019. Early integration of pastoralism and millet cultivation in Bronze Age Eurasia. *Proceedings of the Royal Society B*, 286 (1910), 20191273. <https://doi.org/10.1098/rspb.2019.1273>
- Heron, C., Shoda, S., Breu Barcons, A., Czebreszuk, J., Eley, Y., Gorton, M., Kirleis, W., Kneisel, J., Lucquin, A., Müller, J., Nishida, Y., Son, J.-H., Craig, O.E., 2016. First molecular and isotopic evidence of millet processing in prehistoric pottery vessels. *Scientific Reports*, 6, 38767. <https://doi.org/10.1038/srep38767>
- Isaksson, S., and Nilsson, C., 2018. Analyses of lipid residues in Lusatian pottery from Maciejowice, Poland. *Uppdragsrapport* [online], 327. <http://urn.kb.se/resolve?urn=urn:nbn:se:su:diva-158652>
- Jacob, J., Disnar, J.-R., Boussafir, M., Spadano Albuquerque, A.L., Sifeddine, A., Turcq, B., 2005. Pentacyclic triterpene methyl ethers in recent lacustrine sediments (Lagoa do Caçó, Brazil). *Organic Geochemistry*, 36 (3), 449-461. <https://doi.org/10.1016/j.orggeochem.2004.09.005>
- Jacob, J., Disnar, J.-R., Arnaud, F., Chapron, E., Debret, M., Lallier-Vergès, E., Desmet, M., Revel-Rolland, M., 2008. Millet cultivation history in the French Alps as evidenced by a sedimentary molecule. *Journal of Archaeological Science*, 35, 814-820. <https://doi.org/10.1016/j.jas.2007.06.006>
- Jones, M., Hunt, H., Lightfoot, E., Lister, D., Liu, X., Motuzaite-Matuzeviciute, G., 2011. Food globalization in prehistory. *World Archaeology*, 43 (4), 665-675. <https://doi.org/10.1080/00438243.2011.624764>
- Junno, A., Isaksson, S., Hirasawa, Y., Kato, H., Jordan, P.D., 2020. Evidence of increasing functional differentiation in pottery use among late Holocene maritime foragers in northern Japan. *Archaeological Research in Asia*, 22, 100194. <https://doi.org/10.1016/j.ara.2020.100194>
- Kapcia, M. and Mueller-Bieniek, A., 2019. An insight into Bronze Age subsistence strategy in forested Carpathian foothills, based on plant macroremains. *Archaeological and Anthropological Sciences*, 11(6), 2879-2895. <https://doi.org/10.1007/s12520-018-0720-9>
- Kroll, H., 2010. Die Archäobotanik von Bruszczewo-Darstellung und Interpretation der Ergebnisse. In: J. Müller, J. Czebreszuk, J. Kneisel, eds., *Bruszczewo II: Ausgrabungen und Forschungen einer prähistorischen Siedlungskammer Großpolens*. Studien zur Archäologie in Ostmitteleuropa, 6 (1). Bonn: Dr Rudolf Habelt GmbH, 250-287.
- Kučera, L., Peška, J., Fojtík, P., Barták, P., Kučerová, P., Pavelka, J., Komárková, V., Beneš, J., Polcerová, L., Králík, M., Bednář, P., 2019. First direct evidence of broomcorn millet (*Panicum miliaceum*) in central Europe. *Archaeological and Anthropological Sciences*, 11 (8), 4221-4227. <https://doi.org/10.1007/s12520-019-00798-4>
- Lee-Thorp, J.A., 2008. On isotopes and old bones. *Archaeometry*, 50(6), 925-950. <https://doi.org/10.1111/j.1475-4754.2008.00441.x>
- Leipe, C., Sergusheva, E.A., Müller, S., Spengler, R.N., 3rd, Goslar, T., Kato, H., Wagner, M., Weber, A. W., Tarasov, P.E., 2017. Barley (*Hordeum vulgare*) in the Okhotsk culture (5th-10th century AD) of northern Japan and the role of cultivated plants in hunter-gatherer economies. *PLoS ONE*, 12 (3), e0174397. <https://doi.org/10.1371/journal.pone.0174397>
- Leipe, C., Long, T., Sergusheva, E.A., Wagner, M., Tarasov, P.E., 2019. Discontinuous spread of millet agriculture in eastern Asia and prehistoric population dynamics. *Science Advances*, 5 (9), eaax6225. <https://doi.org/10.1126/sciadv.aax6225>

- Lightfoot, E., Liu, X., Jones, M.K., 2013. Why move starchy cereals? A review of the isotopic evidence for prehistoric millet consumption across Eurasia. *World Archaeology*, 45 (4), 574-623. <https://doi.org/10.1080/00438243.2013.852070>
- Liu, L., Wang, J., Liu, H., 2020. The brewing function of the first amphorae in the Neolithic Yangshao culture, north China. *Archaeological and Anthropological Sciences*, 12, 118. <https://doi.org/10.1007/s12520-020-01069-3>
- Long, T., Leipe, C., Jin, G., Wagner, M., Guo, R., Schröder, O., Tarasov, P.E., 2018. The early history of wheat in China from 14C dating and Bayesian chronological modelling. *Nature Plants*, 4 (5), 272-279. <https://doi.org/10.1038/s41477-018-0141-x>
- Lu, H., Yang, X., Ye, M., Liu, K.-B., Xia, Z., Ren, X., Cai, L., Wu, N., Liu, T.-S., 2005. Millet noodles in Late Neolithic China. *Nature* 437, 967-968. <https://doi.org/10.1038/437967a>
- Miller, N.F., 2015. Rainfall seasonality and the spread of millet cultivation in Eurasia. *Iranian Journal of Archaeological Studies*, 5 (1), 1-10. <https://doi.org/10.22111/IJAS.2015.3015>
- Miller, N.F., Spengler, R.N., Frachetti, M., 2016. Millet cultivation across Eurasia: origins, spread, and the influence of seasonal climate. *The Holocene*, 26(10), 1566-1575. <https://doi.org/10.1177/0959683616641742>
- Miller, M.J., Whelton, H.L., Swift, J.A., Maline, S., Hammann, S., Cramp, L.J., McCleary, A., Taylor, G., Vacca, K., Becks, F., Evershed, R.P., 2020. Interpreting ancient food practices: stable isotope and molecular analyses of visible and absorbed residues from a year-long cooking experiment. *Scientific Reports*, 10, 13704. <https://doi.org/10.1038/s41598-020-70109-8>
- Miyata, Y., Kojima, T., Matsutani, A., Onbe, S., Nishimoto, T., 2007. Oldest common millet from western Japan: carbonized material adhering to pottery from the Ryugasaki A site, Shiga prefecture. *Bulletin of the National Museum of Japanese History*, 137, 255-265.
- Motuzaitė-Matuzevičiūtė, G., Staff, R.A., Hunt, H.V., Liu, X., Jones, M.K., 2013. The early chronology of broomcorn millet (*Panicum miliaceum*) in Europe. *Antiquity*, 87 (338), 1073-1085. <https://doi.org/10.1017/S0003598X00049875>
- Motuzaitė-Matuzevičiūtė, G., Jacob, J., Telizhenko, S., Jones, M.K., 2016. Miliacin in palaeosols from an Early Iron Age in Ukraine reveal in situ cultivation of broomcorn millet. *Archaeological and Anthropological Sciences*, 8(1), 43-50. <https://doi.org/10.1007/s12520-013-0142-7>
- Mueller-Bieniek, A., Nowak, M., Styring, A., Lityńska-Zajac, M., Moskal-del Hoyo, M., Sojka, A., Paszko, B., Tunia, K., Bogaard, A., 2019. Spatial and temporal patterns in Neolithic and Bronze Age agriculture in Poland based on the stable carbon and nitrogen isotopic composition of cereal grains. *Journal of Archaeological Science: Reports*, 27, 101993. <https://doi.org/10.1016/j.jasrep.2019.101993>
- Ohmoto, T., Ikuse, M., Natori, S., 1970. Triterpenoids of the Gramineae. *Phytochemistry*, 9 (10), 2137-2148. [https://doi.org/10.1016/S0031-9422\(00\)85379-0](https://doi.org/10.1016/S0031-9422(00)85379-0)
- Rageot, M., Mötsch, A., Schorer, B., Bardel, D., Winkler, A., Sacchetti, F., Chaume, B., Della Casa, P., Buckley, S., Cafisso, S., Fries-Knoblach, J., Krausse, D., Hoppe, T., Stockhammer, P., Spiteri, C., 2019a. New insights into Early Celtic consumption practices: organic residue analyses of local and imported pottery from Vix-Mont Lassois. *PLoS ONE*, 14 (6), e0218001. <https://doi.org/10.1371/journal.pone.0218001>
- Rageot, M., Mötsch, A., Schorer, B., Gutekunst, A., Patrizi, G., Zerrer, M., Cafisso, S., Fries-Knoblach, J., Hansen, L., Tarpini, R., Krausse, D., Hoppe, T., Stockhammer, P.W., Spiteri, C., 2019b. The dynamics of Early Celtic consumption practices: a case study of the pottery from the Heuneburg. *PLoS ONE*, 14 (10), e0222991. <https://doi.org/10.1371/journal.pone.0222991>
- Reber, E.A. and Evershed, R.P., 2004. How did Mississippians prepare maize? The application of compound-specific carbon isotope analysis to absorbed pottery residues from several Mississippi Valley sites. *Archaeometry*, 46 (1), 19-33. <https://doi.org/10.1111/j.1475-4754.2004.00141.x>

- Reber, E.A., Dudd, S.N., van der Merwe, N.J., Evershed, R.P., 2004. Direct detection of maize in pottery residues via compound specific stable carbon isotope analysis. *Antiquity*, 78 (301), 682-691. <https://doi.org/10.1017/S0003598X00113316>
- Reed, K. and Drnić, I., 2016. Iron Age diet at Sisak, Croatia: archaeobotanical evidence of foxtail millet (*Setaria italica* [L.] P. Beauv.). *Oxford Journal of Archaeology*, 35 (4), 359-368. <https://doi.org/10.1111/ojoa.12101>
- Roffet-Salque, M., Dunne, J., Altoft, D.T., Casanova, E., Cramp, L.J.E., Smyth, J., Whelton, H.L., Evershed, R.P., 2017. From the inside out: upscaling organic residue analyses of archaeological ceramics. *Journal of Archaeological Science: Reports*, 16, 627-640. <https://doi.org/10.1016/j.jasrep.2016.04.005>
- Simonneau, A., Doyen, E., Chapron, E., Millet, L., Vannière, B., Di Giovanni, C., Bossard, N., Tachikawa, K., Bard, E., Albéric, P., Desmet, M., 2013. Holocene land-use evolution and associated soil erosion in the French Prealps inferred from Lake Paladru sediments and archaeological evidences. *Journal of Archaeological Science*, 40, 1636-1645. <https://doi.org/10.1016/j.jas.2012.12.002>
- Spengler, R.N., de Nigris, I., Cerasetti, B., Carra, M., Rouse, L.M., 2018. The breadth of dietary economy in Bronze Age central Asia: case study from Adjı Kui 1 in the Murghab region of Turkmenistan. *Journal of Archaeological Science: Reports*, 22, 372-381. <https://doi.org/10.1016/j.jasrep.2016.03.029>
- Stevens, C.J., Shelach-Lavi, G., Zhang, H., Teng, M., Fuller, G.Q., 2020. A model for the domestication of *Panicum miliaceum* (common, proso or broomcorn millet) in China. *Vegetation History and Archaeobotany*, 30, 21-33. <https://doi.org/10.1007/s00334-020-00804-z>
- Valamoti, S.M., 2016. Millet, the late comer: on the tracks of *Panicum miliaceum* in prehistoric Greece. *Archaeological and Anthropological Sciences*, 8, 51-63. <https://doi.org/10.1007/s12520-013-0152-5>
- Wang, T., Wei, D., Chang, X., Yu, Z., Zhang, X., Wang, C., Hu, Y., Fuller, B.T., 2017. Tian-shanbeilu and the Isotopic Millet Road: reviewing the Late Neolithic/Bronze Age radiation of human millet consumption from north China to Europe. *National Science Review*, 6 (5), 1024-1039. <https://doi.org/10.1093/nsr/nwx015>
- Yang, X., Wan, Z., Perry, L., Lu, H., Wang, Q., Zhao, C., Li, J., Xie, F., Yu, J., Cui, T., Wang, T., Li, M., Ge, Q., 2012a. Early millet use in northern China. *Proceedings of the National Academy of Sciences of the United States of America*, 109 (10), 3726-3730. <https://doi.org/10.1073/pnas.1115430109>
- Yang, X., Zhang, J., Perry, L., Ma, Z., Wan, Z., Li, M., Diao, X., Lu, H., 2012b. From the modern to the archaeological: starch grains from millets and their wild relatives in China. *Journal of Archaeological Science*, 39, 247-254. <https://doi.org/10.1016/j.jas.2011.09.001>
- Zhang, Y., Zhang, D., Yang, Y., Wu, X., 2020. Pollen and lipid analysis of coprolites from Yuhuicun and Houtieying, China: Implications for human habitats and diets. *Journal of Archaeological Science: Reports*, 29, 102135. <https://doi.org/10.1016/j.jasrep.2019.102135>
- Zhao, Z.J., 2011. New archaeobotanical data for the study of the origins of agriculture in China. *Current Anthropology*, 52 (Suppl. 4), S295-S306. <https://doi.org/10.1086/659308>

Tracing millet through biomarker analysis in archaeological sites in alluvial plains: The first miliacin data from the northern Italian Bronze Age

Marta Dal Corso, Marco Zanon, Carl Heron, Mauro Rottoli, Michele Cupitò, Elisa Dalla Longa, Wiebke Kirleis

Abstract

Broomcorn millet has been attested by the biomarker miliacin in the archaeological sediments at the *terramara* site of Fondo Paviani (ca. 1350-1100 BCE), in the Po River valley, northern Italy. This molecule is typical of *Panicum miliaceum* and it was absent in the other modern panicoid grasses analysed for reference in this study. We present the results of investigation of on-site sediments and, together with the available archaeobotanical and isotopic records, incorporate them into the picture of millet consumption and cultivation during the Late Bronze Age in the Po valley.

Introduction

Background on millet and miliacin studies

In the present study, we explore the potential of biomarker analysis in archaeological stratigraphic features. We focus on the biomarker miliacin, a pentacyclic triterpene methyl ether (PTME), currently considered a promising indicator to complement studies on crop diversity in archaeological contexts. The interest in miliacin lies in its ability to trace the presence of a specific crop, namely broomcorn millet (*Panicum miliaceum*), even in the absence of unambiguous botanical evidence (Dubois and Jacob 2016). This method has recently received increased attention, corresponding to that given to the early adoption of millet outside its region of domestication in northern China (Liu *et al.* 2009, 2012), which is indicative of an expansion in contacts, materials and know-how. This topic has gained the attention of many scholars, as testified by the present volume and by the rich literature derived from more than

a decade of archaeobotanical research in central Asia and Europe (e.g. Hunt *et al.* 2008; Motuzaite-Matuzeviciute *et al.* 2013; Miller *et al.* 2016; Filipović *et al.* 2020).

Broomcorn millet as a crop did not reach Europe together with the ‘Neolithic package of founder crops’ composed of emmer, einkorn, wheat and barley (e.g. Rottoli and Castiglioni 2009a; Zohary and Hopf 2012; Rottoli and Pessina 2016; Kirleis 2019; Liu *et al.* 2019). In contrast to these large-grained cereals, which were domesticated in the Fertile Crescent, broomcorn millet belongs to a group of small-grained cereals, usually referred to as millets, which includes drought-resistant species with short growing cycles. In Europe, the first evidence for millets from prehistoric archaeological contexts concerns *Panicum miliaceum* and *Setaria italica* (foxtail millet). A recent dating programme of charred broomcorn millet caryopses from European prehistoric sites established that it diffused as a crop in Europe during the Bronze Age, starting from the 16th century BCE (Filipović *et al.* 2020). Currently, it seems that northern Italy, together with Ukraine, was one of the first regions of Europe where millet cultivation was adopted (Dal Corso *et al.* 2022).

The adoption of a new crop for alimentary purposes has multiple and possibly overlapping cultural implications, ranging from forward planning (i.e. increasing diversity of cultivars as a means to overcome possible crop failures) to short-term human adaptability (i.e. the need to react to ongoing adverse conditions). In addition, millet, as a fast-growing crop, could indicate the adoption of more mobile settlement strategies, or contacts with more mobile human communities, as well as the use of only periodically available land (Motuzaite-Matuzeviciute *et al.* 2013, 2016; Ventresca Miller *et al.* 2020; Dal Corso *et al.* 2022). The study of early millet adoption, therefore, has multifaceted implications for archaeological research, especially as there are multiple ways to trace millet in archaeological contexts, as described below.

Millet consumption can be traced through carbon and nitrogen stable isotope analysis (e.g. Tafuri *et al.* 2009, 2018; Lightfoot *et al.* 2013) conducted on human and animal bones, and they can reveal consumption of millet as staple food or fodder. This method is based on the fact that broomcorn millet uses the C4 photosynthetic pathway and is the first such crop in Europe. Thus, the method is particularly appropriate for studies of prehistoric societies in temperate environments poor in wild panicoids and other wild or cultivated C4 plant species, i.e. where the C4 plant signal identified isotopically can be attributed to millet as a crop. Plant macro-remain analysis is the usual way to detect crops in archaeology. It is made difficult by the fact that millet grains are small and require significant effort to be recovered. In older investigations, the absence of millet remains may be more a reflection of insufficient sampling and processing strategies (e.g. too-large mesh sizes, no flotation) rather than of their genuine absence. Furthermore, unless the grains are fused together (in ‘clumps’), their small size makes millet grains more prone to contaminating underlying, unrelated archaeological features, requiring radiocarbon dating to detect temporally misattributed specimens. In the case of pollen analysis, the morphology of millet pollen grains is usually considered insufficiently diagnostic to avoid confusion with wild, non-economic grass species (Beug 2004). Attempts to detect millet cultivation by means of pollen analysis therefore remain limited (e.g. Cremaschi *et al.* 2016). Phytolith analysis represents a promising research direction in the distinction of *Panicum miliaceum* (Lu *et al.* 2009; Out and Madella 2016), yet it remains a relatively niche research field in temperate environments (e.g. in Italy; Dal Corso *et al.* 2017), which would benefit from the analysis of more wild plants. Given the limitations of the micro- and macro-botanical evidence, any independent data source represents a useful addition to confirm the presence of early millet adoption and cultivation.

In the present paper, our research question is whether it is feasible to use sedimentary miliacin as a molecular tracer of *Panicum miliaceum* at Bronze Age

sites of the Po River valley in northern Italy. In this way, we aim to contribute to the reconstruction of the history of this crop in a very dynamic period and region of Europe.

Miliacin in modern plants and archaeological studies

Miliacin (olean-18-en-3 β -ol methyl ether) is a PTME and is found in high concentrations in the seeds of *Panicum miliaceum*. The value of miliacin as a marker for the presence of millet depends primarily on its unequivocal association with millets in a given study area. Bossard *et al.* (2013) investigated the miliacin content of *Setaria italica* and of multiple *Panicum* and *Pennisetum* species. Miliacin was detected in all *Panicum* species analysed, while it was sparsely present in *Pennisetum*. It was not detected in *Setaria italica*, in contrast with earlier, independent findings (Lu *et al.* 2009). Bossard *et al.* (2013) also tested the presence of miliacin in the green mould *Chaetomium olivaceum* reported by Smetanina *et al.* (2001) and found none. Few other plants are currently known to be a source of miliacin. A possibly non-exhaustive list (sourced from Ohmoto *et al.* 1970; Connor and Purdie 1976, 1981; Bossard *et al.* 2013) includes *Eragrostis ferruginea* (Thunb.) P. Beauv., *Glyceria acutiflora* Torr and different *Chionochloa* species, among others. Most of the species mentioned so far are not native to Europe (Pyankov *et al.* 2010) and they are therefore not relevant to discussing agricultural dynamics in northern Italy during the Bronze Age. In this context, *P. miliaceum* is the most abundant cereal among the known miliacin-producing taxa. Two aspects support the use of sedimentary records of miliacin in some contexts to track past millet agricultural use: the high concentration of miliacin in the grain of *P. miliaceum* relative to other PTMEs and to other grasses, and the high biomass that cultivated broomcorn millet possesses relative to other potential plant sources.

The same inferred relationship between *P. miliaceum* and miliacin has been used as a basis to detect millet cultivation and processing at multiple European sites from the Bronze Age onwards (Jacob *et al.* 2008; Heron *et al.* 2016; Motuzaite-Matuzeviciute *et al.* 2016; Courel *et al.* 2017). Based on previous studies, where sedimentary archives of miliacin have been found in conjunction with a high concentration of other PTMEs (e.g. at a Brazilian lake – Jacob *et al.* 2005), the compound cannot be used as a biomarker for *P. miliaceum*. In contrast, miliacin was the sole detectable PTME in the sediments from lake Le Bourget in France (Jacob *et al.* 2008, 2009), lakes Ledro and Paladru in Italy (Simonneau *et al.* 2013), and in Ukrainian palaeosols thought to have been cultivated for *P. miliaceum* (Motuzaite-Matuzeviciute *et al.* 2013). In conjunction with other evidence, the presence of miliacin in these cases was interpreted as an indicator of the cultivation of *P. miliaceum* in the catchment area.

In this study, we analysed modern grains of *P. miliaceum* as well as grains from other panicoid grass species possibly occurring in Italy. We investigated their miliacin content in order to compare the results with sedimentary miliacin content from the archaeological contexts described here.

Archaeological setting

The archaeological sediment analysed in this study originates from different contexts. First, we sampled two Bronze Age sites. The first site is the pile-dwelling of Bande di Cavriana (Mantua), located within a now-drained lake south of Lake Garda in northern Italy. Sediment samples from different cores from Cavriana (CAV 4-11-12-15, described in Zanon *et al.* 2019) were processed with a 500 μ m sieve, yielding no evidence of millet caryopses. Despite the absence of millet macro-remains, the finds of millet at other northern Italian prehistoric sites (Rottoli and Castiglioni 2009a, 2009b) and the methodological approach used at lake Le Bourget

(Jacob *et al.* 2008) suggested that testing the samples for miliacin could provide information on the possible millet cultivation close to Cavriana.

The samples for miliacin analysis came from an off-site sediment core (CAV 4) taken in the former lake basin and used for palaeoenvironmental investigation (Zanon *et al.* 2019). In the core, deposits were present that partly dated to the Bronze Age occupation related to the pile-dwelling (Zanon *et al.* 2019). Miliacin was not detected in any of the CAV 4 samples. Therefore, the results presented here focus exclusively on the second site studied, the *terramara* of Fondo Paviani near Verona (Fig. 1).

The Terramare culture

The Terramare culture developed in the central Po valley between the transition from MBA1 to MBA2 and the end of RBA¹ (first half of the 16th to ca. mid-12th century BCE) (Cardarelli 2010; Cupitò and Leonardi 2015). It had direct links with the cultural developments in the first part of the MBA in the northern Po valley. In this early phase, mainly pile-dwellings (*palafitte* in Italian) are known, as was the case during the preceding Early Bronze Age Polada culture (Cupitò and Leonardi 2015; Leonardi *et al.* 2015). The Terramare culture is characterised by settlements of subquadrangular layout surrounded by a fortification system consisting of an earthen rampart and a moat; these settlements are called *terramare* (singular *terramara*) (Bernabò Brea *et al.* 1997; Cardarelli 2010; Cupitò 2012; Cupitò and Leonardi 2015). One of the most important and distinctive traits of the culture is the advanced skills in water management and land reclamation (*e.g.* Balista and De Guio 1997; Cremaschi *et al.* 2006, 2016; Bernabò Brea and Cremaschi 2009). Contemporaneous cemeteries are often found near *terramare* settlements, and are characterised by both inhumations and cremations in the northern Po valley and only by cremations in the southern Po valley (*e.g.* De Marinis and Salzani 1997; Cardarelli and Tirabassi 1997; Cupitò and Leonardi 2015). The communities practiced a mixed food economy, relying on both crop cultivation and herding of domesticated animals for subsistence (Nisbet and Rottoli 1997; Riedel 1997; Mercuri *et al.* 2006; Carra 2009; De Grossi Mazzorin 2009, 2015; Rottoli and Castiglioni 2009a, 2009b; Berto and Rottoli 2015; Bertolini *et al.* 2015; Cremaschi *et al.* 2016; Dal Corso 2018). During its more than 400 year long duration, particularly during its final phase, the Terramare culture – specifically the area of the so-called Valli Grandi Veronesi polity (Balista and De Guio 1997; Cupitò and Leonardi 2015; Cupitò *et al.* 2015a) – functioned as the main cultural hub connecting Europe and the Mediterranean world. Thanks to its geographical position, the Terramare area was at the centre of a dense net of routes of exchange and circulation of raw materials (first of all copper and amber), involving north-central Europe, the Alps, peninsular Italy, the Aegean and the Levant (Bellintani 1997, 122-126; Carancini and Peroni 1997; Cardarelli 2010, 452; Bettelli *et al.* 2015, 2017; Cupitò and Dalla Longa 2015; Cupitò and Leonardi 2015; Cupitò *et al.* 2020, 299-301).

The *terramara* of Fondo Paviani was occupied from the advanced MBA to the first phases of the FBA (ca. mid-14th to 11th century BCE; Cupitò *et al.* 2015a). The site is located in the Valli Grandi Veronesi lowlands, in the lower plain north of the Po River. This is a subsiding portion of the Po valley, characterised by Pleistocene alluvial ridges and incised valleys, crossed during the Holocene by spring-fed rivers (Nicosia *et al.* 2011). After a first occupation phase, in which the site was surrounded by a possible palisade and a shallow moat, in the RBA (between the 13th and the

1 Abbreviations for the Bronze Age periods and phases in northern Italy: EBA=Early Bronze Age; MBA=Middle Bronze Age; RBA=Recent Bronze Age; FBA=Final Bronze Age. Further subphases and their correspondence with radiocarbon dating follow Pacciarelli (2001), Cardarelli (2010) and Leonardi *et al.* (2015): EBA1=ca. 2200-1800 BCE; EBA2=ca. 1800-1650 BCE; MBA1=ca. 1650-1550 BCE; MBA2=ca. 1550-1450 BCE; MBA3=ca. 1450-1350/1300 BCE; RBA1=ca. 1350/1300-1200; RBA2=1200-1150 BCE; FBA1-2=ca. 1150-1000 BCE; FBA3=ca. 1000-950/925 BCE.




Figure 1. Location of the site of Fondo Paviani, ca. 40 km southeast of Verona, Italy.

beginning of the 12th century BCE), a peripheral earthen rampart was erected (Balista *et al.* 2012; Cupitò *et al.* 2015a; Dal Corso *et al.* 2017; Dal Corso 2018; Dalla Longa *et al.* 2019). Outside of the rampart, a water basin derived from a senescent branch of the spring-fed Menago River bordered the site, functioning as a large moat (Dal Corso *et al.* 2017; Dal Corso 2018; Dalla Longa *et al.* 2019).

This fortified settlement is one of the largest in the Terramare area. Its construction must have involved the exploitation of existing river channels and alluvial ridges and a partial modification of the surrounding landscape (Deiana *et al.* 2020). In the RBA2, in the first half of the 12th century BCE, the settlement was at its largest, covering ca. 16–20 ha. Typological and archaeometric studies of ceramic, metal, amber and glass artefacts testify that Fondo Paviani was at that time an important centre within a network of contacts between the Aegean-Mycenaean and Levantine worlds (Jones *et al.* 2014; Bettelli *et al.* 2015, 2017), continental Europe, and peninsular Italy (Angelini *et al.* 2015; Cupitò *et al.* 2015a, 2015b, 2015c; Dalla Longa *et al.* 2015; Marcon and Mazzoli, 2015; Strafella *et al.* 2015; Vicenzutto *et al.* 2015). Participation in these exchange networks was probably favoured by the geographical position of the settlement and of the Valli Grandi Veronesi polity of sites depending on it. They were located on the system of spring-fed rivers (the Tione, Tartaro, Tregnone and Menago rivers) that connected the Adige River to the Po River. Fondo Paviani and its polity were thus at the core and, in the RBA, in control (Cupitò *et al.* 2020) of a route connecting north-central Europe and the Alps to the Adriatic coast and the Mycenaean and Levantine cultural worlds.

At the end of the RBA, the Terramare system was hit by a deep crisis (Cardarelli 2010). Different proxies detect environmental instability linked to hydrological and climatic changes at the end of the 2nd millennium BCE, which likely affected the Terramare socio-economic stability, although other, anthropogenic, causes, such as demographic pressure and the subsequent overexploitation of the territory and natural resources, must also be considered (Cremaschi *et al.* 2006, 2016; Cardarelli 2010; Mercuri *et al.* 2011; Ravazzi *et al.* 2012; Dal Corso 2018; Dalla Longa *et al.* 2019). We should emphasise that the crisis affected in particular the Terramare sites of the southern Po valley, for which a true collapse can be detected at the end of the RBA, in the 12th century BCE (Cardarelli 2010). The northern Po valley was also affected by the crisis but, here, a change in the settlement system occurred. There was a hyper-concentration of settlements and the development of a new settlement geometry, as well as the emergence of a culturally-influential phenomenon that is the rise of the Frattesina settlement as a central place (Cupitò and Leonardi 2015; Cupitò *et al.* 2015a, 2020).

Fondo Paviani is one of the sites that displayed resilience. The site continued to be settled until the first phases of the FBA, despite a reduction in size in the later period (Cupitò *et al.* 2015a). In the excavated portion of the site, a significant functional transformation occurred in the first phases of the FBA, in which the area used for dwelling and production was converted into arable land used for gardens (Dalla Longa 2020). Arable plots with ridge-and-furrow system have been identified in the




Figure 2. a. Cereal record from two refuse pits at Fondo Paviani based on charred carpological remains, in this case grains; b. crop-processing by-products from the same stratigraphic units (for Unit 317, see Berto and Rottoli 2015; for Unit 207, see Berto *et al.* in press).

stratigraphy by means of soil micromorphology and geochemical analysis (Nicosia *et al.* 2011; Dal Corso *et al.* 2012; Cupitò *et al.* 2015a).

Evidence of millet at Fondo Paviani

At Fondo Paviani, millet was detected as both charred carpological remains and phytoliths.

Charred cereal grains and other parts of the ears came mostly from the flotation of the fill of postholes, later used as refuse pits, within the settlement. These features date to the transition from the RBA1 to RBA2 (beginning of the 12th century BCE). The taxonomic composition of the cereal record is shown in Figures 2a (grains) and 2b (by-products). In pit fill Unit 317 (Berto and Rottoli 2015), emmer (*Triticum dicoccum*) is the dominant cereal, followed by several other hulled wheats (einkorn [*T. monococcum*], spelt wheat [*T. spelta*], ‘new type’ glume wheat [*T. timopheevii*]) and rare finds of free-threshing wheat and barley. Broomcorn millet is rare in Unit 317 (2 caryopses). The composition of macro-remains from another refuse pit (pit fill Unit 207; Berto *et al.* in press) is very different and shows the dominance of charred Panicoideae grains, with secure identification of broomcorn millet and foxtail millet (Fig. 2). Despite the minor presence of barley and wheats among the grains in Unit 207, the same wheat species are documented in both units (317 and 207) as glume bases and other by-products derived from the cleaning prior to food preparation. In both pits, the record is clearly dominated by cereals (emmer in Unit 317 and millets in Unit 207), probably representing lost food from an advanced stage of processing.

In the sediment from the ditch of the pre-rampart phase (sequence FP10 in Dal Corso *et al.* 2017; Dal Corso 2018) phytolith morphotypes from millet-like inflorescence were found (Fig. 3; for modern material see Lu *et al.* 2009). They are accompanied by more general phytolith morphotypes typical of (but not exclusive to) panicoid grasses, such as grass short cell bilobates, crosses and polylobates (Out and Madella 2016). These remains were particularly abundant in the ash layers in the ditch, likely coming from the dumping of rubbish before or in conjunction with levelling for the base of the peripheral earthen rampart.

In general, the cereal spectrum at Fondo Paviani shows that in the RBA, several different species were cultivated, with an important role given to micro-thermal cereals, usually growing in winter, such as hulled wheat, and drought-resistant macro-thermal cereal, or summer crops, such as millets. From the carpological remains, as well as from the phytolith evidence, we know that *Setaria italica* was also present.




Figure 3. Micro-photographs of phytoliths from the inflorescence of *Panicoideae* (cf. *Panicum miliaceum* glumes) found at Fondo Paviani in the ditch of pre-rampart phase, Unit 9b.

Materials and methods

Modern samples

The investigation of possible miliacin producers has so far involved a relatively limited number of taxa. Furthermore, contrasting reports - such as those on *Setaria italica* and *Chaetomium olivaceum* mentioned above - hinder reliable determination of all possible miliacin sources. In order to address this issue by generating additional data, we tested the miliacin content of modern grains belonging to a subset of taxa within the grass subfamily *Panicoideae*. We focused on hulled grains because of the outer waxing layer present on glumes, where miliacin could be found. Grains of *Panicum miliaceum* were tested in order to evaluate the reliability of the extraction procedure from grains. *Setaria italica* and *Echinochloa crus-galli* were selected due to their documented presence in northern Italian Bronze Age plant assemblages (Tafuri *et al.* 2018). Additional species (*Setaria viridis* and *Echinochloa frumentacea*) were included thanks to the easy access to modern reference material.

Archaeological material from Fondo Paviani


For the present study, 16 sediment samples were analysed (Fig. 4; Table 1). Samples 1-8 originate from the area of the peripheral shallow ditch of the pre-rampart phase. Samples 9-14 come from a posthole, later used as waste pit and filled with charred plant remains, including mostly cereals and cereal by-products (Berto and Rottoli 2015). This waste pit fill (Unit 317, see above) is datable to the RBA, more specifically, to the transition from a late phase of the RBA1 to the RBA2 (Dalla Longa 2020). Finally, samples 15 and 16 were collected from two extensive layers which formed in the open and included trampled palaeo-surfaces located in the same excavation trench as the posthole (Nicosia *et al.* 2011).

Sediment samples measuring ca. 20 g each were collected during fieldwork in 2010, conducted as collaboration between the universities of Kiel and of Padova. The sampling was carried out on freshly cleaned sediment surfaces, with metal tools (knife, spoon) that were washed with water and dried with paper towels between the samples. Dry samples were stored at room temperature in sterile plastic bags labelled on the outside. In the case of the ditch fill, a monolith in a plastic frame was collected and stored in the cold storage at Kiel University (Dal Corso *et al.* 2017; Dal Corso 2018). From each sample, a 1 cm³-subsample of sediment was taken in 2015, in a laboratory at Kiel University.

Strati-graphic unit	Interpretation	Description	References
Ditch of the pre-rampart phase			
Unit 2001	Base of the rampart	Extended silty deposit without archaeological materials	Dal Corso <i>et al.</i> 2017 (Unit D1); Dal Corso 2018
Unit 9a	Ditch fill with domestic waste	Silt and organic debris, including charcoal and some archaeological materials	Dal Corso <i>et al.</i> 2017 (Unit D2); Dal Corso 2018
Unit 9b	Ditch fill with ashes	Discontinuous ash, charcoal and sand lenses	Dal Corso <i>et al.</i> 2017 (Unit D3); Dal Corso 2018
Unit 9c	Ditch fill with domestic waste	Silty loam deposit with charcoal and archaeological materials (e.g. potsherds)	Dal Corso <i>et al.</i> 2017 (Unit D4); Dal Corso 2018
Unit 9c	Ditch fill with ashes	Ash lens with abundant wood charcoal	Dal Corso <i>et al.</i> 2017 (Unit D5); Dal Corso 2018
Unit 2038	Natural fluvial substrate	Calcareous silty clay alluvial deposit	Dal Corso <i>et al.</i> 2017 (Unit D7); Dal Corso 2018
Refuse pit fill 317			
317a	Waste pit fill, uppermost portion	Silty loam deposit with very abundant charred cereals and charcoal	Unpublished
317b	Waste pit fill, central portion	Silty loam deposit with very abundant charred cereals and charcoal	Unpublished
317c	Waste pit fill, central portion	Silty loam deposit with very abundant charred cereals and charcoal	Unpublished
317d	Waste pit fill, lower portion	Mostly charred cereals and by-products	Berto and Rottoli 2015, Fig. 1
321	Waste pit fill, upper eastern portion	Silty loam deposit with very abundant charred cereals and charcoal	Unpublished
322	Waste pit fill, upper eastern portion	Silty loam deposit with very abundant charred cereals and charcoal	Unpublished
Cultural layers			
19a	Uppermost portion of the anthropogenic deposit detected in the extensive layers which formed and were trampled in the open; intra-site gardening/cultivation of arable plots still active in the last phase of settlement (first phases of FBA) but reworking sediments and archaeological materials from the precedent phase (RBA2)	Clay-loam deposit with frequent highly fragmented 2-5 mm artefacts (charcoal, pottery, burnt sediment), wavy top boundary derived from agricultural practice	Nicosia <i>et al.</i> 2011 (top of Unit 4, Horizon 4Aant1)
19b	Central portion of the anthropogenic deposit detected in the extensive layers which formed and were trampled in the open; dwelling layers of human occupation (RBA2)	Silty clay-loam deposit with artefacts of different size classes (charcoal, pottery, burnt sediment, bones, shells)	Nicosia <i>et al.</i> 2011 (central part of Unit 4, Horizon 4Aant2)

Table 1. Description of the litho-stratigraphic units sampled for biomarker analysis at the site of Fondo Paviani.

Figure 4. Field photographs of the archaeological features that have been sampled for miliacin analysis at Fondo Paviani, 2010 campaign. a. the ditch of pre-rampart phase; b. detail of the section of the ditch shown in 4a that was successively sampled with a monolith; c. cultural layers from the excavation trench; d. refuse pit fill 317 and other fill units.



Miliacin extraction

Authentic miliacin was kindly provided by J  r  my Jacob. The methods for miliacin extraction from modern grains and sediment samples follow those presented in Bossard *et al.* (2013). Fresh modern grains from a range of species were obtained from botanical gardens in Germany. Lipids were extracted by ultrasonication (3  ) using dichloromethane (DCM):isopropanol (2:1) and the extracts were combined. The total lipid extract was separated into neutral, acidic and polar fractions on a Pasteur pipette filled with aminopropyl-bonded silica. The neutral fraction was collected after elution with DCM:isopropanol 2:1. The neutral fraction was separated using flash chromatography on silica (activated at 120  C for 24 h, then deactivated with H  O at 5% by wt.) into aliphatic and aromatic hydrocarbons, ethers, esters and ketones, and alcohols by sequential elution with solvent of increasing polarity. All fractions were dried under a stream of N   and stored at   20  C until the analysis using gas chromatography-mass spectrometry (GC-MS). In order to quantify the miliacin in the extracts, a known amount of 5  -cholestane was added as internal standard prior to GC-MS. Miliacin was detected by GC retention time and mass spectrometry (Heron *et al.* 2016).

Results

Miliacin sources

Among all modern specimens tested, miliacin was detected exclusively in *Panicum miliaceum* (Table 2). The absence of detectable miliacin from *Setaria italica* and *Echinochloa crus-galli* is in agreement with the findings of Bossard *et al.* (2013) and Ohmoto *et al.* (1970). Results are shown in the chromatograms in Figure 5. The authentic miliacin standard shows a single peak eluting at 28.157 mins. The molecular ion (M  ) is visible at *m/z* 440 with prominent fragment ions at *m/z* 189 and 204. In Figure 5a, the authentic miliacin standard (upper) is compared with results from the modern samples *Panicum* 1 (middle) and *Panicum* 2 (lower). This confirms the presence of miliacin in the grains of this species through presence of the molecular ion (M  ) at *m/z* 440. The mass spectral data confirm the common origin for the peak. In broomcorn millet, Bossard *et al.* (2013) report yields of 297-476   g miliacin per g   of grain and our results are not inconsistent with these yields.

In Figure 5b, the authentic miliacin standard (upper) is compared with samples of modern *Setaria viridis*, *S. italica* and *S. verticillata*. Once again, the molecular ion (M  ) at *m/z* 440 only is traced: the single peak is from the authentic miliacin, which confirms that no miliacin is present in any of the *Setaria* species analysed. Finally, in Figure 5c, the authentic miliacin standard (upper) is compared with modern samples of *Echinochloa crus-galli* and *E. frumentacea*. The molecular ion (M  ) at *m/z* 440 only is traced; the single peak is from the authentic miliacin only, confirming also in this case that no miliacin is present in the *Echinochloa* species analysed. The very minor peak eluting just before 28 minutes is common to all *Echinochloa* species. It is likely to be a similar compound to miliacin (that is, a triterpene methyl ether).

Figure 5. Extracted ion (m/z 440) chromatograms to test for the presence of miliacin in modern species of Panicoideae. a. authentic miliacin and two Panicum miliaceum extracts; b. authentic miliacin and five Setaria species extracts; c. authentic miliacin and four Echinochloa species extracts. Only the Panicum samples exhibit a chromatographic peak with the same retention time and mass spectral characteristics as authentic miliacin.




Table 2. Results showing the presence or absence of miliacin in the modern species of Panicoideae analysed in this study.

Sample	Plant species (grains)	Weight of grains extracted	Miliacin present	Yield (µg g ⁻¹)
Panicum 1	<i>Panicum miliaceum</i>	122 mg	Yes	703
Panicum 2	<i>Panicum miliaceum</i>	129 mg	Yes	380
Setaria vir 1	<i>Setaria viridis</i>	124 mg	No	-
Setaria vir 2	<i>Setaria viridis</i>	128 mg	No	-
Setaria vert 1	<i>Setaria verticillata</i>	129 mg	No	-
Setaria vert 2	<i>Setaria verticillata</i>	126 mg	No	-
Setaria it 1	<i>Setaria italica</i>	128 mg	No	-
Setaria it 2	<i>Setaria italica</i>	128 mg	No	-
Echinochloa c-g 1	<i>Echinochloa crus-galli</i>	117 mg	No	-
Echinochloa c-g 2	<i>Echinochloa crus-galli</i>	127 mg	No	-
Echinochloa frum 1	<i>Echinochloa frumentacea</i>	123 mg	No	-
Echinochloa frum 2	<i>Echinochloa frumentacea</i>	122 mg	No	-

Miliacin at Fondo Paviani

Miliacin was detected in all of the analysed sediment samples from Fondo Paviani – from the ditch fill (units 2001, 91, 9b, 9c, 2038), the pit (units 317a, 317b, 317c, 317d, 321, 322) and the two extensive layers which formed and were trampled in the open (Units 19a, 19b) (Table 3).

Discussion

Miliacin from modern reference material

From a methodological point of view, this study provides further evidence of miliacin in modern *Panicum miliaceum* grains, with yields of 380-703 µg miliacin per g-1 of grain, which is comparable to those obtained by Bossard *et al.* (2013), who report yields of 297-476 µg per g⁻¹. Miliacin is absent in the grains from the other Panicoideae here analysed, namely *Setaria* sp. (*S. italica*, *S. viridis* and *S. verticillata*) and *Echinochloa* sp. (*E. crus-galli* and *E. frumentacea*). Some of these species are attested in the archaeobotanical record from prehistoric sites in northern Italy (*e.g.* Rottoli 2001; Carra 2009; Rottoli and Castiglioni 2009a, 2009b; Perego 2015; Rottoli and Pessina 2016). Our results from the modern plant material support published data regarding the absence of miliacin in *S. italica* (Bossard *et al.* 2013).

Sample no.	Stratigraphic unit	Miliacin content (µg/g)
<i>Ditch of the pre-rampart phase</i>		
1	Unit 2001 cm ²	<0.01
2	Unit 9a	0.70/1.12
3	Unit 9b (ash layer)	9.01
4	Unit 9b (charcoal layer)	1.34
5	Unit 9b (ash layer)	2.39
6	Unit 9c	8.34
7	Unit 9c (charcoal layer)	2.08
8	Unit 2038	2.08/2.67
<i>Waste pit fill 317</i>		
9	317a	6.23
10	317b	7.47
11	317c	3.88
12	317d	4.59
13	321	7.54
14	322	5.73
<i>Cultural layers</i>		
15	19a	1.57
16	19b	5.73

Table 3. Results showing the amount of miliacin per gram of sediment from the site of Fondo Paviani.




Figure 6. Partial total ion current (TIC) chromatogram showing the presence of miliacin as the sole PTME from a sample in the ditch of the pre-rampart phase, within an ash layer (Unit 9b, sample 3).

Miliacin from archaeological contexts in the floodplain

We applied miliacin extraction procedure on sediment samples from the Late Bronze Age site of Fondo Paviani in the Po valley. This is the first time that the extraction of miliacin from sediment has been undertaken in northern Italy. The feasibility of using this biomarker to attest the presence of millet in archaeological contexts in alluvial plains has been confirmed by this case study, supporting the well-known studies of lake sediments (*e.g.* Jakob *et al.* 2008, 2009; Simmoneau *et al.* 2013) and ‘food crust’ on ceramic containers (*e.g.* Heron *et al.* 2016).

Miliacin in the pre-rampart phase ditch: Millet consumption

The results of biomarker analysis show that miliacin can be traced in all of the samples, although with different yields. To contextualise these results, other materials in the deposit and the formation processes will be described, previously identified through the analysis of plant micro-fossils and soil micromorphology (Dal Corso *et al.* 2017) and using the archaeological data (Cupitò *et al.* 2015a).

In this study, the lowest miliacin yields came from a minerogenic unit constituting the base of the peripheral earthen rampart (sample 1). Here, miliacin is present only in traces. The next-lowest miliacin yield was from the alluvial deposit (sample 8) that forms the natural substrate under the ditch fill. In soil thin-section, this alluvial unit showed moderate porosity and few charcoal fragments (see Unit D7; Dal Corso *et al.* 2017); some sediment may thus have percolated from the anthropogenic deposits above and be responsible for the trace of miliacin. These units concern the ditch fill, where dumps of waste from routine activities are found (Units 9b, 9c; see D3-D5 in Dal Corso *et al.* 2017), together with some washed-in sediment (Unit 9a; see D2 in Dal Corso *et al.* 2017).

The highest miliacin yields in this study are found in ash lenses in the ditch fill (samples 3 and 6, 8.34 and 9.01 $\mu\text{g g}^{-1}$, respectively). These ashes are largely composed of phytoliths derived from chaff of pooid and panicoid cereals and from some grass leaves and stems possibly coming from cereal straw. The pollen record, which is dominated by Cerealia-type pollen (40%) and wild Poaceae pollen (over 30%), supports the phytolith record (Dal Corso *et al.* 2017). Matching the high miliacin content, there are abundant phytoliths (Fig. 3) from panicoid inflorescence comparable with that of broomcorn millet and foxtail millet (Lu *et al.* 2009). Despite being found in the ditch in a secondary deposit, the combination of a number of proxies suggest that these ashes derived from the burning of dung from domestic animals (Dal Corso *et al.* 2017; Dal Corso 2018). This practice possibly relates to the cleaning of stabling areas or from the deliberate use of dung as fuel, although wood charcoal is present too and wood was most likely the primary fuel. Further proxies indicative of herbivore dung are the abundant faecal spherulites in the thin sections, as well as the coprophilous fungal spores (*e.g.* *Sordaria* sp. and *Coniochaeta* sp.) and eggs of the intestinal parasite *Trichuris*. As mentioned above, the associated botanical micro-fossils show a high concentration of grass phytoliths, many still in anatomical position in multicellular aggregates, including crop processing remains, such as chaff and straw (possibly chopped or trampled, considering the cutmarks in the phytolith aggregates), abundant pollen from wild grasses, and pollen from other herbaceous plants (Dal Corso *et al.* 2017; Dal Corso 2018). The latter does not suggest human faecal material; most likely it derives from millet hulls and green forage given to herbivores. Sheep and pig are the most frequent animals in the faunal record at Fondo Paviani (De Grossi Mazzorin 2015). The presence of dung suggests that animals were likely kept inside the settlement at least in some phases of their lives, according to the seasonality of production or processing of meat or secondary products (De Grossi Mazzorin 2015).

Studies in other parts of the world suggest the practice of winter foddering of caprines with millets (e.g. Hermes *et al.* 2019). Further, modern uses of broomcorn millet as fodder include its mixing with barley for feeding pigs or part-grinding the grains for fattening lambs; in general, millet is considered a useful fall-back crop in case of loss of other yields (Baltensperger *et al.* 1995; Berglund 2007; Tran 2015, 2017). At Fondo Paviani, the use of broomcorn millet as fodder is seen only in some pigs exhibiting high $\delta^{13}\text{C}$ values (Tafuri *et al.* 2018).

At some sites in the low and middle Verona Plain, the primary evidence of millet consumption comes from the study of carbon and nitrogen stable isotope composition of osteological remains of humans and animals. These results stem from eight sites – both cemeteries and settlements – covering the entire EBA-RBA (Tafuri *et al.* 2009, 2018). Human remains from the settlement of Dossetto di Nogara (middle-advanced EBA1-EBA2; 1 sample) and the cemeteries of Olmo di Nogara (MBA2-RBA2; 64 samples) and Bovolone (MBA3/RBA1 transitional phase-RBA2; 21 samples) are enriched in ^{13}C , which is here attributed to the consumption of millets within a diet based on terrestrial food. Among these sites, Olmo di Nogara is a particularly useful case study for the present work because, dating from the middle phase of the MBA to the RBA2 (mid-16th to mid-12th century BCE), it is partially contemporaneous with the last occupation phase of Fondo Paviani (RBA). At this cemetery, inhumation burials prevail over the cremation burials and, in the MBA, the existence of elite is evident from the rich grave goods, with swords and daggers buried with men and parures with women, composed of pins made of bronze and amber (Cupitò and Leonardi 2005; Cupitò 2006; Canci *et al.* 2015). According to the anthropological, palaeopathological and isotopic analyses, the social differentiation seen in the grave goods is not reflected in the generally good health condition and nutrition of the individuals buried in the cemetery (Canci *et al.* 2015). The diet of all individuals, including children, contained millet, but that of women was also nitrogen rich and for this reason differed from that of the rest of the population (Tafuri *et al.* 2018). From Fondo Paviani, isotopic analysis were performed only on animal bones: cattle (5 samples), sheep (4) and pig (8) (Tafuri *et al.* 2018). Only pigs show traces of millet consumption. It is possible that crop processing waste was given to these animals, while dehusked millet grains were used as food for humans.

Reconstruction of the beginning of millet consumption in this region requires further investigation. Thus far, few millet grains have been retrieved from some Neolithic and Copper Age contexts (Rottoli and Castiglioni 2009a) and the direct radiocarbon dates for these are missing or unpublished. Among the isotopic studies of the EBA individuals from the area of Verona, an infant from Dossetto di Nogara gave indications of C4 plant consumption (Tafuri *et al.* 2018), unlike the individuals from the cemetery of Arano di Cellore, which did not show such traces (Varalli *et al.* 2016). Future investigations and dating of the EBA macro-botanical, stable isotope and archaeological records will help to better interpret this early millet evidence. The radiocarbon-dated grains of millet available to date suggest that millet was consumed from at least the MBA1 (16th century BCE, after the modelled dates in Filipović *et al.* 2020). For the MBA2 to the RBA period, there is robust isotopic evidence for millet consumption from the cemeteries mentioned above (Tafuri *et al.* 2018).

Miliacin in pit fill 317: A diverse crop assemblage in the RBA

Pit fill 317 was subdivided in the field into several subunits (317a-d) and these all showed the presence of miliacin. Unit 317 is the fill of a refuse pit from the transitional period RBA1-RBA2. At this time, the excavated portion of the settlement features several clay-built open fireplaces, probably used for crop processing (Cupitò *et al.* 2015a). This pit, as well as the other ones (e.g. Unit 207; Fig. 2), was filled after a fire destroyed a possible small, raised wooden granary located in the area, and the charred cereals were removed

from the surface (Cupitò *et al.* 2019). The highest miliacin content in this archaeological context comes from the uppermost portion of the fill. The macro-botanical record is mostly composed of grains of other cereals, all known from the Bronze Age elsewhere in Europe (Stika and Heiss 2013), such as emmer, einkorn, barley, 'new type' glume wheat and free-threshing wheat (Fig. 2). Among the other identified remains are some crop weeds, a few hazelnuts and berries collected from the wild vegetation near the site (Berto and Rottoli 2015). Although broomcorn millet and foxtail millet are not abundant here, they were documented in other features in the same excavation trench (*e.g.* 207; Fig. 2). This broad crop spectrum and the presence of wild fruit and nut suggest a diverse plant-based diet of the inhabitants of Fondo Paviani. Such varied plant use may have been a strategy to buffer poor crop yields, whilst it indicates solid knowledge of plant use and cultivation. Pulses are much less represented than cereals in the studied samples – bitter vetch (*Vicia ervilia*) and common pea (*Pisum sativum*) were recognised (Berto and Rottoli 2015; Berto *et al.* in press).

Miliacin in extensive anthropogenic layer 19b: Domestic waste?

The extensive anthropogenic layer 19b, the samples from which gave high yield of miliacin ($5.73 \mu\text{g g}^{-1}$), belongs to the occupation phase in which the excavated portion of the settlement received dumps of domestic materials, both delimited and scattered (Cupitò *et al.* 2015a; Dalla Longa 2020). Unit 19b can be chronologically ascribed to the RBA2 (Cupitò *et al.* 2015a; Dalla Longa 2020). This layer was excavated in open area and was also recognised in two stratigraphic profiles (although these were not sampled for miliacin), where the anthropogenic units were investigated by soil micromorphology, geochemistry (Nicosia *et al.* 2011) and palynology (Dal Corso *et al.* 2012). These deposits are composed of a series of surface horizons on dry soil, where fine sediment accumulated in the form of material from dug-out features, import from alluvial terraces outside the settlement area, disruption of earth-based construction material and occasional (probably seasonal) overflow from the local drainage (see Unit 4 and horizon 4Anth2; Nicosia *et al.* 2011). The miliacin in this unit probably derives from domestic waste. The earthen construction material in other *terramare* is usually composed of ovicaprid dung (Cardarelli *et al.* 1997; Cremaschi and Ottomano 1998; Cremaschi *et al.* 2004). More work is needed to determine if caprines were foddered with millet at Fondo Paviani. There is no direct evidence of this (Tafari *et al.* 2018), apart from the association of herbivore dung and millet recovered from the ashes in the ditch (Dal Corso *et al.* 2017).

Miliacin in extensive layer 19a: Millet also in the first phases of the Final Bronze Age arable plots?

Unit 19a, overlying Unit 19b, is interpreted as an arable zone within a ridge-and-furrow field system. This is evidenced by the uneven distribution in the soil of finely fragmented, worn anthropogenic materials (ceramics, charcoal, bones, fired clay or daub) with rounded edges, corresponding to manure derived from domestic waste (4Anth1 in Nicosia *et al.* 2011). This unit was also excavated in open area and was recognised in the two stratigraphic profiles, as in the case of Unit 19b. The arable fields were established after a functional change detected in the excavated area, which, as we note above, previously hosted domestic deposits (Unit 19b) (Cupitò *et al.* 2015a; Dalla Longa 2020). Agricultural activities brought up material from the deposits below, and some pedo-relicts show the import of allochthonous soil, brought here on purpose from outside the settlement (Nicosia *et al.* 2011). This unit has a strong reworking of sediments and points to the arable use in the first phases of the FBA, but includes also RBA2 materials and sediments (Cupitò *et al.* 2015a). For this reason, the presence of miliacin here cannot be taken as clearly indicating local millet cultivation.

Conclusions

In the Late Bronze Age *terramara* of Fondo Paviani, in the Po valley, miliacin is present in many anthropogenic deposits. These early results support the validity of miliacin extraction and identification as a tool to track the presence of millet in different sediments from settlement sites. This cereal was previously documented at the site in the form of macro-remains and phytoliths. A multi-proxy approach applied to investigate the ditch fill revealed a concentration of grass and millet phytoliths in ashes, most likely derived from the practice of burning litter or dung from domestic animals fed with millet. Miliacin as well as charred millet caryopses and processing remains were registered in pit fills and in extensive cultural layers formed and trampled in the open. A stable isotope study of the three main domesticates attested at Fondo Paviani (sheep, pig and cattle) only showed evidence of millet consumption in pig, probably because this species thrived on remains of crop processing and other domestic waste. The MBA2-RBA human population living in the region also consumed millet, at least those individuals analysed isotopically, including an exceptional case of an EBA infant fed with millet (Tafuri *et al.* 2018). Broomcorn and foxtail millets were part of the diverse crop spectrum used by the inhabitants of Fondo Paviani, suggesting food economy aimed at preventing harvest failures and at the use of different environments for cultivation. More should be done to investigate winter fodder strategies and the possible use of millet for herbivores at the site.

Further research is required on the long-term behaviour of miliacin in sedimentary contexts (Jacob *et al.* 2008; Courel *et al.* 2017). Compound-specific radiocarbon dating of miliacin has considerable potential (Courel *et al.* 2017). The presence of miliacin in the RBA at Fondo Paviani is in accordance with other evidence signalling consumption of millet during this period (Berto and Rottoli 2015; Tafuri *et al.* 2018). It is not possible to comment on the last settlement phase, dated to the first phases of the FBA. The only layer that has chronological markers attributable to this phase (Unit 19a) also includes materials and sediment pulled up from the RBA2 layer below it (Unit 19b). The miliacin content detected in Unit 19a may then also derive from this reworked sediment. In order to find out if millet was used in the first phases of the FBA, and also to test the method for detection of miliacin within such a deposit, a multi-proxy approach should be applied in an FBA context, where layers specific to this chronological span are present. A suitable site to attempt this could be Frattesina, as it contains both settlement layers and inhumation burials dating to the first phase of the FBA.

Acknowledgements

The authors wish to thank Dr J  r  my Jacob (Laboratoire des Sciences du Climat et de l'Environnement, France) for providing the authentic miliacin standard used in this study.

References

- Balista, C. and De Guio A., 1997. Ambiente ed insediamenti dell'et  del bronzo nelle Valli Grandi Veronesi. In: M. Bernab  Brea, A. Cardarelli, M. Cremaschi, eds., *Le Terramare: la pi  antica civilt  padana*. Catalogo della mostra di Modena. Milano: Mondadori Electa, 137-160.
- Baltensperger, D., Lyon, D., Anderson, R., Holman, T., Stymiest, C., Shanahan, J., Nelson, L., DeBoer, K., Hein, G., Kral, J., 1995. *Producing and Marketing Proso Millet in the High Plains*. EC95-137. Lincoln: University of Nebraska Cooperative Extension.

- Bellintani P., 1997. Frattesina: l'ambra e la produzione vitrea nel contesto delle relazioni transalpine. In: L. Endrizzi and F. Marzatico, eds., *Ori delle Alpi*. Trento: Provincia Autonoma de Trento, 116-129.
- Bernabò Brea, M. and Cremaschi, M., eds., 2009. *Acqua e civiltà nelle Terramare: la vasca votiva di Noceto*. Milano: Skira.
- Bernabò Brea, M., Cardarelli, A., Cremaschi, M., eds., 1997. *Le Terramare: la più antica civiltà padana Catalogo della mostra di Modena*. Catalogo della mostra di Modena. Milano: Mondadori Electa.
- Berglund, D.R., 2007. *Proso Millet in North Dakota*. NDSU A-805. Fargo: North Dakota State University Agriculture and University Extension.
- Berto, F. and Rottoli, M., 2015. Agricoltura e raccolta in un insediamento del Bronzo recente della pianura veronese: il pozzetto US 317 di Fondo Paviani (Verona)-Sca-vi Università di Padova 2007-2012. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria Veneto, 829-832.
- Berto, F., Cupitò, M., Leonardi, G., Rottoli, M., in press. Alimentazione e strategie di sussistenza in un contesto della tarda Età del Bronzo dell'Italia settentrionale: le analisi archeobotaniche del 'pozzetto' US 317 e di altri contesti a Fondo Paviani (VR). Proceedings of Preistoria del cibo. Studi di Preistoria e Protostoria 6. Firenze: Istituto Italiano di Preistoria e Protostoria.
- Bertolini, M., Zanini, S., Thun Hohenstein, U., 2015. Nuovi dati sullo sfruttamento e gestione delle risorse animali tra il Bronzo antico ed il Bronzo recente nei territori del mediobasso Veronese e il basso Polesine. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 321-326.
- Bettelli, M., Cupitò, M., Levi, S.T., Jones, R., Leonardi, G., 2015. Tempi e modi della connessione tra mondo egeo e area padano-veneta: una riconsiderazione della problematica alla luce delle nuove ceramiche di tipo miceneo di Fondo Paviani (Legnago, Verona). In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 377-388.
- Bettelli, M., Cupitò, M., Jones, R., Leonardi, G., Levi, S.T., 2017. *The Po Plain, Adriatic and Eastern Mediterranean in the Late Bronze Age: Fact, Fancy and Plausibility*. AEGAEUM – Annales liégoises et PASpiennes d'archéologie égéenne 41. Liège: Peters Leuven, 165-172.
- Beug, H.J., 2004. *Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete*. München: Verlag Dr. Fridrich Pfeil.
- Bossard, N., Jacob, J., Le Milbeau, C., Sauze, J., Terwilliger, V., Poissonnier, B., Vergès, E., 2013. Distribution of miliacin (olean-18-en-3 β -ol methyl ether) and related compounds in broomcorn millet (*Panicum miliaceum*) and other reputed sources: implications for the use of sedimentary miliacin as a tracer of millet. *Organic Geochemistry*, 63, 48-55. <https://doi.org/10.1016/j.orggeochem.2013.07.012>
- Canci, A., Cupitò, M., Pulcini, M.L., Salzani, L., Fornaciari, G., 2015. La necropoli della media e recente Età del bronzo di Olmo di Nogara (Verona): risultati della ricerca osteoarcheologica, paleochimica e paleodemografica. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 327-340.
- Carancini, G. and Peroni, R., 1997. La koinè metallurgica. In: M. Bernabò Brea, A. Cardarelli, M. Cremaschi, eds., *Le Terramare: la più antica civiltà padana*. Catalogo della mostra di Modena. Milano: Mondadori Electa, 595-601.
- Cardarelli, A., 2010. The collapse of the Terramare culture and growth of new economic and social systems during the Late Bronze Age in Italy. *Scienze dell'Antichità*, 15, 449-520.

- Cardarelli, A., Cremaschi, M., Cattani, M., Labate, D. and Steffé, G., 1997. Nuove ricerche nella terramara di Montale (MO): primi risultati. In: M. Bernabò Brea, A. Cardarelli, M. Cremaschi, eds., *Le Terramare: la più antica civiltà padana*. Catalogo della mostra di Modena. Milano: Mondadori Electa, 224-228.
- Cardarelli, A. and Tirabassi, J. 1997. Le necropoli delle terramare emiliane. In: M. Bernabò Brea, A. Cardarelli, M. Cremaschi, eds., *Le Terramare: la più antica civiltà padana*. Catalogo della mostra di Modena. Milano: Mondadori Electa, 677-697.
- Carra, M., 2009. Alimentazione, ambiente ed economia di sussistenza su base vegetale: studio archeobotanico preliminare dei macroresti provenienti dal sito di Solarolo. *IpoTESI di Preistoria*, 2 (1), 281-291. <https://doi.org/10.6092/issn.1974-7985/1613>
- Connor, H.E. and Purdie, A.W., 1976. Triterpene methyl ether differentiation in *Chionochloa* (Gramineae). *New Zealand Journal of Botany*, 14, 315-326. <https://doi.org/10.1080/0028825X.1976.10428904>
- Connor, H.E. and Purdie, A.W., 1981. Triterpene methyl ethers in *Chionochloa* (Gramineae): distribution in western South Island, New Zealand. *New Zealand Journal of Botany*, 19, 161-170. <https://doi.org/10.1080/0028825X.1981.10425116>
- Courel, B., Schaeffer, P., Adam, P., Motsch, E., Ebert, Q., Moser, E., Féliu, C., Bernasconi, S.M., Hajdas, I., Ertlen, D., Schwartz, D., 2017. Molecular, isotopic and radiocarbon evidence for broomcorn millet cropping in northeast France since the Bronze Age. *Organic Geochemistry*, 110, 13-24. <https://doi.org/10.1016/j.orggeochem.2017.03.002>
- Cremaschi, M., Mercuri, A.M., Torri, P., Florenzano, A., Pizzi, C., Marchesini, M., Zerboni, A., 2016. Climate change versus land management in the Po Plain (northern Italy) during the Bronze Age: new insights from the VP/VG sequence of the terramara Santa Rosa di Poviglio. *Quaternary Science Reviews*, 136, 153-172. <https://doi.org/10.1016/j.quascirev.2015.08.011>
- Cremaschi, M. and Ottomano, C., 1996. Il testimone stratigrafico di Monte Castellaccio e aspetti micromorfologici per lo studio dei processi di formazione del sito. In: M. Pacciarelli, ed., *La Collezione Scarabelli*. Preistoria 2. Imola: Grafis Edizioni, 148-157.
- Cremaschi, M., Ottomano, C., Trombino, L., 2004. Aspetti micromorfologici e pedologici nei processi di formazione della stratigrafia archeologica. In: M. Bernabò Brea and M. Cremaschi, eds., *Il villaggio piccolo della terramara di Santa Rosa di Poviglio e Scavi 1987-1992*. Firenze: Origines, 117-150.
- Cremaschi, M., Pizzi, C., Valsecchi, V., 2006. Water management and land use in the terramare and a possible climatic co-factor in their abandonment: the case study of the terramara of Poviglio Santa Rosa (northern Italy). *Quaternary International*, 151 (1), 87-98. <https://doi.org/10.1016/j.quaint.2006.01.020>
- Cupitò, M., 2006. La necropoli dell'età del bronzo di Povegliano Veronese: rilettura dei dati e nuove ipotesi interpretative a quarant'anni dalla revisione peroniana. In: *Studi di protostoria in onore di Renato Peroni*. Firenze: All'Insegna del Giglio, 30-34.
- Cupitò, M., 2012. Dinamiche costruttive e di degrado del sistema aggere-fossato della terramara di Castione dei Marchesi (Parma): rilettura e reinterpretazione dei dati ottocenteschi. *Rivista di Scienze Preistoriche*, 62, 231-248.
- Cupitò, M. and Dalla Longa, E., 2015. Cultural interaction (and integration?) phenomena between the Terramare world and peninsular Italy in the Middle Bronze Age: the pottery point of view. In: P. Suchowska-Ducke, S. Scott Reiter, H. Vandkilde, eds., *Forging Identities: the Mobility of Culture in Bronze Age Europe: Volume 2*. BAR International Series S2772. Oxford: BAR Publishing, 155-163.
- Cupitò M., Dalla Longa E., Bovolato C., Leonardi G., 2019, *Un'area per la conservazione e la trasformazione dei cereali nel Bronzo recente di Fondo Paviani (Legnago, Verona)*. Presented at the 6th Incontri Annuale di Preistoria e Protostoria, 'Focolari, forni e fornaci tra Neolitico ed Età del Ferro', Bologna, 29 March 2019. Bologna: Università di Bologna, Dipartimento di Storia Culture Civiltà, 101-104. <https://www.iipp.it/vi-iapp-abstract-book/>

- Cupitò, M. and Leonardi, G., 2015. Il Veneto tra Bronzo antico e Bronzo recente. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 201-239.
- Cupitò, M., Leonardi, G., Dalla Longa, E., Nicosia, C., Balista, C., 2015a. Fondo Paviani (Legnago, Verona): il central place della polity delle Valli Grandi Veronesi nella tarda Età del bronzo: cronologia, aspetti culturali, evoluzione delle strutture e trasformazioni paleoambientali. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 357-375.
- Cupitò, M., Angelini, I., Dalla Longa, E., 2015b. Nuovi manufatti in materiale vetroso da Fondo Paviani (Verona) – Scavi Università di Padova 2007-2012: tipocronologia e analisi archeometriche. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 855-859.
- Cupitò, M., Angelini, I., Nimis, P., 2015c. Il torques tipo Canegrate di Fondo Paviani (Verona) alla luce delle indagini archeometriche. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 839-844.
- Cupitò, M., Dalla Longa, E., Balista, C., 2020. From 'Valli Grandi Veronesi system' to 'Frattesina system': observations on the evolution of the exchange system models between Veneto Po valley area and the Mediterranean world during the Late Bronze Age. *Rivista di Scienze Preistoriche* 70, 293-310.
- Dal Corso, M., 2018. *Environmental History and Development of the Human Landscape in a Northeastern Italian Lowland during the Bronze Age: a Multidisciplinary Case-Study*. Universitätsforschungen zur Prehistorischen Archäologie – Band 312, HDL 12. Bonn: Dr. Rudolf Habelt GmbH.
- Dal Corso, M., Marchesini, M., Leonardi, G., Kirleis, W., 2012. Environmental changes and human impact during the Bronze Age in N-Italy: on-site palynological investigation at Fondo Paviani, Verona. In: J. Kneisel, W. Kirleis, M. Dal Corso, N. Taylor, V. Tiedtke, eds., *Collapse or Continuity?* Bonn: Dr. Rudolf Habelt GmbH, 71-84.
- Dal Corso, M., Nicosia, C., Balista, C., Cupitò, M., Dalla Longa, E., Leonardi, G., Kirleis, W., 2017. Bronze Age crop processing evidence in the phytolith assemblages from the ditch and fen around Fondo Paviani, northern Italy. *Vegetation History and Archaeobotany*, 26 (1), 5-24. <https://doi.org/10.1007/s00334-016-0573-z>
- Dal Corso, M., Pashkevych, G., Filipović, D., Liu, X., Motuzaite Matuzeviciute, G., Stobbe, A., Shatilo, L., Videiko, M., Kirleis, W., 2022. Between cereal agriculture and animal husbandry – Millet in the early economy of the north Pontic region. *Journal of World Prehistory*, 35(1).
- Dalla Longa, E., 2020. Uno spazio e tanti usi: analisi funzionale del 'Settore 2/2.1' di Fondo Paviani (Legnago, Verona) tra Bronzo Recente e Bronzo Finale. In: N. Negroni Catacchio, ed., *Atti del XIV Incontro di Studi 'Preistoria e Protostoria in Etruria', Archeologia dell'abitare. Insediamenti e organizzazione sociale prima della città. Dai monumenti ai comportamenti. Ricerche e scavi, 7-9 settembre 2018*. Milano: Centro Studi di Preistoria e Archeologia, 155-167.
- Dalla Longa, E., Cupitò, M., Vidale, M., Levi, S.T., Guida, G., Mariottini, M., Cannavò, V., 2015. Nuove ceramiche con decorazione di tipo appenninico da Fondo Paviani (Verona) – Scavi Università di Padova: inquadramento tipocronologico e indagini archeometriche. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 861-866.
- Dalla Longa, E., Dal Corso, M., Vicenzutto, D., Nicosia, C., Cupitò, M., 2019. The Bronze Age settlement of Fondo Paviani (Italy) in its territory: hydrography, settlement distribution, environment and in-site analysis. *Journal of Archaeological Science: Reports* 28, 102018. <http://dx.doi.org/10.1016/j.jasrep.2019.102018>

- De Grossi Mazzorin, J., 2009. Fauna ed economia animale. In: M. Bernabò Brea and M. Cremaschi, eds., *Acqua e civiltà nelle Terramare: la vasca votiva di Noceto*. Milano: Skira, 170-171.
- De Grossi Mazzorin, J., 2015. Fondo Paviani e Frattesina: economia animale di due central places della tarda Età del bronzo veneta. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 389-400.
- De Marinis, R.C. and Salzani, L., 1997. Le necropoli del Bronzo Medio e Recente nella Lombardia orientale e nel Veneto occidentale. In: M. Bernabò Brea, A. Cardarelli, M. Cremaschi, eds., *Le Terramare: la più antica civiltà padana*. Catalogo della mostra di Modena. Milano: Mondadori Electa, 703-719.
- Deiana, R., Vicenzutto, D., Deidda, G.P., Boaga, J., Cupitò, M., 2020. Remote sensing, archaeological, and geophysical data to study the Terramare settlements: the case study of Fondo Paviani (northern Italy). *Remote Sensing*, 12, 2617.
- Dubois, N. and Jacob, J., 2016. Molecular biomarkers of anthropic impacts in natural archives: a review. *Frontiers of Ecology and Evolution*, 4, 92. <https://doi.org/10.3389/fevo.2016.00092>
- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Klooß, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A., Mueller-Bieniek, A., Nisbet, R., Pashkevich, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H.P., Stobbe, A., Tolar, T., Wasylkowa, K., Wiethold, J., Zerl, T., 2020. New AMS ¹⁴C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 1-18. <https://doi.org/10.1038/s41598-020-70495-z>
- Hermes, T.R., Frachetti, M.D., Doumani Dupuy, P.N., Mar'yashev, A., Nebel, A., Makarewicz, C.A., 2019. Early integration of pastoralism and millet cultivation in Bronze Age Eurasia. *Proceedings of the Royal Society B: Biological Sciences*, 286 (1910), 20191273. <https://doi.org/10.1098/rspb.2019.1273>
- Heron, C., Shoda, S., Breu Barcons, A., Czebreszuk, J., Eley, Y., Gorton, M., Kirleis, W., Kneisel, J., Lucquin, A., Müller, J., Nishida, Y., Son, J., Craig, O.E., 2016. First molecular and isotopic evidence of millet processing in prehistoric pottery vessels. *Scientific Reports*, 6, 38767. <https://doi.org/10.1038/srep38767>
- Hunt, H.V., Vander Linden, M., Liu, X., Motuzaite-Matuzeviciute, G., Colledge, S., Jones, M.K., 2008. Millets across Eurasia: chronology and context of early records of the genera *Panicum* and *Setaria* from archaeological sites in the Old World. *Vegetation History and Archaeobotany*, 17, S5-S18. <https://doi.org/10.1007/s00334-008-0187-1>
- Jacob, J., Disnar, J.-R., Arnaud, F., Chapron, E., Debret, M., Lallier-Vergès, E., Desmet, M., Revel-Rolland, M., 2008. Millet cultivation history in the French Alps as evidenced by a sedimentary molecule. *Journal of Archaeological Science*, 35, 814-820. <https://doi.org/10.1016/j.jas.2007.06.006>
- Jacob, J., Disnar, J.R., Arnaud, F., Gauthier, E., Billaud, Y., Chapron, E., Bardoux, G., 2009. Impacts of new agricultural practices on soil erosion during the Bronze Age in the French Prealps. *The Holocene*, 19, 241-249. <https://doi.org/10.1177/0959683608100568>
- Jones, R., Levi, S.T., Bettelli, M., Vagnetti, L., eds., 2014. Italo-Mycenaean pottery: the archaeological and archaeometric dimensions. Roma: CNR – Istituto di Studi sul Mediterraneo antico.
- Kirleis, W., 2019. *Atlas of Neolithic Plant Remains from Northern Central Europe*. Groningen: Barkhuis Publishing.
- Leonardi, G., Cupitò, M., Baioni, M., Longhi, C., 2015. Northern Italy around 2200 cal BC – From Copper Age to Early Bronze Age: continuity and/or discontinuity? In: H. Meller, H.W. Arz, R. Jung, R. Risch, eds., *2200 BC – Ein Klimasturz*

- als Ursache für den Zerfall der Alten Welt?/2200 BC – A climatic breakdown as the cause for the collapse of the Old World? Tagungen des Landesmuseums für Vorgeschichte Halle 12/I. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte, 283-304.
- Lightfoot, E., Liu, X., Jones, M.K., 2013. Why move starchy cereals? A review of the isotopic evidence for prehistoric millet consumption across Eurasia. *World Archaeology*, 45 (4), 574-623. <https://doi.org/10.1080/00438243.2013.852070>
- Liu, X., Hunt, H.V., Jones, M.K., 2009. River valleys and foothills: changing archaeological perceptions of north China's earliest farms. *Antiquity*, 83, 82-95. <https://doi.org/10.1017/S0003598X00098100>
- Liu, X., Jones, M.K., Zhao, Z., Liu, G., O'Connell, T.C., 2012. The earliest evidence of millet as a staple crop: new light on Neolithic foodways in north China. *American Journal of Physical Anthropology*, 149, 238-290. <https://doi.org/10.1002/ajpa.22127>
- Liu, X., Jones, P.J., Motuzaite Matuzeviciute, G., Hunt, H.V., Lister, D.L., An, T., Przelomska, N., Kneale, C.J., Zhao, Z., Jones, M.K., 2019. From ecological opportunism to multi-cropping: mapping food globalisation in prehistory. *Quaternary Science Reviews*, 206, 21-28. <https://doi.org/10.1016/j.quascirev.2018.12.017>
- Lu, H., Zhang, J., Wu, N., Liu, K.B., Xu, D., Li, Q., 2009. Phytoliths analysis for the discrimination of foxtail millet (*Setaria italica*) and common millet (*Panicum miliaceum*). *PLoS ONE*, 4 (2), e4448. <https://doi.org/10.1371/journal.pone.0004448>
- Marcon, S. and Mazzoli, C., 2015. Le forme di fusione di Fondo Paviani (Verona) – Scavi Università di Padova 2007-2012: caratterizzazione tipologica e petrografica. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 845-848.
- Mercuri, A.M., Accorsi, C.A., Mazzanti, M., Bosi, G., Trevisan Grandi, G., Cardarelli, A., Labate, D., Olmi, L., Torri, P., 2006. Cereal fields from the Middle-Recent Bronze Age, as found in the terramara di Montale, in the Po Plain (Emilia Romagna, northern Italy), based on pollen, seeds/fruits and microcharcoals. In: J.P. Morel, J. Tresserras, J. Carlos Matamala, eds., *The Archaeology of Crop Fields and Gardens*. Bari: Edipuglia, 250-270.
- Mercuri, A.M., Sadori, L., Uzquiano Ollero, P., 2011. Mediterranean and north-African cultural adaptations to mid-Holocene environmental and climatic changes. *The Holocene*, 21, 189-206. <https://doi.org/10.1177/0959683610377532>
- Miller, N.F., Spengler, R.N., Frachetti, M., 2016. Millet cultivation across Eurasia: origins, spread, and the influence of seasonal climate. *The Holocene*, 26, 1566-1575. <https://doi.org/10.1177/0959683616641742>
- Motuzaite-Matuzeviciute, G., Staff, R.A., Hunt, H.V., Liu, X., Jones, M.K., 2013. The early chronology of broomcorn millet (*Panicum miliaceum*) in Europe. *Antiquity*, 87, 1073-1085. <https://doi.org/10.1017/S0003598X00049875>
- Motuzaite-Matuzeviciute, G., Jacob, J., Telizhenko, S., Jones, M.K., 2016. Miliacin in palaeosols from an Early Iron Age in Ukraine reveal in situ cultivation of broomcorn millet. *Archaeological and Anthropological Sciences*, 8, 43-50. <https://doi.org/10.1007/s12520-013-0142-7>
- Nicosia, C., Balista, C., Cupitò, M., Ertani, A., Leonardi, G., Nardi, S., Vidale, M., 2011. Anthropogenic deposits from the Bronze Age site of Fondo Paviani (Verona, Italy): pedomorphic and micropedological characteristics. *Quaternary International*, 243 (2), 280-292. <https://doi.org/10.1016/j.quaint.2010.10.006>
- Nisbet, R. and Rottoli, M., 1997. Le analisi dei macroresti vegetali dei siti dell'età del Bronzo. In: M. Bernabò Brea, A. Cardarelli, M. Cremaschi, eds., *Le Terramare: la più antica civiltà padana*. Catalogo della mostra di Modena. Milano: Mondadori Electa, 745-753.
- Ohmoto, T., Ikuse, M., Natori, S., 1970. Triterpenoids of the Gramineae. *Phytochemistry*, 9, 2137-2148. [https://doi.org/10.1016/S0031-9422\(00\)85379-0](https://doi.org/10.1016/S0031-9422(00)85379-0)

- Out, W.A. and Madella, M., 2016. Morphometric distinction between bilobate phytoliths from *Panicum miliaceum* and *Setaria italica* leaves. *Archaeological and Anthropological Sciences*, 8 (3), 505-521. <https://doi.org/10.1007/s12520-015-0235-6>
- Perego, R., 2015. *Contribution to the Development of the Bronze Age Plant Economy in the Surrounding of the Alps: an Archaeobotanical Case Study of two Early and Middle Bronze Age Sites in Northern Italy (Lake Garda Region)*. Doctoral dissertation, University of Basel.
- Pyankov, V.I., Ziegler, H., Akhani, H., Deigle, C., Luetge, U., 2010. European plants with C4 photosynthesis: geographical and taxonomic distribution and relations to climate parameters. *Botanical Journal of the Linnean Society*, 163 (3), 283-304. <https://doi.org/10.1111/j.1095-8339.2010.01062.x>
- Ravazzi, C., Marchetti, M., Zanon, M., Perego, R., Quirino, T., Deaddis, M., De Amicis, M., Margaritora, D., 2013. Lake evolution and landscape history in the lower Mincio River valley, unravelling drainage changes in the central Po Plain (N-Italy) since the Bronze Age. *Quaternary International*, 288, 195-205. <https://doi.org/10.1016/j.quaint.2011.11.031>
- Rottoli, M., 2001. Analisi archeobotaniche: i macroresti vegetali. In: Frontini P., ed., Castellarò del Vhò: campagne di scavo 1996-1999. Scavi delle Civiche Raccolte Archeologiche di Milano. Milano: Comune di Milano settore cultura musei e mostre raccolte archeologiche e numismatiche, 175-195.
- Rottoli, M. and Castiglioni, E., 2009a. Prehistory of plant growing and collecting in northern Italy, based on seed remains from the early Neolithic to the Chalcolithic (c. 5600-2100 cal BC). *Vegetation History and Archaeobotany*, 18 (1), 91-103. <https://doi.org/10.1007/s00334-007-0139-1>
- Rottoli, M. and Castiglioni, E., 2009b. Indagini sui resti vegetali macroscopici. In: M. Bernabò Brea and M. Cremaschi, eds., *Acqua e civiltà nelle Terramare: la vasca votiva di Noceto*. Milano: Skira, 152-163.
- Rottoli, M. and Pessina, A., 2016. Neolithic agriculture in Italy: an update of archaeobotanical data with particular emphasis on northern settlements. In: S. Colledge and J. Conolly, eds., *The Origins and Spread of Domestic Plants in Southwest Asia and Europe*. Walnut Creek: Left Coast Press, 157-170.
- Simonneau, A., Doyen, E., Chapron, E., Millet, L., Vannièr, B., Di Giovanni, C., Bossard, N., Tachikawa, K., Bard, E., Albéric, P., Desmet, M., Roux, G., Lajeunesse, P., Berger, J.F., Arnaud, F., 2013. Holocene land-use evolution and associated soil erosion in the French Prealps inferred from Lake Paladru sediments and archaeological evidences. *Journal of Archaeological Science*, 40 (4), 1636-1645. <https://doi.org/10.1016/j.jas.2012.12.002>
- Smetanina, O.F., Kuznetsova, T.A., Denisenko, V.A., Pivkin, M.V., Khudyakova, Yu. V., Gerasimenko, A.V., Popov, D. Yu., Il'in, S.G., Elyakov, G.B., 2001. 3 β -Methoxyolean-18-ene (miliacin) from the marine fungus *Chaetomium olivaceum*. *Russian Chemical Bulletin*, 50, 2463-2465. <https://doi.org/10.1023/A:1015068520694>
- Stika, H.P. and Heiss, A.G., 2013. Plant cultivation in the Bronze Age. In: A. Harding and H. Fokkens, eds., *The Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 348-369.
- Strafella, A., Cupitò, M., Angelini, I., Vidale, M., 2015. Le ambre di Fondo Paviani (Verona) – Scavi Università di Padova 2007-2012: inquadramento tipocronologico, analisi archeometriche e analisi paleotecnologica. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 849-854.
- Tafuri, M.A., Craig, O.E., Canci, A., 2009. Stable isotope evidence for the consumption of millet and other plants in Bronze Age Italy. *American Journal of Physical Anthropology*, 139 (2), 146-153. <https://doi.org/10.1002/ajpa.20955>
- Tafuri, M.A., Rottoli, M., Cupitò, M., Pulcini, M.L., Tasca, G., Carrara, N., Bonfanti, F., Salzani, L., Canci, A., 2018. Estimating C₄ plant consumption in Bronze Age north-

- eastern Italy through stable carbon and nitrogen isotopes in bone collagen. *International Journal of Osteoarchaeology*, 28, 131-142. <https://doi.org/10.1002/oa.2639>
- Tran, G., 2015. Proso millet (*Panicum miliaceum*), grain. Feedipedia, a programme by INRAE, CIRAD, AFZ and FAO. <https://www.feedipedia.org/node/722>. [Last updated on October 2, 2015, 16:16]
- Tran, G., 2017. Proso millet (*Panicum miliaceum*), forage. Feedipedia, a programme by INRAE, CIRAD, AFZ and FAO. <https://www.feedipedia.org/node/409>.
- Varalli, A., Moggi-Cecchi, J., Dori, I., Boccone, S., Bortoluzzi, S., Salzani, P., Tafuri, M.A., 2016. Dietary continuity vs. discontinuity in Bronze Age Italy. The isotopic evidence from Arano di Cellore (Illasi, Verona, Italy). *Journal of Archaeological Science: Reports*, 7, 104-113. <https://doi.org/10.1016/j.jasrep.2016.03.047>
- Ventresca Miller, A.R., Spengler, R., Haruda, A., Miller, B., Wilkin, S., Robinson, S., Roberts, P., Boivin, N., 2020. Ecosystem engineering among ancient pastoralists in northern central Asia. *Frontiers in Earth Science*, 8, 168. <https://doi.org/10.3389/feart.2020.00168>
- Vicenzutto, D., Dalla Longa, E., Angelini, I., Artioli, G., Nimis, P., 2015. I manufatti in bronzo del sito arginato di Fondo Paviani (Verona) – scavi Università di Padova 2007-2012: inquadramento tipocronologico e analisi archeometriche. In: G. Leonardi and V. Tinè, eds., *Preistoria e protostoria del Veneto*. Studi di Preistoria e Protostoria 2. Firenze: Istituto Italiano di Preistoria e Protostoria, 833-838.
- Zanon, M., Unkel, I., Andersen, N., Kirleis, W., 2019. Palaeoenvironmental dynamics at the southern Alpine foothills between the Neolithic and the Bronze Age onset: a multi-proxy study from Bande di Cavriana (Mantua, Italy). *Quaternary Science Reviews*, 221, 105891. <https://doi.org/10.1016/j.quascirev.2019.105891>
- Zohary, D., Hopf, M., Weiss, E., 2012. *Domestication of Plants in the Old World: the Origin and Spread of Domesticated Plants in Southwest Asia, Europe, and the Mediterranean Basin*. (4th edition). Oxford: Oxford University Press.

Exploring seed impressions within the fabric of pottery: Using a silicone cast method for reliable identification

Eiko Endo

Abstract

This paper presents an improved method to analyse seed impressions in pottery and describes its effectiveness for distinguishing millets. It may be especially helpful to distinguish grains of millet cultivars, such as foxtail (*Setaria italica*) and broomcorn (*Panicum miliaceum*) millet. This method has rapidly become popular across the Japanese archipelago over the past decade and has strongly improved knowledge about plant utilisation in ancient times. For example, it has revealed the exact timing of the acceptance of crop cultivation during the Jōmon-Yayoi culture transitional period and the diversity in the development of Yayoi agriculture in various parts of the archipelago as a result of differences in social organisation and ecological environments. Based on these achievements, the method represents a valuable alternative to flotation-based studies in the field of archaeobotany. One of the important advantages of the study of seed impressions is that we can get information on plants even from archaeological sites where flotation was not performed. In addition, it can provide direct links between pottery culture and plant use, which are more complicated to explore based on charred seed assemblages. However, due to their complex formation processes, there has been no agreement about how to interpret plant impressions in pottery in relation to human subsistence economies. Therefore, this chapter discusses potential formation processes of plant impressions based on an extensive dataset from the Japanese archipelago.

Introduction

This paper introduces a method to analyse plant impressions (mostly seeds) in pottery and outlines its value for distinguishing grains of different millet species. There is a long history of research concerned with pottery impressions in the study of plant use by prehistoric and early historic cultures. However, nowadays, many archaeobotanists believe that the identification of impressions is often ambiguous and thus are sceptical

about impression-based botanical datasets (Zohary and Hopf 2000, 5; Hunt *et al.* 2008; Stevens *et al.* 2016, 1545). In addition, due to plant impressions' complex formation processes, quantitative analysis of impressions is limited (Willcox and Fornite 1999; McClatchie and Fuller 2014). Some scholars have pointed out that it is possible to discuss activities such as crop processing (*e.g.* threshing) or pottery production by using the impressions of temper (Hovsepyan and Willcox 2008; McClatchie and Fuller 2014; Moskal-del Hoyo *et al.* 2017). However, this paper focuses on seed rather than temper impressions. Although it may be difficult to identify impressions by naked eye and to analyse the data in a quantitative way, impressions are informative in terms of plant use in everyday life, and they contribute to understanding the relationship between humans and plants.

Recently, several improvements in the analytical methodology of plant impression studies have been achieved (*e.g.* Takase 2011; Arobba *et al.* 2017). One innovation, which the present author has been applying successfully, is the analysis of casts made by means of silicone fillings. Another method, using materials other than silicone, is also practised; however, the author considers silicone to be an excellent material due to its transfer function. This method allows reliable identification of imprints to species level if they were formed under appropriate conditions. It may be especially helpful to distinguish different millet species, such as foxtail (*Setaria italica*) and broomcorn (*Panicum miliaceum*) millet, which is often difficult for charred, and thus sometimes deformed, millet grain assemblages due to overlaps in morphology and size (Spengler *et al.* 2013). By analysing the surface structure of the casts, more reliable plant taxonomic identifications are possible.

Refined silicone cast method

The silicone cast method adopted by the author was introduced by Ushino and Tagawa (1991) and improved by Hisa and Katada (2005) to reduce working time in order to facilitate analysis of a large amount of material. Positive casts are made from impressions by means of silicone and are then analysed using scanning electron microscopy (SEM). Thanks to the high-resolution transcription of clay, morphological details of the plant surface can be observed at 500× magnification or higher. Grikpedis (2019), who used this technique in Belarus, noted that the application of SEM to investigate silicone casts of plant impressions facilitates inspection and evaluation not only of the shape and size of imprints, but also their surface elements and, therefore, allows for secure identification of the impressed objects.

Procedure

The detailed procedure is as follows (see Fig. 1):

- a. Search for impressions on the inner and outer surface and cross-section of pottery fragments by naked eye and with the use of a magnifying glass. Pay attention to those impressions that are not fully exposed, because most of their shape is located inside the pottery fabric and is accessible only through a small hole in the surface (Fig. 1A); thanks to the elasticity of silicone, it is possible to also take casts from such deep and overhanging impressions, without destroying the pottery.
- b. Clean the impression to remove soil, using a soft brush.
- c. Coat the impression and the surrounding part of the pottery with acetone containing 5% Paraloid B72 to protect the pottery surface (Fig. 1B).
- d. Fill silicone into the impression (Fig. 1C).
- e. Remove the cast from the impression after hardening (Fig. 1D).
- f. Remove the coating liquid to clean the surface, using 100% acetone.
- g. Analyse the cast using SEM (Fig. 1F).




Figure 1. Main working steps of the seed impression casting method.




Figure 2. Location of impressions of 1451 rice, foxtail millet and broomcorn millet seeds in a combined assemblage of Jōmon-, Yayoi- and Kofun-type pottery.

Processes of formation of plant impressions

As mentioned above, quantitative analyses of plant impressions data have generally not been accepted due to the diverse formation processes that are potentially involved. In the following sections, the author discusses the potential processes of formation of plant impressions based on an extensive dataset from the Japanese archipelago covering the Jōmon (15,000-2500 cal BP) (Noshiro and Sasaki 2014), Yayoi (2500 cal BP-250 CE) and Kofun (250-710 CE) (Mizoguchi 2013) periods, represented by 2951 impressions from 166 archaeological sites.

What was the condition of the pottery when the seeds became impressed within the fabric?

Impressions of crops (*Oryza sativa*, $n=737$; *Setaria italica*, $n=425$; *Panicum miliaceum*, $n=289$) were identified on the inner and outer surfaces, flat base and cross-section of pottery (Fig. 2). They are distributed on various parts of pottery; therefore, crops were likely incorporated into the fabric in the clay state before the pottery obtained its final shape. That impressions were more often detected on the outer surface than the inner surface may be due to the difficulty of observing the inner surface on, for example, vessels with bottlenecks. However, regarding impressions on the flat base of the pottery, it is possible to infer that, here, the incorporation happened after the pot was formed and before the fabric dried (*i.e.* hardened). The impressions of such formation process may be described as impressions ‘on’ pottery.

Where did the seeds become incorporated into the pottery?

In the assemblages analysed, most of the impressions identified to species or genus level were cultivated cereals. The relatively few remaining specimens comprise other cultivated plants or wild plants (Table 1). Given the dominance of crop plants and the low abundance of wild plants, the author presumes that the preparation of the fabric was done inside dwellings, where the crops were stored and cooked. The predominance of crop plants in flotation samples from burnt dwellings from settlements of the Yayoi and succeeding cultures also suggests that crops were stored inside dwellings (e.g. Takase 2004; Takase and Endo 2010). In addition, impressions of household insect pests, such as the maize weevil (Obata *et al.* 2019), support the assumption that fabric for pottery was prepared indoors.

The conditions and preserved features of the identified crops

The dominant features of the crops that are preserved are the caryopsis with lemma and palea (Table 2). Therefore, the author suggests that most seeds were incorporated into fabric during the final stage of dehusking. Furthermore, given the presence of husks of *Oryza sativa*, it seems likely that threshing was done indoors.

Did the grains become incorporated into the fabric intentionally or accidentally?

Based on examples of large amounts of impressions in pottery, some researchers have argued that seeds were sometimes mixed into fabric intentionally. For example, Obata (2016) has identified a large fragment of Jōmon pottery with 526 impressions of perilla (*Perilla frutescens*) using soft X-ray instruments. He argues that this cannot occur without a strong intention to mix perilla with fabric. Such evidence has also occasionally been recorded by the author, raising questions about if/how seeds that were intentionally mixed into clay would have increased the risk of damage during firing of the pot or would have limited the pot's functionality (Fig. 3). The author believes it is debatable whether seeds were intentionally mixed into the fabric or were incorporated by chance. The origin of impressions of crop processing by-products is clearer. If they are detected in large numbers on the entire surface of pottery or daub, it is most likely that they were admixed with the fabric as a temper.

Current results from the study of plant impressions

Due to the lack of charred plant remains obtained through flotation, or the data on them, the replica method has become popular across the Japanese archipelago over the past decade. It has helped reconstruct and discuss plant use in prehistoric times since the impressions in pottery have revealed a diverse utilisation of plants and the advent of agriculture during the Jōmon-Yayoi transitional period. For instance, regarding the use of legumes during the Jōmon period, seed impression studies have demonstrated that continuous utilisation of these plants started in the first half of the Early Jōmon period and that an increase in the seed size started in the Middle Jōmon period (Nakayama 2020). Concerning the spread of agriculture, seed impressions studies have revealed that the crop package of *Oryza sativa*, *Setaria italica* and *Panicum miliaceum* arrived in western Japan at the end of the Final Jōmon

Taxon	Count
<i>Oryza sativa</i>	737
cf. <i>Oryza sativa</i>	122
<i>Setaria italica</i>	425
cf. <i>Setaria italica</i>	49
<i>Panicum miliaceum</i>	289
cf. <i>Panicum miliaceum</i>	75
<i>Setaria viridis</i>	14
Paniceae	12
<i>Perilla</i>	19
cf. <i>Perilla</i>	11
<i>Cannabis</i>	4
<i>Lactuca raddeana</i> var. <i>elata</i>	1
<i>Carpinus tschonoskii</i>	1
cf. <i>Echinochloa</i>	2
<i>Sambucus</i>	2
Fabaceae	2
millipede	1
maize weevil	1
insect	6
shell	3
Total	1776

Table 1. Total counts of identified impressions in Jōmon-, Yayoi- and Kofun-type pottery.

Taxon	Caryopsis	Floret (caryopsis with lemma and palea)	Spikelet	Husk (chaff)
<i>Oryza sativa</i>	64	664		8
<i>Setaria italica</i>	11	14		
<i>Panicum miliaceum</i>	4	282	3	

Table 2. Parts of cereal fruits recognised in the silicone casts of impressions.



Figure 3. A lid with impressions of *Oryza sativa* ($n=3$), *Setaria italica* ($n=21$) and *Panicum miliaceum* ($n=12$) recovered from the Daizuda site (Late Yayoi period), in Nagano Prefecture, Japan.

period and that, in the succeeding Yayoi period, a diverse agriculture developed in various parts of the archipelago as a result of differences in social organisation and ecological environments (Endo 2015). In the western part of the archipelago, *O. sativa* was dominant and accompanied by a small amount of millets. In the Central Highlands, one of the most developed areas of the Jōmon culture, only millet was selected in the first half of the Yayoi period.

Although it is difficult to examine plant impressions data quantitatively, the presence/absence data allow researchers to analyse the relationship between pottery types and cultivated plants. For example, impressions on the Ongagawa-type pottery, which was distributed across western Japan in the Early Yayoi period and has been in Japanese archaeology used as a chronological marker of the start of agriculture, are mainly *O. sativa* accompanied by only little millet. On the other hand, almost all of the impressions on Fusenmon-type pottery, which was distributed across the Central Highlands and is synchronous with Ongagawa-type pottery, are from millets. In addition, during the Middle Yayoi period in the Kanto region in eastern Japan, *O. sativa* dominates in the Miyanodai-type pottery used in the southern part of the Kanto region, while *O. sativa* and millets are mixed in the Kitajima-type pottery used in the northern part of the region (Table 3).

A combination of systematic flotation and use of the silicone-casting method at the Sunadadai site in southern Kanto revealed more than 2500 charred grains of *O. sativa*, but none of millet (Kanagawa Prefecture Buried Cultural Heritage Research Centre [KPBCHR] 1991). It also revealed impressions of eight *O. sativa* caryopses and one *S. italica* grain (Endo 2019). Similarly, the combination of impressions in pottery of rice and millets (Endo 2019) is consistent with the recent results of flotation at Maenakanishi, a site located in northern Kanto, at which mainly Kitajima-type pottery has been recorded (Leipe *et al.* 2020).

Recently, the presented method has been also successfully applied in regions outside Japan, such as the Russian Far East (Fukuda *et al.* 2019), South Korea (Son *et al.* 2010; Nakayama 2014), southern India (Nakayama and Uesugi 2015), Ukraine (Endo 2019) and Belarus (Grikpedis *et al.* 2018; Grikpedis 2019) (Fig. 4).

Differentiation between *Setaria italica* and *Panicum miliaceum*

One of the most important advantages of the method is the opportunity to obtain information on the spectrum of the plants used at archaeological sites where archaeobotanical recovery (*e.g.* flotation) is otherwise not performed or where the preservation of botanical materials is very poor. Even where plant remains have been preserved and recovered, the differentiation of small seeds, particularly those that are morphologically similar, such as grains of millets, can be difficult because charring/carbonisation causes deformations. Silicone casts of impressions are replicas of the fresh (uncarbonised) seed, exhibiting all morphological details. Furthermore, most cereal grains were still covered with lemma and palea when they became incorporated into the fabric, whereas most charred specimens are without lemma and palea. Thus, in silicone casts of impressions, more details of the original material are preserved, which allows a more reliable identification. For distinguishing *Setaria italica* and *Panicum miliaceum* caryopses, the papillae distribution on lemma and palea of *S. italica* is one of the basic criteria. Another criterion is the smooth surface, without papillae, of *S. italica* grains, which is observed only on the lemma and, more specifically, on the part that overlaps with the palea (Fig. 5A). The lemma and palea of *P. miliaceum* have no such papillae and the surface is overall smooth and glossy (Fig. 5B). In addition, it has been shown that *S. italica* is distinguishable from other *Setaria* species, such as its progenitor *S. viridis*, based on the non-ridged base of the papillae on the upper lemma of *S. italica* (Nasu *et al.* 2007). This feature can be also observed in high-quality SEM images of silicone casts (Fig. 5C).

Period	Pottery type	Region	site	Number of observed pottery shards	Number of shards with cereal impressions	<i>Oryza sativa</i>	<i>Setaria italica</i>	<i>Panicum miliaceum</i>
Early Yayoi	Ongagawa	Western Japan	Nishisiga	20,819	95	48		1
			Tamura	423	5	10		
	Fusenmon	Central Highland	Ishigyo	360	47	1	60	26
			Yazaki	132	22		41	2
			Gongendomae		15		5	
			Ohsyuku	182	8		13	2
			Kitakatakitanohara	133	5		4	19
			Fukayamada	42	2		5	9
Middle Yayoi	Miyanodai	Southern Kanto	Ikego	1249	36	30	2	5
			Asukayama	1536	9	9		
	Kitajima	Northern Kanto	Kitajima	1500	24	10	20	1
			Maenakanishi	2373	42	20	24	




Table 3. Identified crop impressions in pottery organised by cultural period, pottery typology and location.

Figure 4. SEM images of *Panicum miliaceum* grain impressions from selected sites in Eurasia: (A) Ishigyo (Nagano Prefecture, Japan); the pottery fragments with impressions are typologically dated to the first quarter of the 1st millennium BCE; (B) Novokiyivka (Ukraine), typologically dated to 1600-1300/1200 BCE; (C) Kamen 6 (Belarus), typologically dated to the 3rd and 2nd millennium BCE; (D) Zhertyj Yar (Russian Far East), typologically dated to the 3rd century BCE-4th century CE.




Figure 5. SEM images of *Setaria italica* and *Panicum miliaceum* grains: (A) *S. italica* with papillae on the lemma and palea; (B) the smooth and glossy surface on the lemma and palea of *P. miliaceum*; (C) the non-ridged base of the papillae on the upper lemma of *S. italica*.

Conclusions

This paper summarises an improved method to study seed impressions in pottery and describes its advantages for identifying small seeds. Seed impressions studies can substantially contribute to our understanding of the relationship between humans and plants in prehistoric times, thus representing a valuable alternative or a complement to flotation-based studies in the archaeobotany. Especially, such studies can provide direct links between pottery making and plant use, which are more complicated to explore based on charred seed assemblages. Further experimental and ethnoarchaeological research is needed to improve our understanding of the formation processes and taphonomy of seed impressions and to allow a more quantitative interpretation of the obtained datasets.

Acknowledgements

The author would like to thank C. Leipe, D. Filipović and the anonymous reviewer for valuable comments that helped to improve the paper and for the English-language editing. This work was supported financially by the JSPS KAKENHI Grant-in-Aid for Scientific Research (C) entitled ‘The timing and route of the spread of East Asia originated millets (*Panicum miliaceum*, *Setaria italica*) to the western steppe’, Grant number 19K01116.

References

- Arobba, D., Panelli, C., Caramiello, R., Gabriele, M., Maggi, R., 2017. Cereal remains, plant impressions and ^{14}C direct dating from the Neolithic pottery of Arene Candide Cave (Finale Ligure, NW Italy). *Journal of Archaeological Science: Reports*, 12, 395-404. <https://doi.org/10.1016/j.jasrep.2017.02.015>
- Endo, E., 2019. (in Japanese) [Yayoi agriculture in the Kanto region]. In: H. Shitara, ed., (in Japanese) [The archaeology of agricultural complex formation: the beginning of farming]. Tokyo: Yuzankaku, 111-125.
- Endo, E., 2015. (in Japanese) [*The start of agriculture in the Japanese archipelago based on cultivated crops*]. Doctoral thesis, Tokyo Metropolitan University. https://tokyo-metro-u.repo.nii.ac.jp/?action=pages_view_main&active_action=repository_view_main_item_detail&item_id=4975&item_no=1&page_id=30&block_id=164
- Endo, E., Nasu, H., Gaskevych, D., Yanevich, A., Pashkevich, G., Videiko, M., 2019. *Ukraine as the Crossroad for Agricultural Dispersal in Eurasia*. Oral presentation at the 18th Conference of the International Work Group for Palaeoethnobotany, 3-8 June 2019, Lecce, Italy.
- Fukuda, M., Kunikita, D., Endo, E., Gorshkov, M., Nasu, H., Kitano, H., 2019. (in Japanese) [Crop utilization and diet in the Pol'tse culture]. In: H. Shitara, ed., (in Japanese) [*Agricultural culture complex*]. Tokyo: Yuzankaku, 71-90.
- Grikpedis, M., 2019. *In the Search of the Earliest Cultivated Plants in Belarus: SEM Analysis of Grain Imprints in Pottery*. Oral presentation at the 18th Conference of the International Work Group for Palaeoethnobotany, 3-8 June 2019, Lecce, Italy.
- Grikpedis, M., Endo, E., Motuzaite Matuzeviciute, G., Kryvaltsevich, M., Tkachova, M., 2018. Plants in pots: SEM research of ceramic silicon casts from the River Prypiat Basin. In: O.V. Lozovskaya, A.A. Vyibornov, E.V. Dolbunova, eds., *Subsistence Strategies in the Stone Age, Direct and Indirect Evidence of Fishing and Gathering. Materials of the international conference dedicated to the 50th anniversary of Vladimir Mikhailovich Lozovski, 15-18 May 2018, Saint Petersburg*. St. Petersburg: ИИМК РАН, 210-213.
- Hisa, Y. and Katada, M., 2005. (in Japanese) [*A technique for making models of impressions on pottery by replication method (trial edition)*]. Fukuoka City: Fukuoka Prefecture Buried Cultural Heritage Research Centre.

- Hovsepian, R. and Willcox, G., 2008. The earliest finds of cultivated plants in Armenia: evidence from charred remains and crop processing residues in pisé from the Neolithic settlements of Aratashen and Aknashen. *Vegetation History and Archaeobotany*, 17, 563-571.
- Hunt, H.V., Linden, M.V., Liu, X., Motuzaite-Matuzeviciute, G., College, S., Jones, M.K., 2008. Millets across Eurasia: chronology and context of early records of the genera *Panicum* and *Setaria* from archaeological sites in the Old World. *Vegetation History and Archaeobotany*, 17, S5-S18. <https://doi.org/10.1007/s00334-008-0187-1>
- Kanagawa Prefecture Buried Cultural Heritage Research Centre (KPBCHR), 1991. *Sunadadai site II*. Yokohama: Kanagawa Prefecture Buried Cultural Heritage Research Centre.
- Leipe, C., Endo, E., Kuramochi, S., Wagner, M., Tarasov, P., 2020. Crop cultivation of Middle Yayoi culture communities (fourth century BCE-first century CE) in the Kanto region, eastern Japan, inferred from a radiocarbon-dated archaeobotanical record. *Vegetation History and Archaeobotany*, 30, 409-421. <https://doi.org/10.1007/s00334-020-00791-1>
- McClatchie, M. and Fuller, D.Q., 2014. Leaving a lasting impression: arable economies and cereal impressions in Africa and Europe. In: C.J. Stevens, S. Nixon, M.A. Murray, D.Q. Fuller, eds., *Archaeology of African Plant Use*. Walnut Creek: Left Coast Press, 259-265.
- Mizoguchi, K., 2013. *The Archaeology of Japan: from the Earliest Rice Farming Villages to the Rise of the State*. Cambridge: Cambridge University Press.
- Moskal-del Hoyo, M., Rauba-Bukowska, A., Lityńska-Zajac, M., Mueller-Bieniek, A., Czekaj-Zastawny, A., 2017. Plant materials used as temper in the oldest Neolithic pottery from south-eastern Poland. *Vegetation History and Archaeobotany*, 26, 329-344. <https://doi.org/10.1007/s00334-016-0595-6>
- Nakayama, S., 2020. (in Japanese) [Legumes and Jōmon people]. Tokyo: Douseisha.
- Nakayama, S., ed., 2014. (in Japanese) [*Origin of grain agriculture in the Japanese Archipelago and the Korean Peninsula*]. Yamanashi: Yamanashi Prefectural Museum.
- Nakayama, S. and Uesugi, A., 2015. (in Japanese) [*Research on cultivated plants in South Asia using the replica method*]. Oral presentation at the 20th Conference of the Japanese Society for West Asian Archaeology, University of Tsukuba, Tsukuba, Japan.
- Nasu, H., Momohara, A., Yasuda, Y., He, J., 2007. The occurrence and identification of *Setaria italica* (L.) P. Beauv. (foxtail millet) grains from the Chengtoushan site (ca. 5800 cal B.P.) in central China, with reference to the domestication centre in Asia. *Vegetation History and Archaeobotany*, 16, 481-494. <https://doi.org/10.1007/s00334-006-0068-4>
- Noshiro, S. and Sasaki, Y., 2014. Pre-agricultural management of plant resources during the Jomon period in Japan – a sophisticated subsistence system on plant resources. *Journal of Archaeological Science*, 42, 93-106. <https://doi.org/10.1016/j.jas.2013.11.001>
- Obata, H., 2016. (in Japanese) [Jōmon people sowing seeds]. Tokyo: Yoshikawa Kobunkan.
- Obata, H., Morimoto, K., Miyanoshita, A., 2019. Discovery of the Jomon era maize weevils in Hokkaido, Japan and its mean. *Journal of Archaeological Science: Reports*, 23, 137-156. <https://doi.org/10.1016/j.jasrep.2018.10.037>
- Son, J.-H., Nakamura, D., Momohara, A., 2010. (in Korean) [Research on plant impressions found on Bronze Age pottery by means of replication]. *Field Archaeology*, 8, 5-34.
- Spengler, R., Chang, C., Tourtellotte, P., 2013. Agricultural production in the central Asian mountains: Tuzusai, Kazakhstan (410-150 B.C.). *Journal of Field Archaeology*, 38, 68-79. <https://doi.org/10.1179/0093469012Z.000000000037>
- Stevens, C.J., Murphy, C., Roberts, R., Lucas, L., Silva, F., Fuller, D.Q. 2016. Between China and South Asia: a middle Asian corridor of crop dispersal and agricultural innovation in the Bronze Age. *The Holocene* 26, 1541-1555. <https://doi.org/10.1177%2F0959683616650268>

- Takase, K., 2004. (in Japanese) [Carbonized seeds from the Suginodo site in Mizusawa city, Iwate prefecture]. *In: (in Japanese) [The excavation report of Suginodo site].* Morioka: Iwate Prefecture Buried Cultural Heritage Research Center, 161-170.
- Takase, K., 2011. Plant seeds recovered from potsherds of the Final Jomon and Yayoi periods: a case study in Iwate and Yamagata prefectures, northeastern Japan. *Meiji University Ancient Studies of Japan*, 3, 41-63.
- Takase, K. and Endo, E., 2010. (in Japanese) [Analysis of carbonized seeds detected from house no. 21 of Yayoi period, no. 31 point of Tagoyama site, Shiki city, Saitama prefecture]. *Bulletin of the Research Institute for Japanese Ancient Studies 2010*, 3-13.
- Ushino, T. and Tagawa, H., 1991. (in Japanese) [Replication method of impressions on the pottery surface]. *Archaeology and Natural Science*, 24, 13-36.
- Willcox, G. and Fornite, S., 1999. Impressions of wild cereal in pisé from the 10th millennium uncal B.P. at Jerf el Ahmar and Mureybet: northern Syria. *Vegetation History and Archaeobotany*, 8, 21-24. <https://doi.org/10.1007/BF02042838>
- Zohary, D. and Hopf, M., 2000. *Domestication of Plants in the Old World* (1st edition). Oxford: Oxford University Press.

Traditional millet cultivation in the Iberian Peninsula: Ethnoarchaeological reflections through the lens of social relations and economic concerns

Andrés Teira-Brión

Abstract

In this paper, I propose a reassessment of some widespread archaeological claims about millet agrarian practices – double rotation and an increase in cereal production – based on conclusions derived from ethnographic analogy. The reflections presented here arose from experiences during fieldwork, which has transformed my perception of the material record of and historical sources on millet farming. The results of the ethnoarchaeological investigation, carried out within the project ‘New cultivars, new landscapes: Agriculture and anthropisation of the first farming societies in northern Iberia (2012-2014)’, have made it possible to recognise the different stages of the *chaîne opératoire* of millet production and processing, as well as millets’ uses in human food and animal feed.

Introduction

In a broad sense, ethnoarchaeology is

‘...the study of how material culture is produced, used and deposited by contemporary societies in relation to the wider social, ideological, economic, environmental and/or technical aspects of the society concerned, and with specific reference to the problems of interpreting archaeological material’ (Sillar 2000, 6).

Over the past two decades, researchers have explored the relevance and meaning of the material past within distinctive disciplinary traditions (Hamilakis and Anagnostopoulos 2009), including approaches to the materiality of ‘primitive’ societies within contemporary modernity (González-Ruibal 2014). Essentially,

however, it is still necessary to distinguish between ethnographic analogy and ethnoarchaeology. According to A. Hernando (1995), analogy is descriptive, establishing specific similarities between two cultures – present and past – under their respective cultural contexts; whereas ethnoarchaeology makes generalisations that aim to be laws, reviewing the ethnographic record in relation to a specific behaviour, process of change, *etc.*, in order to understand in which socio-economic, ideological or environmental conditions a particular model of conduct may appear.

Ethnoarchaeological studies have made important contributions to our knowledge about millet agriculture and processing, based on observation of traditional farmers (*e.g.* Hörander 1995; Lundström-Baudais *et al.* 2002; Hamon and Le Gall 2013; Song *et al.* 2013). These studies have the capacity to connect the pieces of the puzzle we recover from the past with current materials, but not in a direct analogy. The materiality we study at present is not static, immutable or timeless. It is a snapshot of a specific moment in a continuously changing process. The conducts we observe may have been established after the moment we aim to study or at different timescales; therefore, translating these practices into the archaeological record cannot be done uncritically. Indeed, conjectures about current agricultural practices may even transcend into the scientific literature as theories. To avoid this, ethnoarchaeology provides valuable information to reflect on the limitations of archaeological approaches.

A key question in the study of millet farming is what social changes it may have led to. In line with the results of ethnographic studies, two claims have been disseminated among archaeologists: a) short growth cycle of millets involve the production of two cereal harvest per year and, associated with this, b) millet cultivation means an increase in cereal surplus. But, at present, we hardly know how they were cultivated in western Europe by prehistoric communities. These ideas remain as a kind of research artefact towards arguing for a higher complexity and intensification of agricultural practices since the Bronze Age.

Although both proposals are feasible, their generalisation to any past society is problematic. Far from helping to establish the role of millets, they make it difficult to understand their multiple relationships within different agricultural systems. Even though these proposals may be attractive in terms of highlighting millets' potential, they remain speculative and often become stereotypes. In order to acquire agricultural knowledge, the information generated from a single archaeological discipline is often insufficient. Therefore, ethnoarchaeology exists not only to provide similarities or laws of conduct, but also to warn about the problem of transferring practices from the present without evaluating other aspects of the system in which they are enmeshed. An interdisciplinary and multiscale conception and definition of the social and economic contexts of agriculture can allow us to test the appropriateness of the above statements.

Farming millets in the Iberian Peninsula: A case study

Several studies on millets in the Iberian Peninsula have used an ethnographic or ethnoarchaeological approach (Vázquez Varela 1994; Barboff 1995). All these works are focused on the northwestern region, where traditional millet cultivation was carried out until recently. Between 2012 and 2014, I participated in a project led by Lydia Zapata, which made it possible to encounter farmers who were still growing millets or who had only recently abandoned this practice (Moreno-Larrazabal *et al.* 2015). The scope of the study integrated interviews in Spain – in the regions of Asturias and Galicia – and northern Portugal, mainly among elderly people, which allowed us to document farming practices going back to the 1920s. The aim



Figure 1. a) Aitor and Lydia during ethnographic fieldwork in Manuel's foxtail millet plot (Asturias, Spain); b) milling broomcorn millet using an electric mill (Galicia, Spain); c) Delia dehusking *P. miliaceum* with the pilão (Paredes de Coura, Portugal); d) mortar and pisón (pestle) from A Capela (A Coruña, Galicia) (photograph by Fernán López); e) mortar with different type of pestle from A Coruña province (Galicia). The author has the consent of the people depicted in this figure for the dissemination of their images.




Figure 2. a) distribution of *P. miliaceum* and *S. italica* in peasant traditional farming based on ethnobotanical fieldwork; b-d) different plant end uses, as products and by-products; e) double cereal harvest per year with a rotation of rye and broomcorn millet.

of the project was to recognise the *chaîne opératoire* and to determine criteria for interpreting the archaeobotanical evidence from millet products and by-products in relation to actions and tools. In establishing this analogy, the remains were identified for each stage of the processing according to the remains preserved in archaeobotanical assemblages. But at the same time, we reflected on whether contemporaneous millet-related practices have any connection to past ones, when changes in millet use or distribution occurred, and for what reason.

During the ethnographic study, we observed differences in agricultural practices (Fig. 1), primarily related to the species grown and their uses. In Galicia, Spain, and the Minho region of Portugal, broomcorn millet (*Panicum miliaceum*) is grown exclusively. The farmers interviewed did not recognise the foxtail millet (*Setaria italica*) plant. In all cases the grains were the main product, which were destined for two uses: for human food (porridge, pudding or bread) or for animal feed (chickens and hens were fed the whole grains, and cows were fed the flour after calving). The stems of the plants were used as fodder for cattle as a secondary use. Cultivation practices were fairly homogeneous. Sowing was done in June on the land where rye (*Secale cereale*) had been gathered and harvested the previous September or October. A double winter (rye) and spring (broomcorn millet) cereal harvest was therefore carried out in the same plot. Only farmers in the westernmost area of Lugo province documented a single annual rotation with other crops. The way the millet was processed showed one of the most significant differences. Milling of the grains without removing the husks was documented in western areas of the province of A Coruña. The resulting flour had a sort of sandy texture due to the inclusion of some husk fragments that had not been separated out during sieving. Here, although mortars were known, they were not associated with millet processing. Elsewhere in Galicia and Portugal, the grains were not ground, so the use of stone mortars was the usual way for dehusking. For dehusking, two different types of mortars and pestles were used, one in the most northern area (Fig. 1d-e) and another in the southern area (Fig. 1c).

Asturias is the region where we documented the cultivation of foxtail millet (Fig. 2), closely associated with the use of the plant as green fodder for livestock. This plant was very appreciated as fodder to fatten animals in the summer months, when pasture was scarce in the valleys. The grains had a secondary use for feeding the chickens. In no case was human consumption documented. It was common to buy seed grains for sowing at local markets, mainly the one held in the village of Cangas de Narcea. Only two populations of *P. miliaceum* were documented, in maslin plots with *S. italica*. However, farmers did not consider broomcorn millet as a crop, but as a companion plant of foxtail millet. It did not even have a different name.

Reconstructing past cultivation practices using historical records

All crops studied in 2012-2014 (Moreno-Larrazabal *et al.* 2015) were marginal products at the time of the fieldwork. They had survived within a traditional agrarian society that had kept them as a complement for human and animal alimentation. The practices documented can be understood as an example of the resilience of a rural society in the process of transformation towards a market economy. This is visible in the adaptation of processing to current tools. Some of them are easily identifiable, such as the use of the tractor for trampling *P. miliaceum* as a substitute for using cattle or flails. On other occasions, it is complex to identify the exact moment when a practice is being transformed, or even if there is a territorially differentiated behaviour. Grinding in electric mills is clearly a 20th century replacement for grinding in hydraulic mills. However, in certain areas, we do not know when the change to using the flour instead of the whole grains for the same recipes took place. Sometimes, within a homogeneous system of practices, such as the cultivation of *S.*

italica in Cangas de Narcea and Degaña, in Asturias, we find individual specificities. Wetting the grain before sowing was documented in only a few families. We found only one farmer who did not level the ground before sowing. These behaviours can be considered as personal choices but are also a notable source of information. Identifying all these variables is useful for observing the diversity of the practices carried out. They break the strict and uniform picture of crop cultivation. However, it is difficult to know at what point each of these adaptations took place.

The boundaries established by an ethnographic study area can give us a biased notion of the spatial distribution of cultivation. It is therefore important to trace for each behaviour whether it is a modern adaptation or has its roots in remote times. For this purpose, written and archaeological sources provide a multitemporal overview of the distribution of millets and their relationship with other crops within agricultural systems. The need to establish the wealth of countries in modern and contemporary times lay at the origin of national inventories made throughout Europe. One of these sources in the Spanish territory is the *Diccionario Geográfico-Estadístico-Histórico de España y sus posesiones de ultramar*, published by Pascual Madoz (1846-1850). He coordinated a survey to find out about the goods, products and economic activities of each village. The survey was sent to the entire country, with the exception of territories with a specific tax system, such as Navarra, Guipuzkoa or Bizkaia. Although the quality of the information is uneven, as it depended on each local administration, the inventory provides valuable information on the most important crops in each place and the produce supplied to the markets (Fig. 3). In the Madoz inventory, the words *mijo*, commonly used in Spanish for *Panicum miliaceum*, and *panizo*, used for *Setaria italica*, are recorded a total of 478 times (Fig. 3a). However, we cannot associate these words with millet cultivation since *mijo* and *panizo* may also refer to corn (*Zea mays*), as regional substitutes for the more usual name *maíz*. Separating out the populations where maize cultivation is not mentioned provides a more precise distribution of those cases where *maíz* was accompanied by the cultivation of *mijo* (215 times) and/or *panizo* (53 villages) (Fig. 3b). Incorporating this source of information significantly changes the distribution which had been observed in the ethnographic interviews and outlines new social settings. In fact, three or four generations earlier, broomcorn and foxtail millet were widely grown in the Iberian Peninsula. In many of these regions, these crops did not endure in the memory of the farmers. The key reasons for this neglect may be the replacement of millets by other crops and the loss of traditional knowledge orally transmitted from generation to generation.

We must question the extent to which the results of the study of the materiality of these contemporary traditional societies can be transferred to the past. Traditional societies are not the guardians of ancestral practices, but the result of a process of technological, economic, social or environmental adaptation and change. The transformation of traditional agricultural societies in this part of Iberia is not recent, because agriculturalists have adopted many technical and social changes since the 19th century (see Fernández Prieto 1992). Furthermore, traditional agriculture presents social, economic and cultural conducts that cannot necessarily be extrapolated to the recent past. If we focus on the search for archaeological analogy, it may be more interesting to recognise these social settings. In this way, we can consider all available parameters. When it comes to characterising the origin of a behaviour, all behaviours have the same initial value. In other words, farming practices are a portrayal of the objects and the people at the exact moment when they took place, in each specific social and cultural setting.




Figure 3. Millet cultivation in Spain according to the survey conducted by Pascual Madoz (1846-1850). a) references to 'mijo' and 'panizo'; b) filtered nomenclatures to separate out references to millets from probable references to maize.

Diversity of social networks and climatic determinants

Millets are resilient to a variety of agroclimatic challenges, require minimal energy input to yield grain, have a short growth cycle (of 50-90 days) and are well adapted to limited rainfall and poor soils. These characteristics allow their cultivation both in rainy seasons in semi-arid areas and in dry seasons in wet climates. The Iberian Peninsula has high contrasts in rainfall patterns. The southeast collects less than 300 mm per year, while several territories in the northwest and the north reach more than 2500 mm per year. The places studied in the ethnographic research are located within the area of the Atlantic oceanic climate, which has high rainfall. The information extracted from Madoz's survey also indicates that the humid regions of Spain are where cultivation of millets was the most well established until recently (Fig. 4). These are areas with high rainfall (1000-2500 mm per year). Millets are extremely rare in the Mediterranean climate region, except in the northeastern area which receives considerable summer rainfall. In central and southern Iberia, they are cultivated in rainy mountainous areas or in valley bottoms close to river courses. Accordingly, *Panicum miliaceum* and *Setaria italica* found their best agricultural niche in the dry seasons of wet areas, as summer crops. This is corroborated by the sowing seasons. In Asturias, traditional sowing of *S. italica* took place between the end of May and mid-June (Moreno-Larrazabal *et al.* 2015). In Galicia and northwestern Portugal, sowing of *P. miliaceum* took place later, from the first to the last week of June (Vázquez Varela 1994; Moreno-Larrazabal *et al.* 2015).

Although climate is a factor that determines agricultural decisions, a purely climate-based explanation is insufficient to interpret the complexity of the distribution of millet crops. This requires understanding their roles and their interaction with each element of each agricultural system. Communities have sought their own ways of exploiting the land. In Iberia, millet was more related to intensive farming systems than to extensive systems. Broadly speaking, traditional intensive agriculture occupied the northern area, that is, the area with the highest rainfall. Agricultural intensification, with higher inputs of labour, increased fertilisation, frequent rotation and crop diversification allowed high profits on small plots. The land was generally farmed by smallholders, usually from the same family group. Here, thanks to their adaptability, millets found a suitable niche for their cultivation. In the Mediterranean bioclimatic area, rainfall is concentrated in the winter months. Agriculture was focused on an extensive system geared towards the production of winter cereals, but with areas of intensive irrigated production (orchards) and forestry (fruit trees).

Family models, inheritance systems and land ownership or usufruct also exert strong influence on agricultural systems. Although investigating these aspects of archaeology is a difficult task, it is worth doing so ethnographically, because they play a crucial role in the choice of crops. The different systems of inheritance between the coastal and inland areas of Galicia during the Modern Age (16th-18th centuries CE) are a good example. In many areas of inland Galicia, inheritance to a single descendant predominates. Here, an extensive system based on winter cereals, such as rye and wheat, was maintained, together with the exploitation of silviculture species, such as vine and chestnut, which provided economic profitability. The widespread use of fallowing led to the need to preserve the patrimonial goods belonging to the household and to encourage a complex type of family organisation in the domestic group, necessary for extensive agricultural production (Sobrado Correa 2001). A double cereal harvest would mean overexploitation of the land, which was not in the group's best interest in the mid-term. Therefore, in these interior areas of Lugo, millet cultivation involved a single harvest per year.




Figure 4. Rainfall in Spanish millet-growing areas since the 19th century, according to Ninyerola et al. (2005): a) annual rainfall; b) June rainfall.

In the western area of Galicia, influenced by the coast and its surroundings, the inheritance system was more equitable. Generally, all the descendants received a portion of the land, although one of them received twice the amount of the others. Smallholder farming with reduced plot size is a characteristic element of agricultural production, which was balanced by intensive farming, rotation and diversification. In this system, the cultivation of millets would potentially be of greater interest thanks to their flexibility in crop rotation. The straw is of little interest as fodder, as the preferred method of livestock feeding was the production of pasture grass, alternated with the cultivation of oats, maize, vegetables or tubers, among others. This is where annual rye-broomcorn millet rotation was documented ethnographically (Fig. 1e). The written sources also mentioned this association and other rotations, at least since ca. 1600: winter cereal-broomcorn millet-fallow, winter cereal-turnips-broomcorn millet (Saavedra 2018). During the Modern Age, millets represented up to one third of the income received by the ecclesiastical institutions in many Galician coastal regions, until the arrival of maize, which displaced millets in importance starting with the economic crisis of 1627-1632 (Fernández Cortizo 1991).

Early millets in the Iberian Peninsula: Uneven adoption

In the previous chapters, different examples have been given of factors that may have influenced millet agriculture in the past and which point to situations different from an ideal picture of a uniform transmission of farming. In fact, it is common to find different spatial distributions and intensities of millet when analysing archaeobotanical assemblages. Some of the patterns mentioned can be traced back to protohistory (Fig. 5), but others seem to reflect a different trajectory.

Archaeobotanical research indicates the possible early introduction of millets into the Iberian Peninsula around 1650-1300 BCE (cf. Alonso Martínez 2000b; Peña-Chocarro 2000; Tereso *et al.* 2016; Tarongi Chavarri 2017; Jesus *et al.* 2020), although the timing still needs to be tested by direct dating of the grains. Despite finds of some significant archaeobotanical assemblages (Alonso Martínez 2000b), their distribution does not yet provide us information about their differential acceptance by the communities of the Middle Bronze Age and Late Bronze Age.

For the 1st millennium BCE, regional overviews corroborate the great expansion and intensity of millet cultivation (Alonso Martínez 2000a; Pérez Jordá 2013; Tarongi Chavarri 2017; Teira Brión 2019). Territories in the northwest have the highest frequency per sample; here, *Panicum miliaceum* is the second-most ubiquitous crop (24.90%), right after *Triticum* spp. (Teira Brión 2019). Millets were fundamental crops within a balanced system incorporating winter cereals and spring cereals; a poorly diversified leguminous group centred on two species, broad bean (*Vicia faba*) and pea (*Pisum sativum*); and the management of wild forest resources, such as acorns. In other territories, it was also a common cereal. For example, in the Ebro River valley, the ubiquity of *P. miliaceum* is 13.3% and that of *Setaria italica* is 6.24% in the samples from the 1st millennium BCE (Tarongi Chavarri 2017). In the region of València, the ubiquity of broomcorn millet is 11.85% and that of foxtail millet is 5.67% (cf. Pérez Jordá 2013). In contrast, in Upper Andalusia, the frequency of both cereals is low (cf. Montes Moya 2014). Although they are generally well documented in the parts of Iberia with a Mediterranean climate, millets were secondary cereals here, coming after wheat, barley and fruits, such as the vine (*Vitis vinifera*) (Alonso Martínez 2000a; Tarongi Chavarri 2017). In these areas (the Mediterranean and southern regions), occupied by Iron Age communities known as the Iberians, agriculture was based on winter cereals, forestry practices that had been introduced by the Phoenicians (Pérez-Jordá *et al.* 2021), and a greater variability of legumes that can withstand dry




Figure 5. Ubiquity (Ub.) of *Panicum miliaceum* (*P. mil*) and *Setaria italica* (*S. ita*) in Iberian archaeobotanical samples from the Iron Age (data from the databases by Pérez Jordá 2013; Montes Moya 2014; Tarongi Chavarri 2017; Teira Brión 2019), Roman period and Middle Ages (data from the database by Peña-Chocarro et al. 2019).

conditions, such as lentil or chickpea and bitter vetch (Buxó 1997; Alonso Martínez 2000a; Tarongi Chavarri 2017). In the Iberian south, one of the areas yielding the first documented *P. miliaceum* during the Bronze Age, the cultivation of millets seems to have been rare and had already declined during the Iron Age.

For the Roman period and the Middle Ages, we have information from 1353 archaeobotanical samples compiled in a recent synthesis of the entire Iberian Peninsula (Peña-Chocarro *et al.* 2019). Although there are areas that are poorly investigated, the data provide us with an excellent overview for understanding the distribution of millet cultivation. In both periods, millets played a more important role in the northern regions than in the central and southern areas of the peninsula. The high frequency of *Setaria italica* in the eastern region in Roman times and in the northern area in the Middle Ages can be interpreted as being the result of sampling bias, caused by the low number of samples taken and the consequent overrepresentation of some contexts. However, there seems to be no doubt that they were widespread cereals. Archaeological evidence indicates that northern Iberia would have been the area with the highest incidence of millets, and where they would have been more resilient to the changes introduced in agricultural systems over time. Also, it is important to note their significantly lesser importance and near-absence in many settlements in the south.

Despite the unequal evolution observed throughout these periods, we still cannot interpret the relative importance of each crop. Differences in specific practices may have been mediated by the decisions of each society regarding agricultural production, diet and the formation of each social or economic network and cultural tradition. In fact, carpological analyses hardly distinguish between uses, between fallow or crop rotation systems, and they do not tell us whether a double annual harvest is feasible, or even whether uses are similar throughout the Iberian Peninsula or are regionally differentiated. To investigate these questions, we still need the help of interdisciplinary approaches and regional case studies to allow the information obtained to be deciphered in the relevant social context. In this sense, millet research has been revitalised to address the questions of when and how it was consumed, based on the isotopic analysis of human bones. This research brings some interesting results, some of which were also observed using the ethnographic approach and some of which have revealed important dietary novelties. For example, isotopic evidence from the Iron Age village of La Hoya suggests that C4 cereals may have played a role in infant and child feeding, being consumed during the weaning period, probably as porridge (Fernández-Crespo *et al.* 2019). Isotope studies show significant disparities in millet consumption, which reflect: 1) regional and local differences (Lubritto *et al.* 2017; López-Costas and Alexander 2019); 2) preferential ingestion by middle- to lower-class (López-Costas and Müldner 2019) or high-status members (Toso *et al.* 2019); 3) distinct intakes between childhood and adulthood (Dury *et al.* 2019); 4) gender differences in adult diet, with higher C4 values among women (Alaica *et al.* 2019) or men (Toso *et al.* 2019); and 5) food preparation based on a combination of C3 and C4 crops in the diet of dogs (Albizuri *et al.* 2021).

Conclusions

In this text, I have aimed to draw attention to the problems that result from uncritically transferring some ethnography-based statements made in the present to the past. The evolution of millets in the Iberian Peninsula shows different developments based on each agricultural system and the aspects such as the land use, rotation, crops, property and type of family that condition the importance of their incorporation. All these practices involve societies in different ways and require diverse social and productive organisations – aspects that also influence the present that we study through ethnoarchaeology.

Not everything we observe necessarily has ancient roots; some things may be linked to practice transformations that are closer in time. Adaptive changes happen within a short or long period. It is a complex and laborious task to untangle at what point many practices we observe could have taken place in the past. It is the human being who decides on the agricultural system and the obtaining of resources, and who ultimately chooses, harmonises and selects some uses and some crops over others. The examples in the previous chapters indicate behaviours inherent to extensive or intensive farming systems, which cannot be understood in isolation from possible soil and climate conditions (optimal growing period) or independently from the uses given by each community.

Carpological results by themselves do not support certain views which are held in research about the introduction of millets into European agriculture: the hypothesis of double annual cereal production simply because of the coexistence of millets with other cereals, and the hypothesis about the increase in agricultural production. These two statements can only be evaluated by assessing all the elements which make up each agricultural system.

During the past four millennia, since the introduction of millets in the Iberian Peninsula, some trends have endured over the long term, despite the social and technological changes that continue until the present day. We can attribute this to the adaptability of millets, which has allowed them to be resilient to changes in socio-economic models. We should not forget the role of the transmission of traditional knowledge in peasant societies, which implies mechanisms, adaptations and the transmission of know-how that regulate the practices of each community both socially and culturally, and which are used to respond to the environment and guide the management of resources (Berkes *et al.* 2000). The knowledge of millet cultivation would have survived until the arrival of changes leading to a quick or progressive transformation of the agricultural system, until the substitution of their role as essential cereals or as secondary crops. Therefore, one of the aspects we can consider as important is to establish the chronological depth of the practices we are studying.

When studying millet farming, we must apply a multidisciplinary and multiscale approach in order to accommodate different settings. It is in this context that the contribution of ethnoarchaeology can be crucial, giving us guidelines on which social aspects we need to heed and which particular material elements to analyse. Ethnographic results should not be taken only as the standardisation of ideas that highlight those elements that are most interesting from our point of view. The picture created through ethnography is a construct that must contemplate a diversity of practices in a constant process of construction. In conclusion, if millets have a transformative power in agriculture and diet, we must understand how they are embedded in each agricultural system and society.

Acknowledgements

My gratitude goes to all the farmers who have shared their knowledge transmitted from generation to generation and to Aitor Moreno, Lydia Zapata and Itsaso Sopelana for sharing with me this journey through the history of millets. I would also like to acknowledge the generosity of Leonor Peña-Chocarro, Guillem Pérez-Jordà, Natàlia Alonso, Ferran Antolín, João Tereso, Eva Montes Moya and Dani López Reyes for allowing me to use unpublished data produced for the carpological synthesis for the Roman and Middle Ages in the Iberian Peninsula (Peña-Chocarro *et al.* 2019). The present research is included in the actions of the project 'Millet: old crops for new insights: agricultural systems of broomcorn and foxtail millet cultivation in Iberia', supported by a Marie Skłodowska-Curie Individual Fellowship H2020-MSCA-IF-2020 (REA grant agreement ID: 101018935).

References

- Alaica, A.K., Schalburg-Clayton, J., Dalton, A., Kranioti, E., Graziani Echávarri, G., Pickard, C., 2019. Variability along the frontier: stable carbon and nitrogen isotope ratio analysis of human remains from the Late Roman-Early Byzantine cemetery site of Joan Planells, Ibiza, Spain. *Archaeological and Anthropological Sciences*, 11 (8), 3783-3796. <https://doi.org/10.1007/s12520-018-0656-0>
- Albizuri, S., Grandal-d'Anglade, A., Maroto, J., Oliva, M., Rodríguez, A., Terrats, N., Palomo, A., López-Cachero, F.J., 2021. Dogs that ate plants: changes in the canine diet during the Late Bronze Age and the First Iron Age in the northeast Iberian Peninsula. *Journal of World Prehistory*, 34 (1), 75-119. <https://doi.org/10.1007/s10963-021-09153-9>
- Alonso Martínez, N., 2000a. Cultivos y producción agrícola en época ibérica. *Saguntum: Papeles del Laboratorio de Arqueología de Valencia, Extra* 3, 26-46.
- Alonso Martínez, N., 2000b. Registro arqueobotánico de Cataluña Occidental durante el II y I milenio a.n.e. *Complutum*, 11, 221-238.
- Barboff, M., 1995. Le millet au Portugal. In: E. Hörander, ed., *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang, 113-121.
- Berkes, F., Colding, J., Folke, C., 2000. Rediscovery of traditional ecological knowledge as adaptive management. *Ecological Applications*, 10 (5), 1251-1262. <https://doi.org/10.2307/2641280>
- Buxó, R., 1997. *Arqueología de las plantas*. Barcelona: Crítica.
- Dury, G., Lythe, A., Marquez-Grant, N., Garcia-Rubio, A., Graziani, G., Mari, J., Ziríax, M., Schulting, R., 2019. The Islamic cemetery at 33 Bartomeu Vicent Ramon, Ibiza: investigating diet and mobility through light stable isotopes in bone collagen and tooth enamel. *Archaeological and Anthropological Sciences*, 11 (8), 3913-3930. <https://doi.org/10.1007/s12520-018-0644-4>
- Fernández Cortizo, C., 1991. Estrategias familiares y pequeña explotación campesina en la Galicia del siglo XVIII. In: P. Saavedra and R. Villares Paz, eds., *Señores y campesinos en la Península Ibérica, siglos XVIII-XX*. Barcelona: Crítica, 310-345.
- Fernández Prieto, L., 1992. *Labregos con ciencia: estado, sociedade e innovación tecnolóxica na agricultura galega, 1850-1936*. Vigo: Xerais.
- Fernández-Crespo, T., Ordoño, J., Bogaard, A., Llanos, A., Schulting, R., 2019. A snapshot of subsistence in Iron Age Iberia: the case of La Hoya village. *Journal of Archaeological Science: Reports*, 28, 102037. <https://doi.org/10.1016/j.jasrep.2019.102037>
- González-Ruibal, A., 2014. *An Archaeology of Resistance: Materiality and Time in an African Borderland*. Walnut Creek: Left Coast Press.
- Hamilakis, Y. and Anagnostopoulos, A., 2009. What is archaeological ethnography? *Public Archaeology: Archaeological Ethnographies*, 8 (2-3), 65-87. <https://doi.org/10.1179/175355309X457150>
- Hamon, C. and Le Gall, V., 2013. Millet and sauce: the uses and functions of querns among the Minyanka (Mali). *Journal of Anthropological Archaeology*, 32 (1), 109-121. <http://dx.doi.org/10.1016/j.jaa.2012.12.002>
- Hernando Gonzalo, A., 1995. La etnoarqueología, hoy: una vía eficaz de aproximación al pasado. *Trabajos de Prehistoria*, 52 (2), 15-30.
- Hörander, E., ed., 1995. *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang.
- Jesus, A., Tereso, J.P., Gaspar, R., 2020. Interpretative trajectories towards the understanding of negative features using Terraço das Laranjeiras Bronze Age site as a case study. *Journal of Archaeological Science: Reports*, 30, 102222. <https://doi.org/10.1016/j.jasrep.2020.102222>

- López-Costas, O. and Alexander, M., 2019. Paleodiet in the Iberian Peninsula: exploring the connections between diet, culture, disease and environment using isotopic and osteoarchaeological evidence. *Archaeological and Anthropological Sciences*, 11 (8), 3653-3664. <https://doi.org/10.1007/s12520-019-00886-5>
- López-Costas, O. and Müldner, G., 2019. Boom and bust at a medieval fishing port: dietary preferences of fishers and artisan families from Pontevedra (Galicia, NW Spain) during the late medieval and Early Modern period. *Archaeological and Anthropological Sciences*, 11 (8), 3717-3731. <https://doi.org/10.1007/s12520-018-0733-4>
- Lubritto, C., García-Collado, M.I., Ricci, P., Altieri, S., Sirignano, C., Quirós Castillo, J.A., 2017. New dietary evidence on medieval rural communities of the Basque country (Spain) and its surroundings from carbon and nitrogen stable isotope analyses: social insights, diachronic changes and geographic comparison. *International Journal of Osteoarchaeology*, 27 (6), 984-1002. <https://doi.org/10.1002/oa.2610>
- Lundström-Baudais, K., Rachoud-Schneider, A.M., Baudais, D., Poissonnier, B., 2002. Le broyage dans le chaîne de transformation du millet (*Panicum miliaceum*): outils, gestes et écofacts. In: Procopiou, H. and Treuil, R., eds., *Moudre et Broyer: I Méthodes*. Paris: Comité des Travaux historiques et Scientifiques, 150-180.
- Madoz, P., 1846-1850. *Diccionario Geográfico-Estadístico-Histórico de España y sus posesiones de ultramar*. Madrid: Establecimiento tipográfico de P. Madoz y L. Sagasti.
- Montes Moya, E.M., 2014. *Las prácticas agrícolas en la Alta Andalucía a través de los análisis carpológicos (desde la Prehistoria Reciente al s. II d.n.e.)*. Doctoral dissertation, Universidad de Jaén.
- Moreno-Larrazabal, A., Teira-Brión, A., Sopelana-Salcedo, I., Arranz-Otaegui, A., Zapata, L., 2015. Ethnobotany of millet cultivation in the north of the Iberian Peninsula. *Vegetation History and Archaeobotany*, 24 (4), 541-554. <https://doi.org/10.1007/s00334-015-0518-y>
- Ninyerola, M., Pons, X., Roure, J.M., 2005. *Atlas Climático Digital de la Península Ibérica: Metodología y Aplicaciones en Bioclimatología y Geobotánica*. Bellaterra: Universidad Autónoma de Barcelona.
- Peña-Chocarro, L., 2000. Agricultura y alimentación vegetal en el poblado de la Edad del Bronce de Peñalosa (Baños de la Encina, Jaén). *Complutum*, 11, 209-219.
- Peña-Chocarro, L., Pérez-Jordà, G., Alonso, N., Antolín, F., Teira-Brión, A., Tereso, J.P., Montes Moya, E.M., López Reyes, D., 2019. Roman and Medieval crops of the Iberian Peninsula: a first synthesis of seeds and fruits from archaeological sites. *Quaternary International*, 499 (A), 49-66. <https://doi.org/10.1016/j.quaint.2017.09.037>
- Pérez Jordà, G., 2013. *La Agricultura en el País Valenciano entre el VI y el I Milenio a.C.* Doctoral thesis, Universitat de València. <https://roderic.uv.es/handle/10550/31152>
- Pérez-Jordà, G., Peña-Chocarro, L., Pardo-Gordó, S., 2021. Fruits arriving to the West: introduction of cultivated fruits in the Iberian Peninsula. *Journal of Archaeological Science: Reports*, 35, 102683. <https://doi.org/10.1016/j.jasrep.2020.102683>
- Saavedra, P., 2018. El maíz en el sistema agrario y en la alimentación en Galicia, siglos XVII-XIX. *Ohm: Obradoiro de Historia Moderna*, 27, 49-80. <https://doi.org/10.15304/ohm.27.5523>
- Sillar, B., 2000. *Making Pots and Constructing Households: an Ethnoarchaeological Study of Pottery Production, Trade and Use in the Andes*. British Archaeological Reports International Series 883. Oxford: Archaeopress.
- Sobrado Correa, H., 2001. Heredero único y familia troncal en la Galicia interior, 1750-1860. *Studia Historica: Historia Medieval*, 23, 275-303.
- Song, J., Zhao, Z., Fuller, D.Q., 2013. The archaeobotanical significance of immature millet grains: an experimental case study of Chinese millet crop processing. *Vegetation History and Archaeobotany*, 22 (2), 141-152. <https://doi.org/10.1007/s00334-012-0366-y>

- Tarongi Chavarri, M., 2017. Análisis comparativo de los estudios carpológicos de la Depresión del Ebro durante el III y I milenio a.C.: un estado de la cuestión. *Revista d'Arqueologia de Ponent*, 27, 41-59.
- Teira Brión, A., 2019. *Cambio e resiliencia na agricultura e xestión de recursos vexetais no NW da Península Ibérica (1000 a.n.e.-400 d.n.e.)*. Doctoral dissertation, Universidade de Santiago de Compostela. <http://hdl.handle.net/10347/20497>
- Tereso, J.P., Bettencourt, A.M.d.S., Ramil-Rego, P., Teira-Brión, A., López-Dóriga, I., Lima, A., Almeida, R., 2016. Agriculture in NW Iberia during the Bronze Age: a review of archaeobotanical data. *Journal of Archaeological Science: Reports*, 10, 44-58. <http://dx.doi.org/10.1016/j.jasrep.2016.07.011>
- Toso, A., Gaspar, S., Banha da Silva, R., Garcia, S.J., Alexander, M., 2019. High status diet and health in medieval Lisbon: a combined isotopic and osteological analysis of the Islamic population from São Jorge Castle, Portugal. *Archaeological and Anthropological Sciences*, 11 (8), 3699-3716. <https://doi.org/10.1007/s12520-019-00822-7>
- Vázquez Varela, J.M., 1994. O cultivo tradicional do 'mijo', millo miúdo, (*Panicum miliaceum* L.) en Galicia. In: A. Fraguas Fraguas and X.A. Fidalgo Santamariña, eds., *Tecnoloxía Tradicional: Dimensión Patrimonial: Valoración Antropolóxica*. Santiago de Compostela: Consello da Cultura Galega, 117-126.

Versatile usage of millet: Brooms and animal fodder from *Sorghum technicum*

Wiebke Kirleis and Marta Dal Corso

Abstract

On a field survey in northwestern Moldova in August 2019, when searching for archives for palynological investigations, we passed by fields of *Sorghum technicum* and had the good fortune of meeting local farmers on the fields, who introduced us into a very special local tradition. The growing of *Sorghum technicum* is directly linked with the production of sorghum brooms in the nearby village of Cărăcușenii Vechi. Tying sorghum brooms is a highly specialised craft that in modern times is rarely encountered. Therefore, we decided to share our observations and take the opportunity to document small-scale, traditional broom production in Moldova.

Introduction

Sorghum and millets are commonly used for human consumption or as animal fodder. Here we focus on the special breed *Sorghum technicum* and document its cultivation, harvesting and processing in the vicinity of the village of Caracușenii Vechi (Briceni district), in northwestern Moldova, in 2019. The crop's name is already indicative of its main purpose, that is, a technical use. *Sorghum technicum* is the basic resource for organic broom production. The by-products, in this case the leaves and the grains, serve as animal fodder. In Caracușenii Vechi, the tradition of manufacturing sorghum brooms is kept alive in small, family-owned workshops. In addition to Moldova, there are other places in eastern Europe, Thailand and China where organic sorghum broom production takes place, one further example being EcoBrooms, in the Vojvodina, in Serbia (Taylor *et al.* 2019, 414). In Bulgaria, *Sorghum bicolor* is referred to as the crop for broom making (Nedelcheva *et al.* 2007).

Sorghum technicum is a C4 plant, and its growth habitus is reminiscent of maize (*Zea mays*), which is taxonomically and genetically a close relative of all *Sorghum* species. It grows up to 3 m high, and its pendant panicle reaches lengths of 60 cm

Figure 1. *Sorghum technicum*, exhibiting the growth habitus of a typical C4 plant, such as maize (photo: Wiebke Kirleis).



Figure 2. *Sorghum technicum* panicle (photo: Marta Dal Corso).



(Figs. 1 and 2). It is a macro-thermal crop (or summer crop), which at Caracușenii Vechi was seeded in April and harvested at the end of August.

Different from maize, which originates in the Americas, sorghum originates in Africa and was domesticated ca. 3000 BCE (Fuller and Stevens 2018; Winchel *et al.* 2018). From there it spread to India and China (Liu *et al.* 2019), and in a later stage of history it reached the Americas by means of ships transporting enslaved people (Taylor 2019). Botanically, it belongs to the Panicoideae subfamily of the Andropogoneae tribe, which is in the Poaceae family. The classification within the genus *Sorghum* is very complex. According to De Wet and Harlan (1971; USDA Germplasm Resources Information Network), all cultivated sorghum species belong



Figure 3. Single plants of *Sorghum technicum* towering above the arable field (photo: Marta Dal Corso).

to *Sorghum bicolor* ssp. *bicolor*. However, a huge genetic diversity is observed within the genus *Sorghum*, as gene flow occurs freely between cultivars as well as between the genus *Sorghum* and its wild and weedy relatives (Taylor 2019). In Caracușenii Vechi, such diversity is expressed by single plants, towering above the field (Fig. 3).

In the Mansfeld World Database of Agricultural and Horticultural Crops (Schultze-Motel 1986; IPK Gatersleben 2020), *Sorghum technicum* is listed as *Sorghum saccharatum* convar. *technicum* (Koern.) Tzvelev in *Novosti sist. vyss. rast.* vol. 5, 15. 1968, with several synonyms:

1. *Holcus sorghum* var. *technicus* L.H. Bailey, *Gentes Herb.* vol. 1, 132. 1923.
2. *Andropogon sorghum* var. *technicus* Hack. ex A. DC. & C. DC., *Monogr. phan.* vol. 6, 508. 1889.
3. *Sorghum vulgare* var. *technicum* Fiori & Paol., *Fl. Italia* vol. 1, 46. 1896.
4. *Sorghum technicum* Battand. & Trab., *Fl. Algerie Monocot.*, 128. 1895.
5. *Andropogon sorghum* var. *technicus* Koern., *Handb. Getreideb.* vol. 1, 308. 1885.
6. *Sorghum dochna* var. *technicum* Snowden in *Kew Bull.*, 235. 1935.

From millet to broom: *Chaîne opératoire* of broom making in northwestern Moldova

In the vicinity of Caracușenii Vechi, sorghum is grown in monoculture on small, family-owned or leased fields some hectares in extent. In 2019, harvesting took place in late August, before the rainy season started. The long stalks were harvested either by hand, using a sickle or a scythe (Fig. 4), or with the aid of large machines. The stalks cut by the harvesting machine are left for 3-4 days to dry on the field, before being manually picked from the ground (Fig. 5) and transferred to small heaps in a shady location along the edge of the field.

In 2019, the harvesting was organised with several family members gathering in the village and collectively working on the field. Other people from the community joined in for further on-field processing of the plants. Tasks did not appear to be gendered: adult men and women of diverse age classes worked together. The next



Figure 4. Harvesting sorghum with a scythe (photo: Wiebke Kirleis).



Figure 5. Post-mechanical cutting: a family collecting the plants on the arable field (photo: Wiebke Kirleis).



Figure 6a. Collective labour: defoliating the plants on the arable field (photo: Wiebke Kirleis).



Figure 6b. Defoliating the plants on the arable field (photo: Wiebke Kirleis).

step, after collection of the dry stalks and piling them up into small heaps, is to defoliate the stalks. Individuals were responsible for defoliating the stalks of each sorghum heap (Fig. 6a-b).

After the stalks have been defoliated, they are bundled and stacked (Figs. 7-8). The leaves, being a by-product, are piled into heaps beneath the stacked sorghum bundles. The bundled sorghum is then taken to the village (Fig. 8).

The next stages of processing of the sorghum bundles take place in the village. Bundle after bundle of sorghum is treated with a mechanical threshing machine. The panicles are fed into the machine and the grains are released and caught in small containers. The grains serve as animal fodder, and the threshed bundles are



Figure 7. Defoliated bundles, and leaves representing a by-product (photo: Wiebke Kirleis).



Figure 8. Defoliated bundles, detail (photo: Marta Dal Corso).



Figure 9. Transport of the bundles to the village (photo: Wiebke Kirleis).



Figure 10a. Bundles after mechanical threshing, ready for broom production (photo: Wiebke Kirleis).



Figure 10b. Detail: threshed panicles with some remaining fruits (photo: Wiebke Kirleis).

positioned alongside fences along the streets to for further drying before they are distributed to the workshops for tying into brooms (Figs. 10a-b and 11).

The tying of brooms takes place as mass production in small, specialist workshops or is individually organised as a hobby during long winter evenings spent tying nicely decorated prestige brooms (Fig. 12a-b).



Figure 11. Drying of threshed bundles of *Sorghum technicum* in the village of Caracuşenii Vechi, Moldova (photo: Marta Dal Corso).



Figure 12a. Broommakers' workshop (photo: Wiebke Kirleis).




Figure 12b. Special chair with construction for clamping the broom while cording it (photo: Marta Dal Corso).

Tying of the brooms in Caracușenii Vechi starts with combining some bundles and clamping them in a specific chair construction to achieve compact brooms (Fig. 12a-b). In this construction, the bundles are corded to fix the broom's handle. Afterwards, the handle's rough end is cut off. Then the loose ends of the threshed panicles still have to be fashioned into the shape of a broom. Afterwards, they are cut off straight. This is the final stage of the mass production of simple sorghum brooms (Figs. 13-14).



Figure 13. Storage of simple sorghum brooms in the workshop (photo: Marta Dal Corso).



Figure 14. Sale of custom sorghum brooms (photo: Marta Dal Corso).



Figure 15. Prestige sorghum broom with embroidered text, which translates as 'My village with people, hardworking, beautiful, good, Caracusenii Vechi' (photo: Wiebke Kirleis).

To obtain more sophisticated brooms, the handles are sometimes plaited. Further, stitching is done to keep the broom flat. Some brooms even represent prestige items, with embroidery, being one expression of cultural identity. Nowadays these are used to decorate the walls of homes, serve as talisman to bring luck, are given as a souvenir to family members moving away to the cities or abroad, or are sold at traditional handicraft markets (Fig. 15).

Concluding remarks

The aim of this contribution on the uses of *Sorghum technicum* is twofold. First, it documents the cultivation of a specific crop directly linked to traditional handicrafts, in particular the *chaîne opératoire* from sorghum harvesting to the tying of brooms. The technical processing of sorghum reflects different sociocultural and economic aspects of life. It gives insight into plant use and related techniques, related social organisation, and related cultural value. Second, viewed from an archaeological perspective, organic brooms belong to the perishable material culture. They rarely survive in archaeological deposits (Nedelcheva *et al.* 2007; Hurcombe 2014). The documentation presented here is a reminder for the archaeological community when reconstructing former routine activities to consider multiple human-plant relationships that extend beyond dietary aspects. Lateral thinking is required to reconstruct the missing parts.

Acknowledgements

We are very thankful to Grigory, Lucia and Tanja, who were harvesting the fields between Kostantinovka and Caracuşenii Vechi, for insights into *Sorghum* cultivation and processing. Thank-you also to Leova, who showed us his workshop, explained the handicraft of tying the brooms and produced the tiny brooms that served as gifts to the participants of the workshop in Kiel. Our warmest thanks go also to Svetlana Ibens, Kiel, for helping us with translation during our fieldwork. This documentation was produced during archaeological fieldwork carried out within subproject D1 of the SFB 1266, funded by the German Research Foundation – DFG (Project number 2901391021-SFB 1266).

References

- Fuller, D.Q. and Stevens, C.J., 2018. Sorghum domestication and diversification: a current archaeobotanical perspective. In: A. Mercuri, A. D'Andrea, R. Fornaciari, A. Höhn, eds., *Plants and People in the African Past*. Cham: Springer, 427-452. https://doi.org/10.1007/978-3-319-89839-1_19
- IPK Gatersleben, 2020. *Mansfeld World Database of Agricultural and Horticultural Crops*. Webproject at IPK Leibniz Institute Gatersleben [online]. <https://mansfeld.ipk-gatersleben.de/>
- Hurcombe, L.M., 2014. *Perishable Material Culture in Prehistory: Investigating the Missing Majority*. Abingdon: Routledge.
- Liu, X., Jones, P.J., Motuzaite Matuzeviciute, G., Hunt, H.V., Lister, D.L., An, T., Przelomskaef, N., Knealeb, C.J., Zhaog, Z., Jones, M.K., 2019. From ecological opportunism to multi-cropping: mapping food globalisation in prehistory. *Quaternary Science Reviews* 206, 21-28. <https://doi.org/10.1016/j.quascirev.2018.12.017>
- Nedelcheva, A.M., Dogan, Y., Guarrera, P.M., 2007. Plants traditionally used to make brooms in several European countries. *Journal of Ethnobiology and Ethnomedicine*, 3 (1), 1-11. <https://doi.org/10.1186/1746-4269-3-20>

- Schultze-Motel, J., 1986. *Rudolf Mansfeld, Verzeichnis landwirtschaftlicher und gärtnerischer Kulturpflanzen ohne Zierpflanzen*. Berlin: Springer.
- Taylor, J.R.N., 2019. Sorghum and millets: taxonomy, history, distribution, and production. In: J.R.N. Taylor and K.G. Duodu, eds., *Sorghum and Millets: Chemistry, Technology, and Nutritional Attributes*. Cambridge: Elsevier, 1-21.
- Taylor, J.R.N., Zhang, K., Wang, D., 2019. Industrial and nonfood applications. In: J.R.N. Taylor and K.G. Duodu, eds., *Sorghum and Millets: Chemistry, Technology, and Nutritional Attributes*. Cambridge: Elsevier, 393-420.
- U.S. Department of Agriculture. *USDA Germplasm Resources Information Network* [online]. <https://www.ars-grin.gov/>
- De Wet, J.M.J. and Harlan, J.R., 1971. The origin and domestication of *Sorghum bicolor*. *Economic Botany*, 25, 128-135. <https://doi.org/10.1007/BF02860074>
- Winchell, F., Brass, M., Manzo, A., Beldados, A., Perna, V., Murphy, C., Stevens, C., Fuller, D.Q., 2018. On the origins and dissemination of domesticated sorghum and pearl millet across Africa and into India: a view from the Butana Group of the far eastern Sahel. *African Archaeological Review*, 35, 483-505. <https://doi.org/10.1007/s10437-018-9314-2>

CONCLUSION: Early cultivation of millet in Europe: What else and where next? Concluding the workshop proceedings

Dragana Filipović, Marta Dal Corso, Wiebke Kirleis

Millet and what else?

In much of Europe, broomcorn/common/proso millet (*Panicum miliaceum*; Fig. 1) is nowadays still primarily considered as animal feed (e.g. bird feed) and is rarely grown. With the modern-day climate change requiring new agrarian solutions, this and other millets are regaining importance because of their adaptability to sub-optimal to poor growing conditions. There have, for instance, been calls to start cultivation of pearl millet (*Pennisetum glaucum*) or sorghum (*Sorghum*) in the Po valley¹, in northern Serbia, in the Czech Republic (Hermuth *et al.* 2016) and elsewhere in Europe, and also in Asia and Africa², to counter crop failure caused by drought.

Both previous and most recent multidisciplinary research has shown that millet has been cultivated in Europe since the middle of the 2nd millennium BCE, that is, the Middle or the Late/Recent Bronze Age (depending on the part of Europe). It has since then been consumed by both humans and animals. What do we know about the history of this crop on the continent? The workshop that took place in Kiel in November 2019 aimed to start the conversation on the circumstances and consequences of early broomcorn millet cultivation in Europe. The focus was mainly on the Bronze Age subsistence economy in different regions, and on changes in this economy potentially linked with the integration of this new crop into the pre-existing spectrum of cultivars. But participants also shared knowledge on aspects of the recent and modern-day growing of millet in Europe and on other continents; on traditional millet-based cuisine; on conventional and state-of-the-art

1 <https://www.plantagbiosciences.org/people/matteo-petitti/2017/08/27/drought-proof-crops-in-italy-is-millet-the-grain-of-the-future/>

2 One of the workshop participants, Dr Stefania Grando of the International Crops Research Institute for the Semi-arid Tropics (ICRISAT) presented initiatives and projects aimed at promoting the cultivation of resilient crops and educating the farmers. See more at <https://www.icrisat.org/>



Figure 1. Broomcorn millet experimentally grown at the Archäologisch-Ökologisches Zentrum Albersdorf, Schleswig-Holstein, Germany (photo: Wiebke Kirleis).

(i.e. biogeochemical) methods of detecting millet in archaeological layers; and on the biology and physiology of millet. Even animals were brought into the picture, since millet was, and still is, used in animal husbandry.

This book brings together many of the workshop papers, reflecting the array of topics and research areas that the workshop covered. The contributions inform us on the range of cultivated and collected plants from the time before and after the start of millet cultivation in Europe; present the cultural setting in which millet arrived; discuss possible reasons driving the acceptance of this innovation; and reconstruct possible uses of millet and the methods of its cultivation, processing and storage. Not just the plant, but also the animal economy is represented, offering an integrated view of the Bronze Age food economy. Techniques used to trace millet archaeologically continue to be developed or improved, and this book describes the application of a few of them. This broad-based collection of papers demonstrates the complexity of the theme of the workshop, and of the research on the diffusion of agricultural innovations in general. As such, and together with the older and recent, ever-growing literature discussing millets in Eurasia, this book adds another layer to the dynamic picture of the interconnected continents.

The spread of millet is a spectacular display of the power of human interaction. The adaptability and resilience of the plant may have been the main reasons for its successful integration into pre-existing farming systems. This innovation was a cog in the wheel of the economic, political and social transformation in Bronze Age Europe.

Millet in Europe: Where from?

The history of the spread of millet is as complex as the physical features of the land it was carried over and through after its domestication in northeastern China (Fig. 2). Possible roads that millet took across the continents have been proposed based on the isotopic, radiocarbon or archaeobotanical evidence, or a combination thereof (e.g. Spengler *et al.* 2014; Stevens *et al.* 2016; Wang *et al.* 2017), but the location of its last stop in Asia before entering Europe is still unclear.

There is now substantial evidence that the earliest cultivation and domestication of broomcorn millet happened in northeast China, in the lower reaches of the Yellow River, at the latest by around 6000 BCE (Motuzaitė Matuzevičiūtė and Liu 2021). This coincided with and may have led to the formation of Neolithic cultures in northern China. According to the radiocarbon dates on millet grains from the sites in this and




Figure 2. Map of Eurasia indicating locations and regions (marked in brown) with early records of broomcorn millet mentioned in the text (base map © OpenStreetMap contributors; map: Dragana Filipović): 1. Begash, 2. Tasbas, 3. Dali, 4. Pethpuran Teng, 5. Adji Kui, 6. Ojakly, 7. Namcheduri, 8. Kabardinka-2, 9. Chishkho (Чижуха), 10. Guamsky Grot (Гуамский грот), 11. Safjanovo, 12. Sos Höyük, 13. Haftavan, 14. Malyan, 15. Kaneš in Kültepe, 16. Kinet Höyük, 17. Kilise Tepe, 18. Hattuša, 19. Troy, 20. Vinogradnyi Sad (Виноградный сад), 21. Assiros Toumba, 22. Mägura-Buduiasca, 23. Kastonas.

adjacent regions, recently compiled by Leipe *et al.* (2019), it appears that, for a couple of millennia, millet cultivation was restricted to the region of origin. The practice was transferred to north-central China (the upper courses of the Yellow River) by the end of the 4th millennium BCE. The network through which broomcorn millet was transmitted westward extended from the Yellow River via the Gansu Corridor between the Tibetan and the Loess (Huangtu) Plateau, on to the 'Inner Asian Mountain Corridor' (IAMC), stretching between the southwestern slopes of the Altai Mountains and the middle course of the Amu Darya River (present-day Uzbekistan), and bordered by the central Eurasian steppes in the northwest and the Tarim River basin in the southeast (Frachetti 2012; Spengler *et al.* 2014; Jones *et al.* 2016; Stevens *et al.* 2016; Wang *et al.* 2017; Hermes *et al.* 2019; Liu *et al.* 2019).

It appears that it took several centuries for millet to reach the western parts of central Asia, at least based on the dates obtained on millet grains from sites in Kazakhstan, the Kashmir valley and Turkmenistan. About halfway along the IAMC, the AMS-date on charred grains from a burial cist at the pastoralist campsite of Begash in southeastern Kazakhstan (Dzhungar Mountains) demonstrates the presence of broomcorn millet here in the mid- to late 3rd millennium BCE. Millet grains from the neighbouring site of Tasbas are probably of a similar age (Frachetti *et al.* 2010; Spengler *et al.* 2014). The stable isotope evidence from the livestock herded at the nearby settlement of Dali in the early-mid 3rd millennium BCE was interpreted as reflecting the consumption of millet (Hermes *et al.* 2019), implying that millet was already being used (as fodder) in this location at this time. It is suggested that the mobile pastoralists roaming these regions acted as a medium for the spread of the crop (Frachetti *et al.* 2010; Spengler *et al.* 2014). Southeast of this region, similarly early millet dates (from the mid-3rd millennium BCE) were reported from the site of Pethpuran Teng in the Kashmir valley (Yatoo *et al.* 2020).

According to the stable isotope values in humans and livestock from pastoralist sites located in north-central Kazakhstan, to the north and northwest of the IAMC, there was no to very little millet consumption (by animals) in the late 2nd millennium BCE and slightly more (by humans) at the turn of the millennium. Farther to the northwest, in the Trans-Urals, millet is not recorded in the diet of pastoralist groups from the 2nd and first half of the 1st millennium BCE (Motuzaite-Matuzeviciute *et al.* 2015; Wang *et al.* 2017; Ventresca Miller and Makarewicz 2019; Ananyevskaya *et al.* 2020). It is absent in the archaeobotanical record before the very end of the 2nd millennium BCE, and perhaps it is absent until even later (Ryabogina and Ivanov 2011; Rühl *et al.* 2015).

In southern central Asia (present-day Turkmenistan), archaeobotanical studies and direct radiocarbon dates document the presence of broomcorn millet starting from the end of the 3rd millennium BCE at the earliest and with certainty by the second quarter of the 2nd millennium BCE (*e.g.* at the sites of Adjı Kui 1 and Ojakly in the Murghab alluvial fan; Spengler *et al.* 2016, 2018). For much of the 2nd millennium BCE, large parts of southwestern central Asia were home to agricultural communities living in urban centres and to mobile pastoralists occupying small campsites, constituting and interacting within what is known as the Oxus civilisation (Rouse 2020). It is understood that the mobile pastoralists, who engaged in farming as part of their lifestyle, introduced millet to the sedentary populations. The latter incorporated it into their cultivation system, utilising the crop as both human food and animal fodder (Spengler *et al.* 2014; cf. Anthony 2007; Frachetti 2008).

From central Asia, the millet route may have forked to run to the north and south of the Caspian Sea (Jones *et al.* 2016; Wang *et al.* 2017) but perhaps at very different times, since the stable isotopic evidence from northern and central Kazakhstan does not show C4 signals in humans or animals before the late 2nd millennium BCE (Motuzaite-Matuzeviciute *et al.* 2015; Ananyevskaya *et al.* 2020). To the south,

towards the Iranian Plateau, broomcorn millet transmission could have followed long-known north-south trade routes. These are marked by fortified settlements along the northern foothills of the Kopet Dag mountain chain, and they connected the steppe communities with the proto-urban centres in Mesopotamia (e.g. Weeks 2003; Lecomte 2012). Absolute dates are not yet available for archaeobotanical finds of millet from sites along the southwestern coast of the Caspian Sea. Nonetheless, according to the associated archaeological finds, they derived earliest from about 1550 BCE (the site of Haftavan – Nesbitt and Summers 1988; see also Miller 2015; Stevens *et al.* 2016). A few grains recorded as '*Panicum* sp.' from the multi-period site of Malyan in the southern Zagros mountains (roughly end of the 3rd/first half of the 2nd millennium BCE based on the pottery [Miller 1982 cited in Nesbitt and Summers 1988, Table 2]) are likely of uncertain identification, as they were not reported in recent publications (e.g. Miller 2015; Miller *et al.* 2016).

West of the Caspian Sea, in the southern Caucasus region, the earliest recorded archaeobotanical finds of broomcorn millet come from the Late Bronze Age (1500-1100 BCE; Hovsepyan 2015). Recently, a project has been completed that combined archaeobotanical and stable isotopic analyses and radiocarbon dating of grains of broomcorn and foxtail (*Setaria italica*) millet from prehistoric (Neolithic through Iron Age) sites in the Caucasus (Herrscher *et al.* 2018a-b; Martin *et al.* 2021). The results point to the middle of the 2nd millennium BCE (Middle-Late Bronze Age) as the earliest possible time when broomcorn millet could have come into cultivation/use; the earliest obtained date is that on *P. miliaceum* from the site of Namcheduri in Georgia (Poz-66777, 3150±35 BP, 2σ 1501-1305 cal BCE; Martin *et al.* 2021, SI Appendix Table S2). Stable carbon isotope analysis of humans and animals from the Caucasus confirms this finding, particularly in the case of humans, in which a significant increase in $\delta^{13}\text{C}$ values is seen from the Late Bronze Age onwards (Martin *et al.* 2021).

A cross-Caucasian diffusion of millet into Europe is plausible given the continuous communication and cultural connection between the areas south and north of the Caucasus mountains, which intensified from the mid-2nd millennium BCE onwards (Sagona 2018, 378-379, 422). However, just before this time, only mountain plateaus seem to have been occupied in the northern Caucasus, whereas the foothills and the steppe areas immediately to the north were apparently uninhabited, perhaps due to the adverse living conditions that developed here following the 4.2k BP event (Knipper *et al.* 2020, 5). The site of Kabardinka-2 in the Kislovodsk region of the Northern Caucasus yielded millet grains dated to the last quarter of the 2nd millennium BCE (Poz-82838, 2950±30 BP, 2σ 1260-1051 cal BCE; Martin *et al.* 2021).

Farther north, in the lower Don River valley, numerous settlement sites and graves of the Late Bronze Age Srubnaya culture (1900-1500/1400 BCE) are recorded, indicating dense occupation and use of the river valleys and the surrounding dry steppe (Chernykh 1992; Anthony *et al.* 2005; Van Hoof *et al.* 2012). Limited archaeobotanical recovery at Srubnaya culture sites shows the presence of cereals including broomcorn millet, but the remains have not been dated (Лебедева 2005). West of this culture, between the Azov Sea and the lower Dnieper River, lived the partly contemporaneous Late Bronze Age Sabatinovka culture (1600-1300/1200 BCE). Sabatinovka culture sites also yielded millet, and the grains from one of them (Vinogradniy Sad, Ukraine) gave mid-2nd millennium BCE dates, which are currently the earliest in Europe (Filipović *et al.* 2020; Dal Corso *et al.* 2022).

By around 1400 BCE, in the broader region of the Don Delta, the Srubnaya cultural tradition was replaced by the Kobyakovo culture (1400/1300-1000 BCE), which is thought to have developed in the northwestern Caucasus, in the Kuban River valley. The Kobyakovo culture may have facilitated the diffusion of millet to the north. Occasional millet grains have been discovered at a few sites in the Kuban River area (Лебедева 2011, Table 2; Trifonov *et al.* 2017); one from the site of Chishkho was dated

earliest to the 14th century BCE (Poz-82836, 3000 ± 30 BP, 2σ 1377–1126 cal BCE; Martin *et al.* 2021). A little to the south, at the Guamsky Grot rock shelter, a concentration of millet grains was radiocarbon dated to 12th–10th century BCE (Trifonov *et al.* 2017).

North of the northwestern Caucasus, millet grains were discovered at the Kobayakovo culture settlement in Safyanovo on the Don River (Van Hoof *et al.* 2012, 73). According to the results of the stable isotope analysis so far, it was at the very end of the 2nd millennium BCE (start of the Early Iron Age) that millet consumption began on the northern slopes of the Caucasus (Hollund *et al.* 2010; Fuchs 2020; Knipper *et al.* 2020). This was almost half a millennium after millet arrived both in the southern Caucasus and in southwestern Ukraine. It thus remains unclear whether the northern/northwestern Caucasus was an early millet transition zone or whether it received the new crop later than the neighbouring regions. In addition, given the near-synchronous finds of millet in the southern Caucasus and southern Ukraine, it cannot yet be concluded whether the crop translocation proceeded south-north or north-south or whether instead or in addition to the cross-Caucasus route(s), this traffic was expedited via the Black Sea and the Crimean peninsula. That the so far earliest dated find of millet in the southern Caucasus comes from a site on the Black Sea coast (Namcheduri in Georgia; Martin *et al.* 2021) may point to the seaborne crop transfer.

The apparent absence of early (*e.g.* early to mid-2nd millennium BCE) millet south and southwest of the southern Caucasus and the Armenian highlands is, perhaps, illuminating and may suggest that millet cultivation and consumption were initially confined to only some regions. There are several sites in Anatolia with early broomcorn millet finds and their archaeological context indicates that they are unlikely to be older than the mid-2nd millennium BCE (Nesbitt and Summers 1988; Riehl 1999; Miller *et al.* 2016; Stevens *et al.* 2016). The recently obtained radiocarbon dates on millet grains from Sos Höyük, located in the highlands of northeastern Turkey, comply with this impression, as they fall in the last quarter of the 2nd millennium BCE (Poz-66780, 2945 ± 35 BP, 2σ 1261–1039 cal BCE and Poz-66781, 2960 ± 35 BP, 2σ 1276–1051 cal BCE; Martin *et al.* 2021). Stable isotopic measurements on human and animal samples from several 3rd millennium BCE sites (regional Early Bronze Age) in Anatolia do not point to regular consumption of C4 plants; the slightly more positive $\delta^{13}\text{C}$ values in some of the analysed faunal remains probably result from wild C4 forage (Irvine *et al.* 2019), which is abundant in this region and known to have contributed to animal diet from the Early Neolithic (*e.g.* Richards and Pearson 2005; Filipović 2014). It appears that broomcorn millet did not enter the Assyrian trade network operating in the first half of the 2nd millennium BCE (regional Middle Bronze Age) in Anatolia, judging by its absence from the major administrative and trade centre, the ancient city of Kaneš in Kültepe, central Turkey (Fairbairn *et al.* 2013; Fairbairn 2014). There were several millet grains in the Middle Bronze Age layer at Kinet Höyük in northern Levant (<http://www.ademnes.de>), but, without a radiocarbon date and with the later (Iron Age, early 1st millennium BCE) occupation of the site also yielding millet, the age of these finds is uncertain. At the neighbouring Bronze and Iron Age site of Kilise Tepe, millet grains (identified as *Panicum/Setaria* spp.) occur from the 12th century BCE onwards (Bending and Colledge 2007, 588).

For the Hittite kingdom (1650–1190 BCE, regional Late Bronze Age; Bryce 2005), the historical texts describe wheat and barley as staple foods, but do not mention millets (Hoffner 2003; Bryce 2005). This is in line with the absence of broomcorn millet from the immense grain storage facility in the capital of Hattuşa, which burnt in the early 16th century BCE (Neef 2001; Diffey *et al.* 2017), as well as from the archaeobotanical record from much of the second half of the 2nd millennium BCE in Anatolia (Nesbitt and Summers 1988; Fairbairn and Bradley 2008; Dörfler *et al.* 2011; Fairbairn *et al.* 2018). At the northwestern tip of Anatolia, at Troy, small

quantities of broomcorn millet grain were discovered in several occupation layers, but no earlier than phase Troy VI (1700-1250 BCE; Riehl 1999, 9, 46) and mainly in phase VII (starting at *ca.* 1250 BCE; Riehl 1999, 9; <http://www.ademnes.de>). Nesbitt and Summers (1988, 94-95) consider the possibility that broomcorn millet entered Anatolia from the north, via Transcaucasia, as more likely than that it entered from the west (or east), although they acknowledge the scarcity of the archaeobotanical record and the chronological gaps. The gaps still exist, and one should allow for the possibility that the new crop, once in the region, could have circulated within the Aegean Basin both in north-south (Valamoti 2016) and in west-east direction.

The nearest directly dated millet to Troy/Anatolia comes from the Late Bronze Age site of Assiros Toumba, in Greece, 300 km to the northwest, where large grain stores were discovered in dedicated store-rooms (Jones *et al.* 1986); the millet date falls in the 14th century BCE (Wardle *et al.* 2014). Akin to the situation Anatolia, the stable isotope values for animals and humans in mainland and northern Greece do not suggest consumption of millet during the 3rd and most of the first half of the 2nd millennium BCE (*e.g.* Petroutsas and Manolis 2010; Triantafyllou 2015; Nitsch *et al.* 2017). However, Neolithic and Early Bronze Age cattle consumption of C4 plants was noted and, as in Anatolia, explained by grazing on the local C4 vegetation, such as C4-rich coastal salt marshes (Papathanasiou 2015, 40; Nitsch *et al.* 2017). Animal consumption of wild C4 plants probably explains the $\delta^{13}\text{C}$ value of -18‰ in an Early Bronze Age human in the Peloponnese (Richards and Vika 2008). The data for the period 1600-1100 BCE in Greece (the Late Bronze Age in the Aegean Basin) indicate sporadic intake of C4 plant protein (by both animals and humans), but not necessarily or not invariably coming from millet (Petroutsas and Manolis 2010; Triantafyllou 2015).

Similarly, there are isolated cases of a C4 signal in animals (and occasionally in humans) in the coastal areas of Romania and Bulgaria since the Neolithic, and these early occurrences have also been attributed to grazing on the wild C4 vegetation (*e.g.* Honch *et al.* 2006; Balasse *et al.* 2014, 2017; but see Privat *et al.* 2018). For now, the earliest radiocarbon-dated finds of millet from Romania are the grains from Măgura-Buduiasca from the 15th-13th century BCE (Motuzaite Matuzeviciute *et al.* 2013). In Romania and Moldova, the sites of the Late Bronze Age Coslogeni and Noua cultures (second half of the 2nd millennium BCE), both closely linked with the Sabatinovka culture in the North Pontic, yielded archaeobotanical evidence of millet cultivation (Sava 2005; Sava and Kaiser 2011; Dal Corso *et al.* 2022).

On human interaction in Bronze Age Europe

Human interaction, mobility and openness to, or search for, innovations took millet from the places of its domestication and early cultivation to new, distant locales. The recently created spatiotemporal frame for the arrival and spread of broomcorn millet in Europe dates this innovation to the mid-2nd millennium BCE and demonstrates its swift adoption across the continent (Filipović *et al.* 2020). This was in the middle of the Bronze Age, a highly dynamic period in Europe teeming with new developments in nearly every sphere of life and with extensive changes in material production – the time of a wholesale societal transformation. A great deal of this dynamic stemmed from intra- and extra-regional interaction that facilitated the distribution of goods and influences in Europe and beyond. Accordingly, Bronze Age interaction and interconnectedness have been traced archaeologically based on the distribution of raw materials (most prominently metals), artefacts (*e.g.* weapons, utilitarian objects and pieces of jewellery), ornamental styles and mortuary rituals (*e.g.* Gimbutas 1965; Sherratt 1993; Harding 2000, 2013; Kristiansen 2016). Trade in goods, movement of people and spread and exchange of influence

acted as the mechanisms enabling the distribution, which was underpinned by new technological inventions in the domain of maritime and land-based transport, for instance, in shipbuilding and wagon construction (Anthony 1995; Harding 2000; Pare 2004; Kristiansen 2016).

Cultural contact and commercial relations were maintained between both neighbouring and more distant regions (Gimbutas 1965; Bouzek 1985; Jockenhövel 1991; Alberti and Sabatini 2013). Some regions had a more prominent role, such as the Mediterranean, which represented an '*unusually favourable theatre for interaction and transmission*' (Broodbank 2016, 19). Inland, the significance of the Carpathian Basin has been emphasised as a junction of cross-continental communication corridors, an epitome of cultural diversity and a cradle of increasing social hierarchy (Fisch *et al.* 2013). Farther to the north, in the regions assigned to the Nordic Bronze Age, one finds the start- and end-points of long-distance routes of trade of amber and metals respectively (Bergerbrant 2013; Vandkilde 2014; Kristiansen and Suchowska-Ducke 2015). Because rivers and river valleys connected regions, they also served as home to centres emerging along these routes and at crossroads, which initiated or boosted communication and exchange of goods. Excellent examples are the Middle-Late Bronze Age Terramare sites, in the Po Plain in northern Italy (*e.g.* Cardarelli 2009; Jones *et al.* 2014; Dalla Longa *et al.* 2019; Cardarelli *et al.* 2020), and the Late Bronze Age sites in the North Pontic region (*e.g.* Gorbenko and Grebennikov 2009); these were key west and east nodes in the network, and the Carpathian Basin was a central redistribution point.

In the archaeological scholarship on Bronze Age interaction in Europe, the main focus has been on the socio-political and economic connectivity in the context of the development of complex societies and regional hierarchies, and the concentration of power in the emergent political centres (*e.g.* Sherratt and Sherratt 1998; Harding 2000; Kristiansen and Larsson 2005; Knapp and Van Dommelen 2014; Suchowska-Ducke *et al.* 2015). In the political sphere, economic activity is a highly influential factor. Of the various economic domains in the Bronze Age, such as trade/exchange, craft production and flow of information/ideas, the subsistence economy appears to have received comparatively less research attention. There are, however, many studies documenting and discussing the incorporation of new plant or animal species into farming systems in Europe in the Bronze Age and the (resulting) changes in subsistence strategies (*e.g.* Bökönyi 1974; Sherratt 1981; Halstead 1987, 1996; Harding 1989, 2000; Behre 1998; Benecke 1998; de Hingh 2000; Greenfield 2005; Halstead and Isaakidou 2011, 2017; Bartosiewicz 2013; Stika and Heiss 2013; Kneisel *et al.* 2015; Effenberger 2018; Dal Corso *et al.* 2019; papers in this volume). Here too, inter-cultural contacts are seen as the means by which the innovations spread. Whereas identifying the 'source' and the 'recipient' of new materials, technologies or foodstuffs is of great importance for tracing the spread of materials and influences, scholars agree that a uni-directional flow between the origin and the destination is unlikely. Rather, the interacting sides actively shaped the nature, scale and content of the interactions (*e.g.* Sherratt 1993; Sherratt and Sherratt 1998; Harding 2000; Kristiansen and Larsson 2005; Galaty *et al.* 2014; Kristiansen 2016).

There have also been studies exploring spatiotemporal correlations between new developments in food economy and those in material culture, demography, ritual and environment (*e.g.* Valamoti 2016; Kneisel *et al.* 2019; Filipović *et al.* 2020; papers in this volume). This is challenging, not least because of the multiple, asynchronous chronologies, different classification and description systems, and varied terminology used for the same cultural period in different regions and vice versa. All the more challenging is understanding the nature of the relationships (if any) between changes and trends seen in different aspects of the Bronze Age archaeological record, including archaeobotanical. As the old adage goes, 'correlation does not imply causation' and so how does one determine economic, ecological or social reasons for, and ramifications of, the uptake of a new resource, such as a foreign cereal crop?

Millet as the theme of the workshop and the proceedings

The workshop in Kiel was an excellent occasion for a scientific exchange on ‘all things millet’ during direct interaction among a relatively small group of specialists, who are diverse in their expertise or in their regional research focus. Knowing what the interactions were like among the many different cultural entities living in Europe in the later part of the Bronze Age (cf. Fokkens and Harding 2013), and what their interactions with their natural environment were like (*e.g.* Capuzzo *et al.* 2018; Feeser *et al.* 2019), provides a necessary background to the knowledge of *when* millet arrived. The proceedings are an attempt to address social relations, food and materials economy and associated technologies, and the natural environment as reasons behind, or consequences of, the beginnings of millet cultivation in Europe. One of the major conclusions was that studies focusing on local or regional practices are needed to come closer to a general understanding of the *why* and *how* of the early millet cultivation.

This book offers several such studies, some looking at local, *e.g.* settlement-level evidence of millet production and use, and others taking a broader, regional or supra-regional perspective and looking at the entire plant or animal component of food economies. They emphasise the necessity for and demonstrate the potential to push further and refine the broad, continental-level picture of millet cultivation. It is with this smaller-scale, but more focused type of research that we can identify people’s needs, choices and perceptions with respect to food production strategies, and to begin to understand how individual, environmental and social dynamics may have influenced decisions such as acceptance or rejection of an innovation.

Another set of contributions in this volume outlines additional methodologies and perspectives used in the millet-centred research that, as illustrated by the findings described in these papers, can provide information not only on the cultivation and consumption of different millet species as food, but also on their uses as fodder and raw material. Just like was the case with crops known from the Neolithic, the multi-functionality of broomcorn and other millets could have been at the core of their widespread incorporation into plant economies.

Knowledge on the early history of millet in Europe: Where are we now and where do we go next?

Bioarchaeology in general has been almost saturated with millet studies or related studies, and almost every month there is a new publication dealing with this crop in one way or another.

Major archaeological interest in millet started with a focus on Asia (and China in particular) within the ERC-funded Food Globalisation in Prehistory (FOGLIP project), directed by Martin Jones and conducted at Cambridge University, UK (from a much longer list, Jones *et al.* 2011, 2016; see <http://www.foglip.mcdonald.cam.ac.uk/>). This project offered valuable guidance for the millet-centred research that has been and can be carried out in Europe. There had previously been some interest in the appearance of millet in Europe as well as in its archaeobotanical and archaeological context (*e.g.* Harding 1989; Marinval 1992; Hörander 1995; Hunt *et al.* 2008). With the refining of traditional methods, such as radiocarbon dating, and the development and application of new ones, such as stable isotopic, biochemical and genetic approaches, the scope has been enlarged and the detail improved. Thanks to this, even though it still lags behind that in China for example, the knowledge

of early millet cultivation and consumption in Europe has grown substantially in recent years. The Millet Dating Programme (Filipović *et al.* 2020), carried out in the Collaborative Research Centre 1266, at Kiel, brought together the expertise of 40 archaeobotanists and archaeologists from multiple countries; it was a key milestone, and it laid the groundwork for future work on the history of millet in Europe.

Through AMS radiocarbon dating of a large number of millet grains, this programme established that the crop first appeared in Europe around the mid-2nd millennium BCE – no earlier than the late 17th century BCE in the southeastern Europe (Ukraine) and no earlier than the late 13th century BCE in north-central Europe (northern Germany). There are now more than 100 archaeological sites with radiometrically dated millet grains (Motuzaite-Matuzeviciute *et al.* 2013; Dreslerová *et al.* 2020; Filipović *et al.* 2020; Dal Corso *et al.* 2022; Toulemonde *et al.*, this volume). Figure 3 shows the sites that have yielded 2nd millennium BCE AMS-dates on millet. In a number of other AMS-dated cases, the ‘early’ millet finds turned out to be (significantly) younger than the archaeological layers in which they were found. The intrusive finds are, as a rule, small in number (one to several grains in a deposit or a feature) and they almost always come from multi-period sites or those disturbed in recent times (by, for instance, modern agricultural activity). Large, high-density millet finds (termed ‘concentrations’) are much more likely to correspond to the age of the archaeological layer or context they came from. Several papers in this volume present the cases where millet finds fit the expected date (based on archaeological periodisation), and several present cases where there is a mismatch (Reed *et al.*; Hellmund; Toulemonde *et al.*). Thus, the caution raised in some previous studies is reiterated here:

1. Before suggesting the age of broomcorn millet finds based on the dating of other materials from the site, researchers should assess the possibility of contamination/intrusiveness, by considering the density of the finds, as well as the nature of the site (whether single-phase or multi-phases occupation), its proximity to later sites, and modern-day disturbance of the find location.

Combined, the radiocarbon dates and archaeobotanical finds of millet document the start of *cultivation* of this crop in Europe. The stable isotope analyses of human and animal tissues and the biomarker (miliacin) records from Europe also point to the mid-2nd millennium BCE as also the onset of the *consumption* of millet (*e.g.* Tafuri *et al.* 2009; Gerling 2015; Heron *et al.* 2016; Gamarra *et al.* 2018; Cavazzuti *et al.* 2019; Pospieszny *et al.* 2021; Standall *et al.* this volume). Taken together, these strands of evidence suggest that:

2. Millet arrived in Europe as a crop and was recognised as such; it seems unlikely that it was first a tolerated weed or accidental inclusion, as previously thought based on the low presence of ‘early’ millet at pre-Bronze Age sites. The status of crop, perhaps in some places similarly or more important than other staples, may in part explain the faster diffusion and adoption of the new cultivar compared with the spread of Neolithic crops several millennia earlier.

It has been stated before and emphasised in the papers here (Reed *et al.*; Hellmund; Pashkevych; Toulemonde *et al.*) that taking up millet and other new Bronze Age plant resources may have been a risk-buffering measure, a strategy that guaranteed sufficient food. This has been seen as particularly relevant in areas considered agriculturally marginal (*e.g.* with poor or shallow soils and/or low rainfall) or where millet may have enabled colonisation of parts of the landscape unsuitable for winter crops or more demanding crops (*e.g.* areas in wetlands accessible only during the dry/summer season). The contribution by Toulemonde *et al.* underscores how the areas uninhabited during the Early Bronze Age were colonised in the Middle-Late Bronze Age and how the incoming groups cultivated significant amounts of millet. This may have been the case in northern Germany too (Feaser *et al.* this volume). We can, therefore, propose that:




Figure 3. Sites in Europe from which broomcorn millet grains probably or possibly derive from the 2nd millennium BCE. The selection is based on the upper bound of 2a cal BCE range of the ^{14}C AMS-dates published in Filipović et al. (2020 [Supplementary Data]) and papers in this volume (Tolar and Pavlin; Toulemonde et al.); calibration using OxCal v4.4.2 and IntCal20 (Bronk Ramsey 2009; Reimer et al. 2020). Shaded areas indicate the geographical division used in Filipović et al. (2020) and the years BCE-labels note the modelled start of millet cultivation (base map © OpenStreetMap contributors; map: Dragana Filipović, Anna Carina Lange: 1. (Rhine-) Altenrheine, 2. Assiros Toumba, 3. Bodegaw 19, 4. Balloy, 5. Binningen, 6. (Friedberg-)Bruchenbrücken, 7. Butzow 10, 8. Cornești-Iarcuri, 9. Courcery, 10. Mačkovac- Crišnjevi, 11. Custaza, 12. Dikly Sad (Дикіу Сад), 13. Dragomelj, 14. Fajsz 18, 15. Fechenheim (Frankfurt am Main), 16. Gingsheim, 17. Gomolava, 18. Großbahren 4, 19. Grosuplje, 20. Guamsky Grot (Гуамський грот), 21. Hagnau(-Burg), 22. Hérange, 23. Hisar (Leskovač), 24. Ipš, 25. Jülich-Güsten, 26. Knittlingen 5, 27. (Lauda-)Königshofen, 28. Lavagnone, 29. Lipnik 5, 30. Lüdensen 6, 31. Lutomiensk, 32. Măgura-Buduiasca, 33. Maszkowice, 34. Mählow 5, 35. Neudorf, 36. (Hundsburg-) Olbetal, 37. Orehova vas, 38. Mačkovac-Oštrovi, 39. Posewalk 109, 40. Pécel 02, 41. Pisek-Sever, 42. Radis 1, 43. Réau, 44. Reznei, 45. Rullstorf 5, 46. Santa Giulia, 47. Schwerin 81, 48. Sézannes, 49. Sovi převís, 50. Stülfried an der March, 51. Szczepiadło 17, 52. Valečov 1, 53. Velim, 54. Vinogradnyi Sad (Виноградний Сад), 55. Vráblet-Velké Lehembý, 56. Wahlstedt, 57. Warendorf(-Einen), 58. (Hünenburg bei) Watenstedt, 59. Witow 1, 60. Zahájí.

3. Millet may have encouraged or facilitated (re-)colonisation of uninhabited or less inhabited areas and unused or underused parts of the landscape, such as those we tend to label as agriculturally marginal. Perhaps we can infer that the colonising groups indeed relied on millet to a great degree because of its good response to a range of growing conditions. This premise could lead to a conclusion that early millet growers in Europe were aware and took advantage of the potential of the new crop. Millet may have gone from being a choice to being a necessity, or it may have been both, depending on the context.

A general decrease in humidity in the course of the Bronze Age and increasing openness of the landscape have been mentioned as having stimulated or accelerated the adoption of millet (e.g. papers in this volume by Bartosiewicz; Filatova; Feeser *et al.*; see also Cremaschi *et al.* 2006, 2016; Tafuri *et al.* 2009; Dal Corso 2018; Feeser *et al.* 2019; Gumnior and Stobbe 2021). Perhaps the regions which gave the earliest radiocarbon dates in Europe were affected by these environmental trends and millet cultivation ‘saved’ parts of them from becoming decolonised. Approaching millet as an ‘enabler crop’ foregrounds its possible contribution to maintaining or increasing population numbers in some places. The middle and late 2nd millennium BCE indeed witnessed apparently growing numbers of settlements or cemeteries in, for instance, the North Pontic region, the Po River valley, the Carpathian Basin, northeastern France, and north-central Europe (Cardarelli 2009; Capuzzo *et al.* 2018; Kneisel *et al.* 2019; Pospieszny *et al.* 2021; Dal Corso *et al.* 2022; Toulemonde *et al.* this volume). With this in mind, and also that millet was effectively diversifying farming and cooking niches, we can hypothesise that:

4. The ‘millet effect’, embedded in the overall diversification in the Bronze Age of farming and culinary niches, as well as economic and social relations, extended beyond food production and consumption, into symbolism and ritual. There are in Europe quite a few examples of the early finds of millet in contexts that may suggest its potentially ritual use or symbolic role, such as within or in association with burials or ‘ritual pits’ (e.g. Bartík and Hajnalová 2004; Palmer 2007; Filipović *et al.* 2020, Supplementary Data; in this volume: Hellmund; Reed *et al.*; Toulemonde *et al.*).

Millet, so what: Why adopt millet into food production systems already relying on a range of crops?

The Middle and Late Bronze Age archaeobotanical records from different parts of Europe contain remains of a wide range of cultivated and wild plants found in settlements and cemeteries. Besides crops used from the Neolithic onwards, such as several types of wheat (*Triticum*) and different variants of barley (*Hordeum*), lentil (*Lens culinaris*), pea (*Pisum sativum*) and flax (*Linum usitatissimum*), several new species, in addition to broomcorn millet, were taken into cultivation in the later part of the period or towards its end, including gold-of-pleasure (*Camelina sativa*) and broad bean (*Vicia faba*), and perhaps *Lallemantia*, foxtail millet (*Setaria italica*) and safflower (*Carthamus*). The Middle Bronze Age was in some places characterised by a greater range of pulse crops, which, besides the earlier lentil and barley, also included bitter vetch (*Vicia ervilia*) and grass pea (*Lathyrus sativus*). Spelt wheat (*Triticum spelta*) seems to have been (more) regularly cultivated too. For wild resources, many Late Bronze Age sites show evidence of the collection of oak (acorns; *Quercus*), chenopods (*Chenopodium*) and perhaps wild crucifers (*Brassicaceae*) and large-seeded grasses (Poaceae) (e.g. Harding 1989; Jones and Valamoti 2005; Gyulai 2010; Pashkevych 2012; Stika and Heiss 2013; Kroll 2016; Perego 2017; Effenberger 2018; Šálková *et al.* 2019; papers in this volume). The fact that such a wide spectrum of edible plants was in use suggests that these were the times of plenty.

Food can become more abundant when environmental conditions are favourable. On the other hand, diversification of the subsistence base can be seen as a strategy for countering scarcity, which can, among other things, result from unfavourable growing conditions of plants due to extreme weather or harsh climate. The late Holocene is characterised by an overall decline in temperature, with warm or cold anomalies in some regions (e.g. Bradley and Bakke 2019). The Bronze Age overlaps with the Subboreal period, which is known for its unstable climate and at least two significant and adverse oscillations in the (Eastern) Mediterranean – the 4.2 ka BP and the 3.2 ka BP ‘events’ (documented as ‘megadroughts’; Kaniewski and van Campo 2017; Weiss 2017). It is unclear what consequences, if any, these and other climatic anomalies would have had in terms of plant food availability and acquisition in other parts of Europe.

Because the internal climate systems vary among regions, and the general (e.g. sub-continental) oscillations affect these regions in dissimilar ways, regional or micro-regional perspectives are necessary for the identification of possible correlations among climate, environment and plant consumption. The growing number of well-dated palaeoclimatic, palaeo-environmental and archaeological proxies create a platform for collaboration among these disciplines (Capuzzo *et al.* 2018; Kneisel *et al.* 2019; Weiberg *et al.* 2019; Schirrmacher *et al.* 2020). It is, therefore, to be expected that the recognition of any possible links between the developments in these different spheres will ensue (e.g. Gumnior and Stobbe 2021).

As highlighted in the introduction to this volume and in some of the contributions, the biology and ecology of broomcorn millet could have made it a highly attractive member of the Bronze Age crop-suite. Whether or not the first millet cultivators in Europe became instantly aware of the advantageous properties of millet – in other words, whether the knowledge of these properties arrived in Europe along with the crop or was acquired – cannot be inferred without more details on the growing conditions, both those imposed by the climate and weather and those created by people. Was millet a sought-after buffer crop from the start (as a fast-growing, drought-resistant, low-demanding supplement or alternative to the ‘traditional’ cultigens, wheat and barley) or had there been a phase of ‘learning’ its potential, an initial episode in the ‘cultural history of millet use’ (Weber and Fuller 2008)? Marijke Van der Veen contends that

‘... the introduction of a new crop may be started by allocating just a small part of the available land to it, allowing an assessment over one or more years before making the decision whether to switch wholesale to the new crop.’ (Van der Veen 2010, 3)

A much more populated ‘early millet map’ of Europe could shed some light on this issue, and would likely emphasise how varied the picture is depending on the local ecological and cultural factors, which likely differed within and between macro- and micro-regions. There have already been some indications of the uneven adoption or consumption of millet in the early days of its presence in Europe. For instance, stable isotopic data from the Bronze Age in the Po Basin, in northern Italy, show that the same community included both individuals who consumed millet and those who did not (Tafuri *et al.* 2009, 2018), and that among those who did, different individuals ate different amounts of millet (Cavazzuti *et al.* 2019). A similar situation is observed in southern Poland, where the stable isotope analysis of a series of Bronze Age burials revealed that individuals buried in the upland zone (the northern fringes of the Carpathian Basin) consumed millet at the time when members of the same community buried in the lowlands to the north did not. Both cultural and natural factors (elevation, latitude) may have dictated these differences (Pospieszny *et al.* 2021).

Curiously, according to the radiocarbon dates on millet (Filipović *et al.* 2020), latitudes north of ca. 50 parallel North in north-central Europe received or accepted millet later than those south of it. This apparent delay could be due to the natural constraints (i.e. cold climate, low availability of daylight) or different cultural

perceptions of millet, or both. With respect to the former, it would be worth taking a plant perspective and consider what kinds of adaptations (e.g. genetic mutations) were necessary for the millet plant to develop landraces able to succeed in conditions that differed from one area to another within its region of dispersal.

Instead of or in addition to natural factors necessitating broadening of the subsistence base, a long list of social and cultural factors and relations tend to shape the spread and adoption of introduced foods (Palmer and Van der Veen 2002; Van der Veen 2003, 2010; Hastorf 2017; Twiss 2019). They depend on, and at the same determine, the role of the innovations, which leads us to the next question.

Millet for what: Food, drink, fodder, symbol, currency, or all of the above?

Both millet and other contemporary crops have been used for multiple biological and technical purposes, i.e. human food or drink, animal graze or fodder, construction and craft material. Therefore, millet can archaeologically or historically be found in any of these roles (e.g. Moreno-Larrazabal *et al.* 2015; Teira-Brión this volume). Site- or region-specific studies explore the share of millet in food production and consumption, and we can expect further exciting discoveries in this area.

Culturally determined cooking methods and tastes influence the way millets are processed and prepared for consumption as food or drink – for instance dehusking, (par)boiling, malting, milling, mixing with other foods (e.g. de Garine 2001; Kohler-Schneider 2001, 153; Lundström-Baudais *et al.* 2002; Weber and Fuller 2008; Pashkevych this volume). Some archaeological evidence of millet processing and millet-containing food or drink has already been discovered (see Standall *et al.* this volume; Popov *et al.* 2018; Heiss *et al.* 2019). If not millet eating, certainly millet drinking would have in some instances been symbolically charged, such as at rituals, festivals and similar occasions of conspicuous consumption (cf. Palmer and Van der Veen 2002; Twiss 2019), although in some modern-day traditional communities, millet beer is perceived as food and consumed both day to day and on special occasions (de Garine 2001).

Stable carbon and nitrogen isotope measurements from Europe demonstrate that both humans and animals consumed millet from the very first days of its presence on the continent (e.g. Tafuri *et al.* 2009; Cavazzuti *et al.* 2019), humans doing so both directly and via animals. This suggests that millet had a similar dietary standing as other crops, and, as the biomolecular and anatomical analysis of food residues document, that it was apparently combined in meals with other ingredients, both of animal and of plant origin (Heron *et al.* 2016; Popov *et al.* 2018). Given its mild, indistinct taste, it could have been used to bulk up foods and drinks and give them a sticky texture, as in the Ukrainian/Russian dish *kulich* (Pashkevych this volume); or the ‘barley-millet-brome grass risotto’ in Early Iron Age Stillfried (Kohler-Schneider 2001, 153). It may also have been combined with animal foodstuffs as in Late Bronze Age Bruszczewo (Heron *et al.* 2016). And it may also have been used as a weaning food, such as at Iron Age La Hoya (Fernández-Crespo *et al.* 2019). The surviving tradition of millet cooking among the Cossack groups in Ukraine and of millet farming in northwestern Iberia (Moreno-Larrazabal *et al.* 2015; Teira-Brión this volume) exemplifies how the role(s) and importance of millet in diet construct its cultural meaning and significance and give it a place in cultural identity.

Approaching millet as a cultural category creates a possibility to examine the cultural affinity and status of millet farmers and/or consumers (cf. Palmer and Van der Veen 2002). As the modern-day examples described by Kirleis *et al.* (this volume) aptly illustrate, millet production and consumption can reflect individual or group social status, or a desire for social distinction; they can define and delineate economic, social and dietary identity.

Cropping system		Definition
Monoculture	Increased following mechanisation	One crop grown on the same plot for successive years.
Monocropping		One crop grown on one plot in a single year.
Multiple (,mixed') cropping		Two or more crops grown on the same plot in one year, simultaneously or sequentially.
	Sequential	More than one crop grown per plot per year, in sequence. Single or two/three crops grown per stage in the sequence.
	Intercropping	Two or more crops grown simultaneously on one plot (usually a combination of cereals and pulses).
	Mixed	Simultaneous cropping with random distribution.
	Row	Two or more crops per plot sown singly in separate rows.
	Strip	Two or more crops per plot sown singly in separate strips.
Relay intercropping		Simultaneous cropping for only part of crop life-cycles.
Ratoon cropping		Cultivation of crop regrowth after harvest.

Table 1. Summary of types of cropping system, modified after Buttler 1999, Table 24.1.

An interesting case described by Roland Hardenberg is that of the Dongria Kondh indigenous tribe now living in Odisha state, eastern India, whose members use pearl, finger and other locally grown millets as a currency – to pay for labour (Hardenberg 2021). This reminds us that early millet production need not necessarily have been geared (only) towards local consumption and that in some contexts millet could have been purely a cash crop (*sensu* Sherratt 1999). That its grain is lightweight and has a profitable ratio of grain sown to grain harvested would have been advantageous from the perspective of exchange. This aspect of millet re-opens an area of research on the value of staple crops as exchangeable items, as currencies. Further, it has implications for the understanding of the modes and mechanisms via which millet (and other organic commodities) were distributed, and how the networks circulating the products were formed and shaped by internal and external socio-economic and environmental factors. Coming from afar, broomcorn millet could have initially been perceived as an exotic food in parts of Europe (cf. Sherratt 1999; Van der Veen 2003; Boivin *et al.* 2012) and could have represented a valuable tradeable good.

We are inclined to believe that the new crop increased food security and, by extension, eased life for the agriculture-reliant prehistoric communities; that its adoption brought about a good thing and may have saved lives in times or places of food shortage. We should, however, not forget that the opportunity to produce more food would have increased the level of exploitation of the land and the workload of those working the land. Apart from thorough weeding early in the growing cycle, millet crops require little field management compared with most of the cereals, but their processing is similarly laborious to that of hulled wheats (Lundström-Baudais *et al.* 2002; Harvey and Fuller 2005; Weber and Fuller 2008). Where millet was sown regularly, and after winter or early spring crops, it would have added another set of agricultural tasks at sowing and harvesting time and required re-scheduling of the annual cycle revolving around winter crops. This would have placed pressure on the food producers.

We should be examining the archaeological and land-use records for signs of reorganisation of agricultural production at the time of and after the start of cultivation of millet and other new crops of the Bronze Age. Ann Buttler has conveniently summarised historical and ethnographic evidence of systems producing multiple crops, and described their advantages (Table 1; Buttler 1999). Her work offers an extremely useful framework for inspecting the archaeological and related evidence and inferring agricultural solutions. Some directions already transpire in the contributions to this volume. The archaeological (spatial and temporal) co-occurrence of millet and broad bean emphasised by Hellmund (this

volume) and also to some extent by Reed *et al.* (this volume) is a peculiar one and brings to mind intercropping of pearl millet and cowpea (*Vigna unguiculata*) in West Africa (Herrmann 2017; Styring *et al.* 2019). Intensification of arable production could have been done in this and other ways, including varied (*i.e.* more strategic) use of field manure, expansion into previously less-used or unused parts of the landscape, flexible fallowing and crop rotation regimes, use of more efficient tools and technologies. Thus, *ecological opportunism*, when viewed as a reason behind the adoption of millet in Eurasia, likely extended beyond the widening choice of crops and included pragmatic approaches to securing food and maximising returns (Jones *et al.* 2011; Liu *et al.* 2019).

In addition to the diversification, seen both in the range of plants cultivated or gathered and in the range of production techniques, we should expect to see indications of specialisation, probably aligned with agricultural tasks. Burhan Ulaş (2021) informs us that, until recently, there were specialised harvesters in southeastern Anatolia, hired in season throughout the region. Further back in the past, there also were specialised agricultural workers for sowing, threshing and grain-mill operating. They were paid in agricultural products (wheat, bulgur, flour). While moving around, they transmitted their knowledge and grains of different wheat varieties. In other words, specialised seasonal workforce served as vectors for the diffusion of agricultural techniques and resources in eastern Anatolia. This observation becomes even more interesting when we consider that this region is home to different Christian and Muslim ethnic groups (Ulaş 2021). This and other similar examples are particularly inspiring in the context of the traditional association of early millet (cultivation and/or consumption) with various mobile human groups that are a hallmark of the prehistory of central Asia and parts of eastern Europe (Frachetti *et al.* 2010; Spengler *et al.* 2014; Hermes *et al.* 2019; Dal Corso *et al.* 2022; Pashkevych this volume).

Conclusions and outlook

On its arrival, the new crop was incorporated into the economic, socio-environmental and ritual spheres. The papers in this book address different aspects of the introduction of millet in Europe. They also ask and spark a host of new questions about the cultural and environmental history of the integration of millet into food production systems, its economic, dietary, symbolic and ritual roles. Building on the already existing knowledge, possibilities open up for more in-depth investigations on broomcorn millet and other agricultural innovations in prehistoric Europe.

Despite the growing research on the arrival of millet and its early cultivation and consumption in Europe, the question *Where (exactly) from* is still open, and we have outlined several different possibilities here and in other papers (Filipović *et al.* 2020; Dal Corso *et al.* 2022). Answering this question requires further archaeobotanical finds and direct dating of broomcorn millet from regions situated at the interface of Asian and European cultural interaction.

Acknowledgements

The authors are indebted to the colleagues who were invited and/or took part, or otherwise assisted, in the initiation and completion of the *Millet Dating Programme*, the millet-themed workshop at Kiel and its proceedings. The authors are more than grateful to the many colleagues who aided in various ways. In particular, for their help with putting the workshop and the proceedings together, warm thanks are due to Almuth Alsleben, Marian Berihuete-Azorín, Jelena Bulatović, Nora Claussen, Yasmin Dannath, Walter Dörfler, Hermann Gorbahn, Felix Krause, Anna Carina

Lange, Ingo Feeser, John Meadows, Giedre Motuzaitė Matuzevičiūtė, Hiro Nasu, Suzanne Needs-Howarth, Roisin O'Droma, Merle Oelbüttel, Hendrik Raese, Kelly Reed, Tanja Reiser, Iris Reitzig, Ulrich Schmölcke, Janina Stampe, Robert Staniuk, Nicole Taylor, Tjaša Tolar, Françoise Toulemonde, Julian Wiethold, Fiona Walker-Friedrichs and Tanja Zerl. Funding for the *Millet Dating Programme*, the workshop and this book was provided by the German Research Foundation (DFG; project number 2901391021 – SFB 1266).

References

- Alberti, M.E. and Sabatini, S., eds., 2013. *Exchange Networks and Local Transformations: Interaction and Local Change in Europe and the Mediterranean from the Bronze Age to the Iron Age*. Oxford: Oxbow Books.
- Ananyevskaya, E., Akhatov, G., Loman, V., Dmitriev, E., Ermolayeva, A., Evdokimov, A., Garbaras, A., Goryachev, A., Kukushkin, A., Kukushkin, I., Kurmankulov, Z., Logvin, A., Lukpanova, Y., Onggar, A., Sakenov, S., Sapolaite, J., Shevnina, I., Usmanova, E., Utubayev, Z., Varfolomeev, V., Voyakin, D., Yarygin, S., Motuzaitė Matuzeviciute, G., 2020. The effect of animal herding practices on the diversity of human stable isotope values in north central Asia. *Journal of Archaeological Science: Reports*, 34, 102615. <https://doi.org/10.1016/j.jasrep.2020.102615>
- Anthony, D.W., 1995. Horse, wagon & chariot: Indo-European languages and archaeology. *Antiquity*, 69 (264), 554-565. <https://doi.org/10.1017/S0003598X00081941>
- Anthony, D.W., 2007. *The Horse, the Wheel, and Language: How Bronze-Age Riders from the Eurasian Steppes Shaped the Modern World*. Princeton: Princeton University Press.
- Anthony, D.W., Brown, D., Brown, E., Goodman, A., Khokhlov, A., Kosintsev, P., Kuznetsov, P., Mochalov, O., Murphy, E., Peterson, D., Pike-Tay, A., Popova, L., Rosen, A., Russell, N., Weisskopf, A., 2005. The Samara Valley Project: Late Bronze Age economy and ritual in the Russian steppes. *Eurasia Antiqua*, 11, 395-417.
- Balasse, M., Tornero, C., Bréhard, S., Ughetto-Monfrin, J., Voinea, V., Bălăşescu, A., 2014. Cattle and sheep herding at Cheia, Romania, at the turn of the fifth millennium cal BC: A view from stable isotope analysis. *Proceedings of the British Academy*, 198, 115-142. <https://doi.org/10.5871/bacad/9780197265758.001.0001>
- Balasse, M., Bălăşescu, A., Tornero, C., Fremondeau, D., Hovsepyan, R., Gillis, R., Popovici, D., 2017. Investigating the scale of herding in Chalcolithic pastoral communities settled along the Danube River in the 5th millennium BC: a case study at Borduşani-Popină and Hârşova-tell (Romania). *Quaternary International*, 436, 29-40. <https://doi.org/10.1016/j.quaint.2015.07.030>
- Bartík, J. and Hajnalová, M., 2004. Hromadný nález keramiky z Lozorna [Large find of pottery from Lozorno]. *Zborník Slovenského Národného Múzea XCVIII – Archeológia*, 14, 9-42.
- Bartosiewicz, L., 2013. Animals in Bronze Age Europe. In: H. Fokkens and A. Harding, eds., *Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 328-347.
- Behre, K.E., 1998. Landwirtschaftliche Entwicklungslinien und die Veränderung der Kulturlandschaft in der Bronzezeit Europas. In: B. Hänsel, ed., *Mensch und Umwelt in der Bronzezeit Europas*. Kiel: Oetker-Voges Verlag, 91-109.
- Bending, J. and Colledge, S., 2007. The archaeobotanical assemblages. In: N. Postgate and D. Thomas, eds., *Excavations at Kilise Tepe, 1994-98: from Bronze Age to Byzantine in Western Cilicia*. Cambridge: McDonald Institute Monographs, 583-595.
- Benecke, N., 1998. Haustierhaltung, Jagd und Kult mit Tieren im bronzezeitlichen Mitteleuropa. In: B. Hänsel, ed., *Mensch und Umwelt in der Bronzezeit Europas*. Kiel: Oetker-Voges Verlag, 61-75.

- Bergerbrant, S., 2013. Migration, innovation and meaning: sword depositions on Lolland, 1600-1100 BC. In: M.E. Alberti and S. Sabatini, eds., *Exchange Networks and Local Transformations: Interaction and Local Change in Europe and the Mediterranean from the Bronze Age to the Iron Age*. Oxford: Oxbow Books, 146-155.
- Boivin, N., Fuller, D.Q., Crowther, A., 2012. Old World globalization and the Columbian exchange: comparison and contrast. *World Archaeology*, 44 (3), 452-469. <https://doi.org/10.1080/00438243.2012.729404>
- Bökönyi, S., 1974. *History of Domestic Mammals in Central and Eastern Europe*. Budapest: Akadémiai Kiadó.
- Bouzek, J., 1985. *The Aegean, Anatolia and Europe: Cultural Interrelations in the Second Millennium B.C.* Studies in Mediterranean Archaeology 29. Göteborg: Paul Åströms Förlag/Prague: Academia.
- Bradley, R.S. and Bakke, J., 2019. Is there evidence for a 4.2 ka BP event in the northern North Atlantic region? *Climate of the Past*, 15, 1665-1676. <http://dx.doi.org/10.5194/cp-15-1665-2019>
- Bronk Ramsey, C., 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon*, 51 (1), 337-360. <https://doi.org/10.1017/S0033822200033865>
- Broodbank, C., 2016. The transmitting sea: a Mediterranean perspective. In: E. Kiriati and C. Knappett, eds., *Human Mobility and Technological Transfer in the Prehistoric Mediterranean*. Cambridge: Cambridge University Press, 18-30.
- Buttler, A., 1999. Traditional seed cropping systems in the temperate Old World: models for antiquity. In: C. Gosden and J. Hather, eds., *The Prehistory of Food: Appetites for Change*. London: Routledge, 452-467.
- Bryce, T., 2005. *Life and Society in the Hittite World*. Oxford: Oxford University Press.
- Cardarelli, A., 2009. The collapse of the Terramare culture and growth of new economic and social systems during the Late Bronze Age in Italy. *Scienze dell'Antichità: Storia Archeologia Antropologia*, 15, 449-519.
- Cardarelli, A., Cavazzuti, C., Fritzl, M., Gavranovic, M., Hajdu, T., Kiss, V., Köhler, K., Kulcsár, G., Melis, E., Rebay-Salisbury, K., Szabó, G., Szeverényi, V., 2020. The connections between the plains of the Po and the Danube during the Bronze Age seen through the spread of the 'urnfield model'. *Rivista die Scienze Preistoriche LXX S1 – Italia tra Mediterraneo ed Europa: Mobilità, Interazioni e Scambi*. Firenze: Istituto Italiano di Preistoria e Protostoria, 231-243.
- Capuzzo, G., Zanon, M., Dal Corso, M., Kirleis, W., Barcelo, J.A., 2018. Highly diverse Bronze Age population dynamics in central-southern Europe and their response to regional climatic patterns. *PLoS ONE*, 13 (8), e0200709. <https://doi.org/10.1371/journal.pone.0200709>
- Cavazzuti, C., Skeates R., Millard, A.R., Nowell, G., Peterkin, J., Brea, M.B., Cardarelli, A., Salzani, L., 2019. Flows of people in villages and large centres in Bronze Age Italy through strontium and oxygen isotopes. *PLoS ONE*, 14 (1), e0209693. <https://doi.org/10.1371/journal.pone.0209693>
- Chernykh, E.N., 1992. *Ancient Metallurgy in the USSR: the Early Metal Age*. Cambridge: Cambridge University Press.
- Cremaschi, M., Pizzi, C., Valsecchi, V., 2006. Water management and land use in the Terramare and a possible climatic co-factor in their abandonment: the case study of the Terramara of Poviglio Santa Rosa (northern Italy). *Quaternary International*, 151, 87-98. <https://doi.org/10.1016/j.quaint.2006.01.020>
- Cremaschi, M., Mercuri, A.M., Torri, P., Florenzano, A., Pizzi, C., Marchesina, M., Zerboni, A., 2016. Climate change versus land management in the Po Plain (northern Italy) during the Bronze Age: new insights from the VP/VG sequence of the Terramara Santa Rosa di Poviglio. *Quaternary Science Reviews*, 136, 153-172. <https://doi.org/10.1016/j.quascirev.2015.08.011>

- Dal Corso, M., 2018. *Environmental History and Development of a Northeastern Italian Lowland During the Bronze Age: a Multi-Disciplinary Case-Study*. Universitätsforschungen zur Prehistorischen Archäologie 312. Bonn: Dr. Rudolf Habelt GmbH.
- Dal Corso, M., Kirleis, W., Kneisel, J., Taylor, N., Wieckowska-Lüth, M., Zanon, M., eds., 2019. *How's Life? Living Conditions in the 2nd and 1st Millennia BCE*. Leiden: Sidestone Press.
- Dal Corso, M., Pashkevych, G., Filipović, D., Liu, X., Motuzaitė-Matuzeviciute, G., Stobbe, A., Shatilo, L., Videiko, M., Kirleis W., 2022. Between cereal agriculture and animal husbandry – millet in the early economy of the North Pontic region. *Journal of World Prehistory*.
- Dalla Longa, E., Dal Corso, M., Vicenzutto, D., Nicosia, C., Cupitò, M., 2019. The Bronze Age settlement of Fondo Paviani (Italy) in its territory: hydrography, settlement distribution, environment and in-site analysis. *Journal of Archaeological Science: Reports*, 28, 102018. <https://doi.org/10.1016/j.jasrep.2019.102018>
- Diffey, C., Neef, R., Bogaard, A., 2017. The archaeobotany of large-scale hermetic cereal storage at the Hittite capital of Hattusha. In: A. Schachner, ed., *Innovation versus Beharrung: Was macht den Unterschied des hethitischen Reichs im Anatolien des 2. Jahrtausends v. Chr.?* Byzas 23. Istanbul: Ege Yayınları, 185-202.
- Dörfler, W., Herking, C., Neef, R., Pasternak, R., von den Driesch, A., 2011. Environment and economy in Hittite Anatolia. In: H. Genz and D.P. Mielke, eds., *Insights into Hittite History and Archaeology*. Colloquia Antiqua 2, 99-124.
- Dreslerová, D., Kozáková, R., Metlička, M., Brychová, V., Bobek, P., Čišecký, Č., Demján, P., Lisá, L., Pokorná, A., Michálek, J., Strouhalová, B., Trubač, J., 2020. Seeking the meaning of a unique mountain site through a multidisciplinary approach: the late La Tène site at Sklářské valley, Šumava Mountains, Czech Republic. *Quaternary International*, 542, 88-108. <https://doi.org/10.1016/j.quaint.2020.03.013>
- Effenberger, H., 2018. The plant economy of the northern European Bronze Age – more diversity through increased trade with southern regions. *Vegetation History and Archaeobotany*, 27, 65-74. <https://doi.org/10.1007/s00334-017-0621-3>
- Fairbairn, A., 2014. Preliminary archaeobotanical investigations of plant production, consumption, and trade at Bronze Age Kültepe-Kanesh. In: L. Atici, F. Kulakoğlu, G. Barjamovic, A. Fairbairn, eds., *Current Research in Kültepe-Kanesh*. Journal of Cuneiform Studies – Supplemental Series 4. Atlanta: Lockwood Press, 177-193.
- Fairbairn, A.S. and Bradley, K., 2008. Archaeobotany at Kaman-Kalehöyük 2007. *Kaman-Kalehöyük*, 17, 195-199.
- Fairbairn, A., Kulakoğlu, F., Atici, L., 2013. Archaeobotanical evidence for trade in hazelnut (*Corylus* sp.) at Middle Bronze Age Kültepe (c. 1950-1830 B.C.), Kayseri Province, Turkey. *Vegetation History and Archaeobotany*, 23, 167-174. <https://doi.org/10.1007/s00334-013-0403-5>
- Fairbairn, A.S., Wright, N.J., Weeden, M., Barjamovic, G., Matsumura, K., Rasch, R., 2018. Ceremonial plant consumption at Middle Bronze Age Büklükale, Kırıkkale Province, central Turkey. *Vegetation History and Archaeobotany*, 28, 327-346. <https://doi.org/10.1007/s00334-018-0703-x>
- Feaser, I., Dörfler, W., Kneisel, J., Hinz, M., Dreibrodt, S., 2019. Human impact and population dynamics in the Neolithic and Bronze Age: multi-proxy evidence from north-western central Europe. *The Holocene*, 29, 1596-1606. <https://doi.org/10.1177%2F0959683619857223>
- Fernández-Crespo, T., Ordoño, J., Bogaard, A., Llanos, A., Schulting, R., 2019. A snapshot of subsistence in Iron Age Iberia: the case of La Hoya village. *Journal of Archaeological Science: Reports*, 28, 102037. <https://doi.org/10.1016/j.jasrep.2019.102037>
- Filipović, D., 2014. *Early Farming in Central Anatolia – an Archaeobotanical Study of Crop Husbandry, Animal Diet and Land Use at Neolithic Çatalhöyük*. Oxford: Archaeopress.

- Filipović, D., Meadows, J., Dal Corso, M., Kirleis, W., Alsleben, A., Akeret, Ö., Bittmann, F., Bosi, G., Ciută, B., Dreslerová, D., Effenberger, H., Gyulai, F., Heiss, A.G., Hellmund, M., Jahns, S., Jakobitsch, T., Kapcia, M., Kloof, S., Kohler-Schneider, M., Kroll, H., Makarowicz, P., Marinova, E., Märkle, T., Medović, A., Mercuri, A.M., Mueller-Bieniek, A., Nisbet, R., Pashkevych, G., Perego, R., Pokorný, P., Pospieszny, Ł., Przybyła, M., Reed, K., Rennwanz, J., Stika, H.-P., Stobbe, A., Tolar, T., Wasylkowa, K., Wiethold, J., Zerl, T., 2020. New AMS ^{14}C dates track the arrival and spread of broomcorn millet cultivation and agricultural change in prehistoric Europe. *Scientific Reports*, 10, 13698. <https://doi.org/10.1038/s41598-020-70495-z>
- Fischl, K.P., Kiss, V., Kulcsár, G., Szeverényi, V., 2013. Transformations in the Carpathian Basin around 1600 B.C. In: H. Meller, F. Bertemes, H.R. Bork, R. Risch, eds., *1600 – Kultureller Umbruch im Schatten des Thera-Ausbruchs? / 1600 – Cultural Change in the Shadow of the Thera-Eruption?* Tagungen des Landesmuseums für Vorgeschichte Halle 9. Halle (Saale): Landesamt für Denkmalpflege und Archäologie Sachsen-Anhalt, Landesmuseum für Vorgeschichte Halle (Saale), 355-371.
- Fokkens, H. and Harding, A., eds., 2013 *Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press.
- Frachetti, M.D., 2008. *Pastoralist Landscapes and Social Interaction in Bronze Age Eurasia*. Berkeley: University of California Press.
- Frachetti, M.D., 2012. Multiregional emergence of mobile pastoralism and nonuniform institutional complexity across Eurasia. *Current Anthropology*, 53 (1), 2-38. <https://doi.org/10.1086/663692>
- Frachetti, M.D., Spengler, R.S., Fritz, G.Y., Mar'yashev, A.N., 2010. Earliest evidence of broomcorn millet and wheat in the central Eurasian steppe region. *Antiquity*, 84, 993-1010. <https://doi.org/10.1017/S0003598X0006703X>
- Fuchs, K., 2020. *Interdisciplinary Analysis of the Cemetery Kudachurt 14: Evaluating Indicators of Social Inequality, Demography, Oral Health and Diet during the Bronze Age Key Period 2200-1650 BCE in the Northern Caucasus*. Leiden: Sidestone Press.
- Galaty, M.L., Tomas, H., Parkinson, W.A., 2014. Bronze Age European elites: from the Aegean to the Adriatic and back again. In: A.B. Knapp and P. Van Dommelen, eds., *The Cambridge Prehistory of the Bronze and Iron Age Mediterranean*. Cambridge: Cambridge University Press, 157-177.
- Gamarra, B., Howcroft, R., McCall, A., Dani, J., Hajdu, Z., Nagy, E.G., Szabo, L.D., Domboróczki, L., Pap, I., Raczky, P., Marcsik, A., Zoffmann, Z.K., Hajdu, T., Feeney, R.N.M., Pinhasi, R., 2018. 5000 years of dietary variations of prehistoric farmers in the Great Hungarian Plain. *PLoS ONE*, 13 (5), e0197214. <https://doi.org/10.1371/journal.pone.0197214>
- Garine, E. de, 2001. An ethnographic account to the many roles of millet beer in the culture the Duupa agriculturalists (Poli Mountains, northern Cameroon). In: I. Garine and V. Garine, eds., *Drinking: An Anthropological Approach*. Oxford: Berghan, 191-204.
- Gerling, C., 2015. *Prehistoric Mobility and Diet in the West Eurasian Steppes 3500 to 300 BC: an Isotopic Approach*. Topoi – Berlin Studies of the Ancient World 25. Berlin: Walter de Gruyter.
- Gimbutas, M., 1965. *Bronze Age Cultures in Central and Eastern Europe*. The Hague: Mouton & Co.
- Gorbenko, K.V. and Grebennikov, Y.S., 2009. The 'Dykyi Sad' fortified settlement as a uniting link in the context of economic, political and cultural relation of 1200-1000 BC (the Baltic shore, the Pontic region and the Mediterranean). *Baltic-Pontic Studies*, 14, 375-390.
- Greenfield, H., 2005. A reconsideration of the Secondary Products Revolution in south-eastern Europe: on the origins and use of domestic animals for milk, wool, and traction in the central Balkans. In: J. Mulville and A.K. Outram, eds.,

- The Zooarchaeology of Fats, Oils, Milk and Dairying*. Proceedings of the 9th Conference of the International Council of Archaeozoology, Durham, August 2002. Oxford: Oxbow, 14-31.
- Gumnior, M. and Stobbe, A., 2021. First palynological results from lowland sites in the Romanian Banat and their implications for settlement and land use dynamics in the southeastern Carpathian Basin. *Quaternary International*, 583, 48-61. <https://doi.org/10.1016/j.quaint.2021.02.005>
- Gyulai, F., 2010. *Archaeobotany in Hungary: Seed, Fruit, Food and Beverage Remains in the Carpathian Basin: an Archaeobotanical Investigation of Plant Cultivation and Ecology from the Neolithic to the Late Middle Ages*. Budapest: Archaeolingua.
- Halstead, P., 1987. Man and other animals in later Greek prehistory. *Annual of the British School of Archaeology at Athens*, 82, 71-83.
- Halstead, P., 1996. Pastoralism or household herding? Problems of scale and specialisation in early Greek animal husbandry. *World Archaeology*, 28 (1), 20-42. <https://doi.org/10.1080/00438243.1996.9980329>
- Halstead, P. and Isaakidou, V., 2011. Revolutionary secondary products: the development and significance of milking, animal-traction and wool-gathering in later prehistoric Europe and the Near East. In: T. Wilkinson, S. Sherratt, J. Bennet, eds., *Interweaving Worlds: Systemic Interactions in Eurasia, 7th to the 1st Millennia BC*. Oxford: Oxbow, 61-76.
- Halstead, P. and Isaakidou, V., 2017. Sheep, sacrifices, and symbols: animals in Later Bronze Age Greece. In: U. Albarella, M. Rizzetto, H. Russ, K. Vickers, S. Viner-Daniels, eds., *The Oxford Handbook of Zooarchaeology*. Oxford: Oxford University Press, 113-126.
- Hardenberg, R., (2021). The rise of a new millet assemblage in India. In: T. Schade, B. Schweizer, S. Teuber, R. Da Vela, W. Frauen, M. Karami, D. Kumar Ojha, K. Schmidt, R. Sieler, M. Toplak, eds., *Exploring Resources: on Cultural, Spatial and Temporal Dimensions of ResourceCultures*. RessourcenKulturen 13. Tübingen: Tübingen University Press.
- Harding, A.F., 1989. Interpreting the evidence for agricultural change in the Late Bronze Age in northern Europe. In: H.-A. Nordström and A. Knape, eds., *Bronze Age Studies*. Transactions of the British-Scandinavian Colloquium in Stockholm, May 10-11, 1985. Museum of National Antiquities Studies 6. Stockholm: Museum of National Antiquities, 173-181.
- Harding, A.F., 2000. *European Societies in the Bronze Age*. Cambridge: Cambridge University Press.
- Harding, A., 2013. World systems, cores and peripheries in pre-historic Europe. *European Journal of Archaeology*, 16, 378-400. <https://doi.org/10.1179/1461957113Y.0000000032>
- Harvey, E.L. and Fuller, D.Q., 2005. Investigating crop processing using phytolith analysis: the example of rice and millets. *Journal of Archaeological Science*, 32, 739-752. <https://doi.org/10.1016/j.jas.2004.12.010>
- Hastorf, C.A., 2017. *The Social Archaeology of Food*. Cambridge: Cambridge University Press.
- Heiss, A.G., Antolín, F., Azorín, M.B., Biederer, B., Erlach, R., Gail, N., Griebel, M., Linke, R., Lochner, M., Marinova, E., Oberndorfer, D., Stika, H.-P., Valamoti, S.M., 2019. The Hoard of the Rings: 'Odd' annular bread-like objects as a case study for cereal-product diversity at the Late Bronze Age hillfort site of Stillfried (Lower Austria). *PLoS ONE*, 14 (6), e0216907. <https://doi.org/10.1371/journal.pone.0216907>
- Herrmann, L., 2017. Elemente von Landwirtschaftssystemen aus sich entwickelnden Ländern Afrikas und Asiens zu Unterstützung der Modellbildung für den neolithischen Anbau in Zentraleuropa. In: J. Lechterbeck and E. Fischer, eds., *Festschrift für Manfred Rösch*. UPA 300. Bonn: Dr. Rudolf Habelt GmbH, 165-180.

- Hermes, T.R., Frachetti, M.D., Doumani Dupuy, P.N., Mar'yashev, A., Nebel, A., Makarewicz, C.A., 2019. Early integration of pastoralism and millet cultivation in Bronze Age Eurasia. *Proceedings of the Royal Society, B: Biological Sciences*, 286 (1910), 20191273. <https://doi.org/10.1098/rspb.2019.1273>
- Hermuth, J., Janovská, D., Hlásná Čepková, P., Ušák, S., Stražil, Z., Dvořáková, Z., 2016. Sorghum and foxtail millet – promising crops for the changing climate in central Europe. In: P. Konvalina, ed., *Alternative Crops and Cropping Systems*. London: IntechOpen Limited, 3-28.
- Heron, C., Shoda, S., Barcons, A.B., Czebreszuk, J., Eley, Y., Gorton, M., Kirleis, W., Kneisel, J., Lucquin, A., Müller, J., Nishida, Y., Son, J., Craig, O.E., 2016. First molecular and isotopic evidence of millet processing in prehistoric pottery vessels. *Scientific Reports*, 6, 38767. <https://doi.org/10.1038/srep38767>
- Herrscher, E., André, G., Bodet, C., Chataigner, C., Decaix, A., Goude, G., Hamon, C., Le Mort, F., Lyonnet, B., Martin, L., Messenger, E., Oberlin, C., Ollivier, V., Poulmarc'h, M., Sermet, C., Vila, E., 2018a. The origins of millet cultivation in the Caucasus: archaeological and archaeometric approaches. *Préhistoires Méditerranéennes*, 6, 1-27.
- Herrscher, E., Poulmarc'h, M., Pecqueur, L., Jovenet, E., Benecke, N., Decaix, A., Lyonnet, B., Guliyev, F., André, G., 2018b. Dietary inferences through stable isotope analysis at the Neolithic and Bronze Age in the southern Caucasus (sixth to first millennium BC, Azerbaijan): from environmental adaptation to social impacts. *American Journal of Physical Anthropology*, 164, 1-20. <https://doi.org/10.1002/ajpa.23718>
- Hingh, A.E. de, 2000. *Food Production and Food Procurement in the Bronze Age and Early Iron Age (2000-500 BC)*. Archaeological Studies, Leiden University 7. Leiden: Leiden University.
- Hoffner, H.A., 2003. Daily life among the Hittites. In: R. Averbeck, M.W. Chavalas, D.B. Weisberg, eds., *Life and Culture in the Ancient Near East*. Bethesda: CDL Press, 95-118.
- Hollund, H.I., Higham, T., Belinskij, A., Korenovskij, S., 2010 Investigation of palaeodiet in the north Caucasus (south Russia) Bronze Age using stable isotope analysis and AMS dating of human and animal bones. *Journal of Archaeological Science*, 37, 2971-2983. <https://doi.org/10.1016/j.jas.2010.08.009>
- Honch, N.V., Higham, T.F.G., Chapman, J., Gaydarska, B., Hedges, R.E.M., 2006. A palaeodietary investigation of carbon ($^{13}\text{C}/^{12}\text{C}$) and nitrogen ($^{15}\text{N}/^{14}\text{N}$) in human and faunal bones from the Copper Age cemeteries of Varna I and Durankulak, Bulgaria. *Journal of Archaeological Science*, 33, 1493-1504. <https://doi.org/10.1016/j.jas.2006.02.002>
- Hörander, E., ed., 1995. *Millet – Hirse – Millet*. Actes du Congrès d'Aizenay, 18-19 août 1990. Grazer Beiträge zur Europäischen Ethnologie 4. Frankfurt am Main: Peter Lang.
- Hovsepian, R., 2015. On the agriculture and vegetal food economy of Kura-Araxes culture in the south Caucasus. *Paléorient*, 41, 69-82.
- Hunt, H.V., Vander Linden, M., Liu, X., Motuzaite-Matuzeviciute, G., Colledge, S., Jones, M.K., 2008. Millet across Eurasia: chronology and context of early records of the genera *Panicum* and *Setaria* from archaeological sites in the Old World. *Vegetation History and Archaeobotany*, 17, S5-S18. <https://dx.doi.org/10.1007%2Fs00334-008-0187-1>
- Irvine, B., Erdal, Y.S., Richards, M.P., 2019. Dietary habits in the Early Bronze Age (3rd millennium BC) of Anatolia: a multi-isotopic approach. *Journal of Archaeological Science: Reports*, 24, 253-263. <https://doi.org/10.1016/j.jasrep.2019.01.015>
- Jockenhövel, A., 1991. Räumliche Mobilität von Personen in der mittleren Bronzezeit im westlichen Mitteleuropa. *Germania*, 69, 49-62.
- Jones, G. and Valamoti, S.M., 2005. *Lallemantia*, an imported or introduced oil plant in Bronze Age northern Greece. *Vegetation History and Archaeobotany*, 14, 571-577. <https://doi.org/10.1007/s00334-005-0004-z>

- Jones, G., Wardle, K., Halstead, P., Wardle, D., 1986. Crop storage at Assiros. *Scientific American*, 254 (3), 96-103.
- Jones, M., Hunt, H., Lightfoot, E., Lister, D., Liu, X., Motuzaite-Matuzeviciute, G., 2011. Food globalization in prehistory. *World Archaeology*, 43 (4), 665-675. <https://doi.org/10.1080/00438243.2011.624764>
- Jones, M., Hunt, H., Kneale, C., Lightfoot, E., Lister, D., Liu, X., Motuzaite-Matuzeviciute, G., 2016. Food globalisation in prehistory: the agrarian foundations of an interconnected continent. *Journal of the British Academy*, 4, 73-87. <https://doi.org/10.5871/jba/004.073>
- Jones, R.E., Levi, S.T., Bettelli, M., Vagnetti, L., 2014. *Italo-Mycenaean Pottery: the Archaeological and Archaeometric Dimensions*. Rome: CNR Istituto di Studi sul Mediterraneo Antico.
- Kaniewski, D. and van Campo, E., 2017. 3.2 ka BP megadrought and the Late Bronze Age collapse. In: H. Weiss, ed., *Megadrought and Collapse: From Early Agriculture to Angkor*. New York: Oxford University Press, 161-182. <https://doi.org/10.1093/oso/9780199329199.003.0005>
- Knapp, A.B. and Van Dommelen, P., eds., 2014. *The Cambridge Prehistory of the Bronze and Iron Age Mediterranean*. Cambridge: Cambridge University Press.
- Kneisel, J., Dal Corso, M., Kirleis, W., Taylor, N., Tiedke, V., 2015. The third food revolution? Common trends in economic and subsistence strategies in Bronze Age Europe. In: J. Kneisel, W. Kirleis, M. Dal Corso, H. Scholz, N. Taylor, V. Tiedtke, eds., *The Third Food Revolution? Setting the Bronze Age Table: Common Trends in Economic and Subsistence Strategies in Bronze Age Europe*. Proceedings of the International Workshop 'Socio-Environmental Dynamics over the Last 12,000 Years: The Creation of Landscapes III' (15th-18th April 2013) in Kiel. Universitätsforschungen zur prähistorischen Archäologie 623. Bonn: Dr. Rudolf Habelt GmbH, 275-287.
- Kneisel, J., Dörfler, W., Dreibrodt, S., Schaefer-Di Maida, S., Feeser, I., 2019. Cultural change and population dynamics during the Bronze Age: integrating archaeological and palaeoenvironmental evidence for Schleswig-Holstein, northern Germany. *The Holocene*, 29, 1607-1621. <https://doi.org/10.1177%2F0959683619857237>
- Knipper, C., Reinhold, S., Gresky, J., Berezina, N., Gerling, C., Pichler, S.L., Buzhilova, A.P., Kantorovich, A.R., Maslov, V.E., Petrenko, V.G., Lyakhov, S.V., Kalmykov, A.A., Belinskiy, A.B., Hansen, S., Alt, K.W., 2020. Diet and subsistence in Bronze Age pastoral communities from the southern Russian steppes and the north Caucasus. *PLoS ONE*, 15 (10), e0239861. <https://doi.org/10.1371/journal.pone.0239861>
- Kohler-Schneider, M., 2001. *Verkohlte Kultur- und Wildpflanzenreste aus Stillfried and der March als Spiegel bronzezeitlicher Landwirtschaft im Weinviertel, Niederösterreich*. Vienna: Verlag der Österreichischen Akademie der Wissenschaften.
- Kristiansen, K., 2016. Interpreting Bronze Age trade and migration. In: E. Kiriati and C. Knappett, eds., *Human Mobility and Technological Transfer in the Prehistoric Mediterranean*. Cambridge: Cambridge University Press, 154-180.
- Kristiansen, K. and Larsson, T.B., 2005. *The Rise of Bronze Age Society: Travels, Transmission and Transformations*. Cambridge: Cambridge University Press.
- Kristiansen, K. and Suchowska-Ducke, P., 2015. Connected histories: the dynamics of Bronze Age interaction and trade 1500-1100 BC. *Proceedings of the Prehistoric Society*, 81, 361-392. <https://doi.org/10.1017/ppr.2015.17>
- Kroll, H., 2016. Die Pflanzenfunde von Feudvar. In: H. Kroll and K. Reed, eds., *Die Archäobotanik*. Würzburg: Würzburg University Press, 37-194.
- Lacomte, O., 2012. *Ulug Depe: a Forgotten City in Central Asia: MAFTUR, Mission Archéologique Franco-Turkmène*. Paris: MAE, CNRS.
- Leipe, C., Long, T., Sergusheva, E.A., Wagner, M., Tarasov, P.E., 2019. Discontinuous spread of millet agriculture in eastern Asia and prehistoric population dynamics. *Science Advances*, 5 (9), eaax6225. <https://doi.org/10.1126/sciadv.aax6225>

- Liu, X., Jones, P.J., Motuzaite Matuzeviciute, G., Hunt, H.V., Lister, D.L., An, T., Przelomska, N., Kneale, C.J., Zhao, Z., Jones, M.K., 2019. From ecological opportunism to multi-cropping: mapping food globalisation in prehistory. *Quaternary Science Reviews*, 206, 21-28. <https://doi.org/10.1016/j.quascirev.2018.12.017>
- Lundström-Baudais, K., Rachoud-Schneider, A.-M., Baudais, D., Poissonnier, B., 2002. Le broyage dans la chaîne de transformation du millet (*Panicum miliaceum*): outils, gestes et écofacts. In: H. Procopiou and R. Treuil, eds., *Moudre et broyer: L'interprétation fonctionnelle de l'outillage de mouture et de broyage dans la Préhistoire et l'Antiquité: I: Méthodes*. Paris: Éditions du CTHS, 180-208.
- Marinval, P., 1992. Archaeobotanical data on millets (*Panicum miliaceum* and *Setaria italica*) in France. *Review of Palaeobotany and Palynology*, 73, 259-270. [https://doi.org/10.1016/0034-6667\(92\)90062-L](https://doi.org/10.1016/0034-6667(92)90062-L)
- Martin, L., Messenger, E., Bedianashvili, G., Rusishvili, N., Lebedeva, E., Longford, C., Hovsepyan, R., Bitadze, L., Chkadua, M., Vanishvili, N., Le Mort, F., Kakhiani, K., Abramishvili, M., Gogochuri, G., Murvanidze, B., Giunashvili, G., Licheli, V., Salavert, A., Andre, G., Herrscher, E., 2021. The place of millet in food globalization during Late Prehistory as evidenced by new bioarchaeological data from the Caucasus. *Scientific Reports* 11, 13124. <https://doi.org/10.1038/s41598-021-92392-9>
- Miller, N.F., 1982. Economy and Environment of Malayan, a third millennium B.C. Urban Center in Southern Iran. Doctoral dissertation, University of Michigan.
- Miller, N.F., 2015. Rainfall seasonality and the spread of millet cultivation in Eurasia. *Iranian Journal of Archaeological Studies*, 5, 1-10. <https://dx.doi.org/10.22111/ijas.2015.3015>
- Miller, N.F., Spengler, R.N., Frachetti, M., 2016. Millet cultivation across Eurasia: origins, spread, and the influence of seasonal climate. *The Holocene*, 26, 1566-1575. <https://doi.org/10.1177%2F0959683616641742>
- Moreno-Larrazabal, A., Teira-Brión, A., Sopelana-Salcedo, I., Arranz-Otaegui, A., Zapata, L., 2015. Ethnobotany of millet cultivation in the north of the Iberian Peninsula. *Vegetation History and Archaeobotany*, 24, 541-554. <https://doi.org/10.1007/s00334-015-0518-y>
- Motuzaitė-Matuzevičiūtė, G. and Liu, X., 2021. Prehistoric agriculture in China: food globalization in prehistory. In: H.H. Shugart, ed., *Oxford Research Encyclopedia of Environmental Science*. Oxford: Oxford University Press, 1-23. <https://doi.org/10.1093/acrefore/9780199389414.013.168>
- Motuzaitė-Matuzeviciute, G., Staff, R.A., Hunt, H.V., Liu, X., Jones, M.K., 2013. The early chronology of broomcorn millet (*Panicum miliaceum*) in Europe. *Antiquity*, 87, 1073-1085. <https://doi.org/10.1017/S0003598X00049875>
- Motuzaitė-Matuzeviciute, G., Lightfoot, E., O'Connell, T.C., Voyakin, D., Liu, X., Loman, V., Svyatko, S., Usmanova, E., Jones, M.K., 2015. The extent of cereal cultivation among the Bronze Age to Turkic period societies of Kazakhstan determined using stable isotope analysis on bone collagen. *Journal of Archaeological Science*, 59, 23-34. <https://doi.org/10.1016/j.jas.2015.03.029>
- Neef, R., 2001. Getreide im Silokomplex an der Poternenmauer (Boğazköy) – erste Aussagen zur Landwirtschaft. *Archäologischer Anzeiger* 2001/3, 335-341.
- Nesbitt, M. and Summers, G.D., 1988. Some recent discoveries of millet (*Panicum miliaceum* L. and *Setaria italica* (L.) P. Beauv.) at excavations in Turkey and Iran. *Anatolian Studies*, 38, 85-97. <https://doi.org/10.2307/3642844>
- Nitsch, E., Andreu, S., Creuzieux, A., Gardeisen, A., Halstead, P., Isaakidou, V., Karathanou, A., Kotsachristou, D., Nikolaidou, D., Papanthimou, A., Petridou, C., Triantaphyllou, S., Valamoti, S.M., Vasileiadou, A., Bogaard, A., 2017. A bottom-up view of food surplus: using stable carbon and nitrogen isotope analysis to investigate agricultural strategies and diet at Bronze Age Archontiko and

- Thessaloniki Toumba, northern Greece. *World Archaeology*, 49 (1), 105-137. <https://doi.org/10.1080/00438243.2016.1271745>
- Palmer, C., 2007. Carbonized plant macrofossils from Velim Skalka: preliminary observations. In: A. Harding, R. Šumberová, C. Knüsel, A. Outram, eds., *Velim: Violence and Death in Bronze Age Bohemia*. Prague: Institute of Archaeology of the Academy of Sciences of the Czech Republic, 137-141.
- Palmer, C. and Van der Veen, M., 2002. Archaeobotany and the social context of food. *Acta Palaeobotanica*, 42 (2), 195-202.
- Papathanasiou, A., 2015. Stable isotope analysis in Neolithic and Bronze Age Greece: an overview. In: A. Papathanasiou, M.P. Richards, S.C. Fox, eds., *Archaeodiet in the Greek World: Dietary Reconstruction from Stable Isotope Analysis*. Hesperia Supplementary Volume 49. Athens: The American School of Classical Studies at Athens, 25-55.
- Pare, C.F.E., 2004. Die Wagen der Bronzezeit in Mitteleuropa. In: M. Fansa and S. Burmeister, eds., *Rad und Wagen: Der Ursprung einer Innovation: Wagen im Vorderen Orient und Europa*. Mainz: Von Zabern, 355-372.
- Pashkevych, G., 2012. Environment and economic activities of Neolithic and Bronze Age populations of the northern Pontic area. *Quaternary International*, 261, 176-182. <https://doi.org/10.1016/j.quaint.2011.01.024>
- Perego, R., 2017. Contribution to the Development of the Bronze Age Plant Economy in the Surrounding of the Alps: an Archaeobotanical Case Study of Two Early and Middle Bronze Age Sites in Northern Italy (Lake Garda Region). Doctoral dissertation, University of Basel.
- Petroutsas, E.I. and Manolis, S.K., 2010. Reconstructing Late Bronze Age diet in mainland Greece using stable isotope analysis. *Journal of Archaeological Science*, 37, 614-620. <https://doi.org/10.1016/j.jas.2009.10.026>
- Popov, H., Marinova, E., Hristova, I., Iliev, S., 2018. Plant food from the Late Bronze and Early Iron Age hilltop site Kush Kaya, eastern Rhodope Mountains, Bulgaria: insights on the cooking practices. In: M. Ivanova, B. Athanassov, V. Petrova, D. Takorova, P.W. Stockhammer, eds., *Social Dimensions of Food in the Prehistoric Balkans*. Oxford: Oxbow, 263-277.
- Pospieszny, Ł., Makarowicz, P., Lewis, J., Górski, J., Taras, H., Włodarczak, P., Szczepanek, A., Ilchyshyn, V., Jagodinska, M.O., Czebreszuk, J., Muzolf, P., Nowak, M., Polańska, M., Juras, A., Chyleński, M., Wójcik, I., Lasota-Kuś, A., Romaniszyn, J., Tunia, K., Przybyła, M.M., Grygiel, R., Matoga, A., Makowiecki, A., Goslar, T., 2021. Isotopic evidence of millet consumption in the Middle Bronze Age of east-central Europe. *Journal of Archaeological Science*, 126, 105292. <https://doi.org/10.1016/j.jas.2020.105292>.
- Privat, K., Sobotkova, A., Bakardzhiev, S., Russeva, V., 2018. Excavation and palaeodietary analysis of Bronze Age human remains from Boyanovo, Yambol Province. In: S.A. Ross, A. Sobotkova, J. Tzvetkova, G. Nekhrizov, S. Connor, eds., *The Tundzha Regional Archaeology Project: Surface Survey, Palaeoecology, and Associated Studies in Central and Southeast Bulgaria, 2009-2015 Final Report*. Oxford: Oxbow, 182-190.
- Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., van der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S., 2020. The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0-55 cal kBP). *Radiocarbon*, 62 (4), 725-757. <https://doi.org/10.1017/RDC.2020.41>
- Richards, M.P. and Pearson, J.A., 2005. Stable isotope evidence of diet at Çatalhöyük. In: I. Hodder, ed., *Inhabiting Çatalhöyük: Reports from the 1995-99 Seasons*:

- Çatalhöyük Project Vol. 4*. Cambridge: McDonald Institute Monographs/London: British Institute of Archaeology at Ankara, 313-321.
- Richards, M.P. and Vika, E., 2008. Stable isotope results from new sites in the Peloponnese: Sykia, Kalamaki and Spaliareika. In: H. Martlew, Y. Tzedakis, M. Jones, eds., *Archaeology Meets Science: Biomolecular and Site investigations in Bronze Age Greece*. Oxford: Oxbow, 231-235.
- Riehl, S., 1999. *Bronze Age Environment and Economy in the Troad: the Archaeobotany of Kumtepe and Troy*. BioArchaeologica 2. Tübingen: Mo Vince Verlag.
- Rouse, L.M., 2020. Silent partners: archaeological insights on mobility, interaction and civilization in central Asia's past. *Central Asian Survey*, 39 (3), 398-419. <https://doi.org/10.1080/02634937.2020.1769024>
- Rühl, L., Herbig, C., Stobbe, A., 2015. Archaeobotanical analysis of plant use at Kamennyi Ambar, a Bronze Age fortified settlement of the Sintashta culture in the southern Trans-Urals steppe, Russia. *Vegetation History and Archaeobotany*, 24, 413-426. <https://doi.org/10.1007/s00334-014-0506-7>
- Ryabogina, N.E. and Ivanov, S.N., 2011. Ancient agriculture in western Siberia: problems of argumentation, paleoethnobotanic methods, and analysis of data. *Archaeology, Ethnology and Anthropology of Eurasia*, 39 (4), 96-106. <https://doi.org/10.1016/j.aear.2012.02.011>
- Sagona, A., 2018. *The Archaeology of the Caucasus*. Cambridge: Cambridge University Press.
- Šálková, T., Chvojka, O., Hlášek, D., Jiřík, J., John, J., Novák, J., Kovačikova, L., Beneš, J., 2019. Crops along the trade routes? Archaeobotany of the Bronze Age in the region of south Bohemia (Czech Republic) in context with longer distance trade and exchange networks. *Archaeological and Anthropological Sciences*, 11, 5569-5590. <https://doi.org/10.1007/s12520-019-00893-6>
- Sava, E., 2005. Viehzucht und Ackerbau in der Noua-Sabatinovka-Kultur. In: B. Horejs, R. Jung, E. Keiser, B. Teržan, eds., *Interpretationsraum Bronzezeit: Festschrift Bernhard Hänsel*. Bonn: Dr. Rudolf Habelt GmbH, 143-159.
- Sava, E. and Kaiser, E., 2011. Поселение с «золянками» у села Одая-Мичурин, Республика Молдова (археологические и естественнонаучные исследования) / *Die Siedlung mit, Aschelhügeln beim Dorf Odaia-Miciurin, Republik Moldova (archäologische und naturwissenschaftliche Untersuchungen)*. Tyragetia XIX. Chişinău: Muzeul Național de Arheologie și Istorie a Moldovei.
- Schirrmacher, J., Kneisel, J., Knitter, D., Hamer, W., Hinz, M., Schneider, R.R., Weinelt, M., 2020. Spatial patterns of temperature, precipitation, and settlement dynamics on the Iberian Peninsula during the Chalcolithic and the Bronze Age. *Quaternary Science Reviews*, 233, 106220. <https://doi.org/10.1016/j.quascirev.2020.106220>
- Sherratt, A., 1981. Plough and pastoralism: aspects of the secondary products revolution. In: I. Hodder, G. Isaac, N. Hammond, eds., *Pattern of the Past: Studies in Honour of David Clarke*. Cambridge: Cambridge University Press, 261-305.
- Sherratt, A., 1993. What would a Bronze Age world system look like? Relations between temperate Europe and the Mediterranean in later prehistory. *Journal of European Archaeology*, 1 (2), 1-58. <https://doi.org/10.1179/096576693800719293>
- Sherratt, A., 1999. Cash-crops before cash: organic consumables and trade. In: C. Gosden and J. Hather, eds., *The Prehistory of Food: Appetites for Change*. London: Routledge, 13-34.
- Sherratt, A. and Sherratt, S., 1998. Small worlds: interaction and identity in the ancient Mediterranean. In: E.H. Cline and D. Harris-Cline, eds., *The Aegean and the Orient in the Second Millennium*. Austin: University of Texas Press, 329-342.
- Spengler, R., Frachetti, M.D., Doumani, P., Rouse, L., Cerasetti, B., Bullion, E., Mar'yashev, A., 2014. Early agriculture and crop transmission among Bronze Age mobile pastoralists of central Eurasia. *Proceedings of the Royal Society, B: Biological Sciences*, 281, 20133382. <https://doi.org/10.1098/rspb.2013.3382>

- Spengler, R.N., Ryabogina, N., Tarasov, P.E., Wagner, M., 2016. The spread of agriculture into northern central Asia: timing, pathways, and environmental feedbacks. *The Holocene*, 26, 1527-1540. <https://doi.org/10.1177%2F0959683616641739>
- Spengler, R.N., de Nigris, I., Cerasetti, B., Carra, M., Rouse, L.M., 2018. The breadth of dietary economy in Bronze Age central Asia: case study from Adjı Kui 1 in the Murghab region of Turkmenistan. *Journal of Archaeological Science: Reports*, 22, 372-381. <https://doi.org/10.1016/j.jasrep.2016.03.029>
- Stevens, C.J., Murphy, C., Roberts, R., Lucas, L., Silva, F., Fuller, D.Q., 2016. Between China and South Asia: a middle Asian corridor of crop dispersal and agricultural innovation in the Bronze Age. *The Holocene*, 26, 1541-1555. <https://doi.org/10.1177%2F0959683616650268>
- Stika, H.-P. and Heiss, A.G., 2013. Plant cultivation in the Bronze Age. In: H. Fokkens and A. Harding, eds., *Oxford Handbook of the European Bronze Age*. Oxford: Oxford University Press, 340-361.
- Styring, A.K., Höhn, A., Linseele, V., Neumann, K., 2019. Direct evidence for agricultural intensification during the first two millennia AD in northeast Burkina Faso. *Journal of Archaeological Science*, 108, 104976. <https://doi.org/10.1016/j.jas.2019.104976>
- Suchowska-Ducke, P., Reiter, S.S., Vandkilde, H., eds., 2015. *Forging Identities: The Mobility of Culture in Bronze Age Europe*. Report from a Marie Curie Project 2009-2012 with Concluding Conference at Aarhus University, Moesgaard 2012. BAR International Series 2771. Oxford: BAR Publishing.
- Tafuri, A.M., Craig, O.E., Canci A., 2009. Stable isotope evidence for the consumption of millet and other plants in Bronze Age Italy. *American Journal of Physical Anthropology*, 139, 146-153. <https://doi.org/10.1002/ajpa.20955>
- Tafuri, M.A., Rottoli, M., Cupitò, M., Pulcini, M.L., Tasca, G., Carrara, N., Bonfanti F., Salzani, L., Canci, A., 2018. Estimating C4 plant consumption in Bronze Age north-eastern Italy through stable carbon and nitrogen isotopes in bone collagen. *International Journal of Osteoarchaeology*, 28, 131-142. <https://doi.org/10.1002/oa.2639>
- Triantaphyllou, S., 2015. Stable isotope analysis of skeletal assemblages from prehistoric northern Greece. In: A. Papathanasiou, M.P. Richards, S.C. Fox, eds., *Archaeodiet in the Greek World: Dietary Reconstruction from Stable Isotope Analysis*. Hesperia Supplementary Volume 49. Athens: The American School of Classical Studies at Athens, 57-75.
- Trifonov, V.A., Shishlina, N.I., Yu Lebedeva, E., van der Plicht, J., Rishko, S.A., 2017. Directly dated broomcorn millet from northwestern Caucasus: tracing the Late Bronze Age route into the Russian steppe. *Journal of Archaeological Science: Reports*, 12, 288-294. <https://doi.org/10.1016/j.jasrep.2017.02.004>
- Twiss, K.C., 2019. *The Archaeology of Food: Identity, Politics, and Ideology in the Prehistoric and Historic Past*. Cambridge: Cambridge University Press.
- Ulaş, B., 2021. Traditional wheat cultivation in south-eastern Anatolia and its comparison to the archaeological context. *Genetic Resources and Crop Evolution*, 68, 151-184. <https://doi.org/10.1007/s10722-020-00977-6>
- Valamoti, S.M., 2016. Millet, the late comer: on the tracks of *Panicum miliaceum* in prehistoric Greece. *Archaeological and Anthropological Sciences*, 8, 51-63. <https://doi.org/10.1007/s12520-013-0152-5>
- Van der Veen, M., 2003. When is food a luxury? *World Archaeology*, 34 (3), 405-427. <https://doi.org/10.1080/0043824021000026422>
- Van der Veen, M., 2010. Agricultural innovation: invention and adoption or change and adaptation? *World Archaeology*, 42 (1), 1-12. <http://dx.doi.org/10.1080/00438240903429649>
- Van Hoof, L., Dally, O., Schlöfel, M., 2012. Staying home or staying with your cattle? Different reactions to environmental changes in the Late Bronze Age of the lower Don area (southern Russia). In: W. Bebermeier, R. Hebenstreit, E. Kaiser, J. Krause, eds., *Landscape Archaeology*. eTopoi Journal for Ancient Studies 3, 71-75.

- Vandkilde, H., 2014. Breakthrough of the Nordic Bronze Age: transcultural warriorhood and a Carpathian crossroad in the sixteenth century BC. *European Journal of Archaeology*, 17 (4), 602-633. <https://doi.org/10.1179/1461957114Y.00000000064>
- Ventresca Miller, A.R. and Makarewicz, C.A., 2019. Intensification in pastoralist cereal use coincides with the expansion of trans-regional networks in the Eurasian steppe. *Scientific Reports*, 9, 8363. <https://doi.org/10.1038/s41598-018-35758-w>
- Weiberg, E., Bevan, A., Kouli, K., Katsianis, M., Woodbridge, J., Bonnier, A., Engel, M., Finné, M., Fyfe, R., Maniatis, Y., Palmisiano, A., Panajiotidis, S., Roberts, N.C., Shennan, S., 2019. Long-term trends of land use and demography in Greece: a comparative study. *The Holocene*, 29 (5), 742-760. <https://doi.org/10.1177%2F0959683619826641>
- Wang, T., Wei, D., Chang, X., Yu, Z., Zhang, X., Wang, C., Hu, Y., Fuller, B.T., 2017. Tian-shanbeilu and the Isotopic Millet Road: reviewing the Late Neolithic/Bronze Age radiation of human millet consumption from north China to Europe. *National Science Review*, 6 (5), 1024-1039. <https://doi.org/10.1093/nsr/nwx015>
- Weiss, H. ed., 2017. *Megadrought and Collapse: from Early Agriculture to Angkor*. New York: Oxford University Press.
- Wardle, K., Higham, T., Kromer, B., 2014. Dating the end of the Greek Bronze Age: a robust radiocarbon-based chronology from Assiros Toumba. *PLoS ONE*, 9 (9), e106672. <https://doi.org/10.1371/journal.pone.0106672>
- Weber, S.A. and Fuller, D.Q., 2008. Millets and their role in early agriculture. *Prāgdhārā*, 18, 69-90.
- Weeks, L.R., 2003. *Early Metallurgy of the Persian Gulf: Technology, Trade, and the Bronze Age World*. Boston: Brill Academic Publishers.
- Yattoo, M.A., Spate, M., Betts, A., Pokharia, A.K., Shah, M.A., 2020. New evidence from the Kashmir valley indicates the adoption of East and West Asian crops in the western Himalayas by 4400 years ago. *Quaternary Science Advances*, 2, 100011. <https://doi.org/10.1016/j.qsa.2020.100011>
- Лебедева, Е.Ю., 2005. Археоботаника и изучение земледелия эпохи бронзы в Восточной Европе [Archeobotany and the study of Bronze Age agriculture in Eastern Europe]. *OPUS: Междисциплинарные исследования в археологии*, 4, 50-68.
- Лебедева, Е.Ю., 2011. Первые результаты археоботанических исследований на археологических памятниках Адыгеи [The first results of archaeobotanical research at the archaeological sites of Adygea]. *Аналитические исследования лаборатории естественнонаучных методов*, 2, 244 -257.

List of authors (in the order of appearance):

Wiebke Kirleis

Institute of Prehistoric and Protohistoric
Archaeology
Kiel University
Johanna-Mestorf-Str. 2-6, D-24118 Kiel, Germany
wiebke.kirleis@ufg.uni-kiel.de

Dragana Filipović

Institute of Prehistoric and Protohistoric
Archaeology
Kiel University
Johanna-Mestorf-Str. 2-6, D-24118 Kiel, Germany
d.filipovic@ufg.uni-kiel.de

Marta Dal Corso

Institute of Prehistoric and Protohistoric
Archaeology
Kiel University
Johanna-Mestorf-Str. 2-6, D-24118 Kiel, Germany
mdalcorso@ufg.uni-kiel.de

Galyna Pashkevych

National Museum of Natural Sciences
National Academy of Sciences in Ukraine
Bul. Bohdan Khmelnytsky 15, 01030 Kyiv, Ukraine
galina.pashkevich@gmail.com

Sofia Filatova

Institute of Prehistoric and Protohistoric
Archaeology
Kiel University
Johanna-Mestorf-Str. 2-6, D-24118 Kiel, Germany
s.filatova@ufg.uni-kiel.de

László Bartosiewicz

Osteoarchaeological Research Laboratory
Department of Archaeology and Classical Studies
Stockholm University
Lilla Frescativägen 7, 106 91 Stockholm, Sweden
laszlo.bartosiewicz@ofl.su.se

Kelly Reed

Oxford Martin School
University of Oxford
34 Broad Street, Oxford OX1 3BD, UK
kellyreed@hotmail.co.uk

Jacqueline Balen

Archaeological Museum in Zagreb
19 Nikola Šubić Zrinski Square, P.O. Box 13, 10000
Zagreb, Croatia
jbalen@amz.hr

Ivan Drnić

Archaeological Museum in Zagreb
19 Nikola Šubić Zrinski Square, P.O. Box 13, 10000
Zagreb, Croatia
idrnic@amz.hr

Sara Essert

Department of Biology
University of Zagreb
Rooseveltova trg 6, 10000 Zagreb, Croatia
sara.essert@biol.pmf.hr

Hrvoje Kalafatić

Institute of Archaeology
Jurjevska 15, 10000 Zagreb, Croatia
hrvoje.kalafatic@iarh.hr

Marija Mihaljević

Nova Gradiška Municipal Museum
Trg kralja Tomislava 7, 35400 Nova Gradiška, Croatia
marija.mihaljevic@gmng.hr

Emily Zavodny

Department of Anthropology
University of Florida
Gainesville, FL 32611, USA
ezavodny@ufl.edu

Tjaša Tolar

Scientific Research Centre of the Slovenian Academy of Sciences and Arts (ZRC SAZU)
Institute of Archaeology
Novi trg 2, 1000 Ljubljana, Slovenia
tjasa.tolar@zrc-sazu.si

Primož Pavlin

Scientific Research Centre of the Slovenian Academy of Sciences and Arts (ZRC SAZU)
Institute of Archaeology
Novi trg 2, 1000 Ljubljana, Slovenia
primoz.pavlin@zrc-sazu.si

Monika Hellmund

Office for Heritage Management and Archaeology
Saxony-Anhalt – State Museum
of Prehistory in Halle (Saale)
Richard-Wagner-Str. 9, D-06114 Halle (Saale), Germany
mhellmund@lda.stk.sachsen-anhalt.de

Françoise Toulemonde

67 rue des Sablons, 78750 Mareil-Marly, France
*Associate member of UMR 7209 MNHN-CNRS, Archéozoologie, Archéobotanique: Sociétés, Pratiques et Environnements (AASPE)
Paris, France
francoise.toulemonde@gmail.com

Julian Wiethold

Institut national de recherches archéologiques préventives (Inrap)
Centre de recherches archéologiques de Metz, Laboratoire archéobotanique
12 rue de Méric, 57063 Metz Cedex 2, France
&
UMR 6298 ArteHiS
6 Boulevard Gabriel, 21000 Dijon, France
julian.wiethold@inrap.fr

Emmanuelle Bonnaire

Archéologie Alsace
11 rue Champollion, 67600 Sélestat, France
*Associate member of UMR 7209 MNHN-CNRS, Archéozoologie, Archéobotanique: Sociétés, Pratiques et Environnements (AASPE)
Paris, France
emmanuelle.bonnaire@archeologie.alsace

Geneviève Daoulas

Institut national de recherches archéologiques préventives (Inrap)

Centre de recherches archéologiques de Châlons-en-Champagne,
38 rue des Dats, 51520 Saint-Martin-sur-Le-Pré, France
&
UMR 7209 MNHN-CNRS, Archéozoologie, Archéobotanique: Sociétés, Pratiques et Environnements (AASPE)
Paris, France
genevieve.daoulas@inrap.fr

Marie Derreumaux

Centre de Recherche Archéologique de la Vallée de l'Oise (CRAVO)
17 rue James de Rothschild, 60200 Compiègne, France
&
UMR 7209 MNHN-CNRS, Archéozoologie, Archéobotanique: Sociétés, Pratiques et Environnements (AASPE)
Paris, France
mariecarpo@free.fr

Frédérique Durand

Institut national de recherches archéologiques préventives (Inrap)
ZI Les Pinsons, 13 rue du Négoce 31650 Saint-Orens-de-Gameville, France &
UMR 5608 TRACES
Toulouse, France
frederique.durand@inrap.fr

Bénédicte Pradat

Institut national de recherches archéologiques préventives (Inrap)
148 avenue Maginot, 37100 Tours, France
&
UMR 7209 MNHN-CNRS, Archéozoologie, Archéobotanique: Sociétés, Pratiques et Environnements (AASPE)
Paris, France
benedicte.pradat@inrap.fr

Oriane Rousselet

Bibracte, Centre archéologique européen
58370 Glux-en-Glenne
o.rousselet@bibracte.fr

Caroline Schaal

Université Bourgogne-Franche-Comté
16 route de Gray, 25000 Besançon, France
UMR 6249 Chrono-environnement
25000 Besançon, France
Caroline.Schaal@univ-fcomte.fr

Véronique Zech-Matterne

Centre National de la Recherche Scientifique (CNRS)
 UMR 7209 MNHN-CNRS, Archéozoologie,
 Archéobotanique: Sociétés, Pratiques et
 Environnements (AASPE)
 55 rue Buffon, 75005 Paris, France
 veronique.zech@mnhn.fr

Ingo Feeser

Institute of Prehistoric and Protohistoric Archaeology
 Kiel University
 Johanna-Mestorf-Str. 2-6, D-24118 Kiel, Germany
 ifeeser@ufg.uni-kiel.de

Stefanie Schaefer-Di Maida

Institute of Prehistoric and Protohistoric Archaeology
 Kiel University
 Johanna-Mestorf-Str. 2-6, D-24118 Kiel, Germany
 sschaefer@ufg.uni-kiel.de

Stefan Dreibrodt

Institute for Ecosystem Research
 Kiel University
 Olshausenstrasse 75, D-24118 Kiel, Germany
 sdreibrodt@ecology.uni-kiel.de

Jutta Kneisel

Institute of Prehistoric and Protohistoric Archaeology
 Kiel University
 Johanna-Mestorf-Str. 2-6, D-24118 Kiel, Germany
 jutta.kneisel@ufg.uni-kiel.de

Edward A. Standall

BioArch, Department of Archaeology
 University of York
 Wentworth Way, York YO10 5DD, UK & Department
 of Scientific Research
 British Museum
 Great Russell Street, London WC1B 3DG, UK
 EStandall@britishmuseum.org

Oliver E. Craig

BioArch, Department of Archaeology
 University of York
 Wentworth Way, York YO10 5DD, UK
 Oliver.Craig@york.ac.uk

Carl Heron

Department of Scientific Research
 The British Museum
 Great Russell Street, London WC1B 3DG, UK
 cheron@britishmuseum.org

Marco Zanon

Institute of Prehistoric and Protohistoric Archaeology
 Kiel University
 Johanna-Mestorf-Str. 2-6, D-24118 Kiel, Germany
 marco@zanon.xyz

Mauro Rottoli

Dipartimento di Lettere e Filosofia
 Università degli Studi di Trento
 Via Tommaso Gar 14, 38122 Trento, Italy &
 Laboratorio di Archeobiologia – Musei Civici di Como
 Piazza Medaglie d'Oro 6, 22100 Como, Italy
 archeobotanica@gmail.com

Michele Cupitò

Dipartimento dei Beni Culturali
 Università degli Studi di Padova
 Piazza Capitaniato 7, 35139 Padova, Italy
 michele.cupito@unipd.it

Elisa Dalla Longa

Dipartimento dei Beni Culturali
 Università degli Studi di Padova
 Piazza Capitaniato 7, 35139 Padova, Italy
 elisa.dallalonga@unipd.it

Eiko Endo

Center for Obsidian and Lithic Studies
 Meiji University
 1-1 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-0064,
 Japan
 endosalt@yahoo.co.jp

Andrés Teira-Brión

School of Archaeology
 University of Oxford
 36 Beaumont Street, Oxford, OX1 2PG, UK
 andres.teirabrion@arch.ox.ac.uk

STPAS: Scales of Transformation in Prehistoric and Archaic Societies

The book series 'Scales of Transformation in Prehistoric and Archaic Societies' (STPAS) is an international scientific series that covers major results deriving from or being associated with the research conducted in the Collaborative Research Centre 'Scales of Transformation: Human-Environmental Interaction in Prehistoric and Archaic Societies' (CRC 1266). Primarily located at Kiel University, Germany, the CRC 1266 is a large interdisciplinary project investigating multiple aspects of socio-environmental transformations in ancient societies between 15,000 and 1 BCE across Europe.

Volume 1

Das Jungneolithikum in Schleswig-Holstein

Sebastian Schultrich | 2018

ISBN: 9789088907425

Format: 210x280mm | 506 pp. | Language: German | 43 illus. (bw) | 103 illus. (fc)

Keywords: Late Neolithic, Single Grave Culture, Corded Ware Culture, transformation, solid stone axe, battle axe, fragments of axes | Jungneolithikum, Einzelgrabkultur, Schnurkeramische Kultur, Transformation, Felsgesteinäxte, Streitäxte, Axtfragmente

Volume 2

Embracing Bell Beaker

Adopting new ideas and objects across Europe during the later 3rd millennium BC (c. 2600-2000 BC)

Jos Kleijne | 2019

ISBN: 9789088907555

Format: 210x280mm | 300 pp. | Language: English | 91 illus. (fc)

Keywords: archaeology; Late Neolithic; Bell Beaker phenomenon; settlement archaeology; innovation; network analysis; mobility; prehistoric potter

Volume 3

Habitus?

The Social Dimension of Technology and Transformation

Edited by Sławomir Kadrow & Johannes Müller | 2019

ISBN: 9789088907838

Format: 210x280mm | 232 pp. | Language: English | 15 illus. (bw) | 65 illus. (fc)

Keywords: European prehistory; archaeology; habitus; technology; transformation; social dimension; ethnoarchaeology

Volume 4

How's Life?

Living Conditions in the 2nd and 1st Millennia BCE

Edited by Marta Dal Corso, Wiebke Kirleis, Jutta Kneisel, Nicole Taylor, Magdalena Wieckowska-Lüth, Marco Zanon | 2019

ISBN: 9789088908019

Format: 210x280mm | 220 pp. | Language: English | 29 illus. (bw) | 43 illus. (fc)

Keywords: Bronze Age, domestic archaeology, household archaeology, daily life, routine activities, diet, waste, violence, health, natural resources, food production

Volume 5

Megalithic monuments and social structures

Comparative studies on recent and Funnel Beaker societies

Maria Wunderlich | 2019

ISBN: 9789088907869

Format: 210x280mm | 382 pp. | Language: English | 114 illus. (bw) | 246 illus. (fc)

Keywords: Megalithic graves, monumentality, Funnel Beaker Complex, ethnoarchaeology, Sumba, Nagaland, social organisation, cooperation

Volume 6

Gender Transformations in Prehistoric and Archaic Societies

Edited by Julia Katharina Koch & Wiebke Kirleis | 2019

ISBN: 9789088908217

Format: 210x280mm | 502 pp. | Language: English | 114 illus. (bw) | 58 illus. (fc)

Keywords: academic fieldwork; gender archaeology; social archaeology; environmental archaeology; history of archaeology; Mesolithic; Neolithic; Bronze Age; Iron Age; Europe; South-west Asia; Central Asia

Volume 7

Maidanets'ke

Development and decline of a Trypillia mega-site in Central Ukraine

René Ohlrau | 2020

ISBN: 9789088908484

Format: 210x280mm | 326 pp. | Language: English | 141 illus. (bw) | 93 illus. (fc)

Keywords: settlement archaeology; prehistoric archaeology; early urbanism; geophysical survey; paleodemography; Trypillia; mega-site

Volume 8

Detecting and explaining technological innovation in prehistory

Edited by Michela Spataro & Martin Furholt | 2020

ISBN: 9789088908248

Format: 210x280mm | 248 pp. | Language: English | 22 illus. (bw) | 37 illus. (fc)

Keywords: archaeology; prehistory; technology; innovation; invention; tradition; chaîne opératoire; knowledge acquisition; knowledge transfer; Neolithic; Bronze Age; Iron Age; ethnography; ceramic; metal; bone

Volume 9

Archaeology in the Žitava Valley I

The LBK and Želiezovce settlement site of Vráble

Edited by Martin Furholt, Ivan Cheben, Johannes Müller, Alena Bistáková,

Maria Wunderlich & Nils Müller-Scheeßel | 2020

ISBN: 9789088908972

Format: 210x280 | 546 pp. | Language: English | 50 illus. (bw) | 157 illus. (fc)
 Keywords: European Early Neolithic; LBK, settlement patterns; social organization; social conflict; village and neighbourhood structures; burial rites; enclosure; excavation report

Volume 10

Hellenistic Architecture and Human Action

A Case of Reciprocal Influence

Edited by Annette Haug & Asja Müller | 2020

ISBN: 9789088909092

Format: 210x280mm | 208 pp. | Language: English | 38 illus. (bw) | 29 illus. (fc)

Keywords: Classical archaeology; Hellenistic architecture; agency; perception

Volume 11

Interdisciplinary analysis of the cemetery Kudachurt 14

Evaluating indicators of social inequality, demography, oral health and diet during the Bronze Age key period 2200-1650 BCE in the Northern Caucasus

Katharina Fuchs | 2020

ISBN: 9789088909030

Format: 210x280mm | 406 pp. | Language: English | 25 illus. (bw) | 137 illus. (fc)

Keywords: North Caucasian archaeology; Bronze Age; burial practice; social inequality; human remains; palaeopathology; oral health; C and N stable isotopes

Volume 12

Tripolye Typo-chronology

Mega and Smaller Sites in the Sinyukha River Basin

Liudmyla Shatilo | 2021

ISBN: 9789088909511

Format: 210x280mm | 422 pp. | Language: English | 100 illus. (bw) | 90 illus. (fc)

Keywords: Prehistoric archaeology; Tripolye; chronology; mega-sites; Sinyukha River Basin; pottery; carbon dating; typochronology; Eastern Europe

Volume 13

Vom Kollektiv zum Individuum

Transformationsprozesse am Übergang vom 4. zum 3. Jahrtausend v. Chr. in der Deutschen Mittelgebirgszone

Clara Drummer | 2022

ISBN: 9789464270129

Format: 210x280mm | 343 pp. | Language: German | 53 illus. (bw) | 95 illus. (fc)

Keywords: Neolithic; Germany; Lower Mountain Range; migrations; identities; social transformations; Corded Ware; Bell Beaker; funerary archaeology; ancient DNA

Volume 14

Millet and What Else?

The Wider Context of the Adoption of Millet Cultivation in Europe

Edited by Wiebke Kirleis, Marta Dal Corso & Dragana Filipović | 2022

ISBN: 9789464270150

Format: 210x280mm | 328 pp. | Scales of Transformation 14 | Series: Scales of

Transformation | Language: English | 61 illus. (bw) | 13 illus. (fc) | Keywords:

archaeology; millet; Europe prehistory; Bronze Age; archaeobotany; zooarchaeology; miliacin; prehistoric agriculture

MILLET AND WHAT ELSE?

The wider context of the adoption of millet cultivation in Europe

Broomcorn/common/proso millet (*Panicum miliaceum*) is a cereal crop that originated in East Asia and was transferred westward to Europe, where it was introduced in the mid-2nd millennium BCE, at the height of the Bronze Age. Archaeobotanists from the Collaborative Research Centre 1266, supported by many colleagues, conducted a large-scale programme of radiocarbon dating of millet grains from prehistoric Europe. They discovered that the spread of this crop on the continent happened quickly, extending far and wide.

What do we know about the (pre)history of this crop in Europe? The workshop organised by the CRC 1266 at Kiel University in 2019 encouraged the discussion on the circumstances and consequences of early broomcorn millet cultivation in Europe. This book brings together many of the workshop papers and reflects the diverse topics and research areas covered. The contributions inform us on the range of cultivated and collected plants from the time before and after the start of millet cultivation in Europe; present the cultural setting in which millet arrived; discuss possible reasons driving the acceptance of this innovation; and reconstruct possible uses of millet and the methods of its cultivation, processing and storage. Not just the plant economy, but also the animal economy is represented, since millet was and is grown for both humans and animals. Techniques used to trace millet archaeologically are continually being developed or improved, and this book describes the application of a few of them.

This broad-based compilation of papers adds another layer to the dynamic picture of the Bronze Age and the interconnected continent. It also illustrates the complexity of the research on the diffusion of agricultural innovations.



sidestonepress

ISBN: 978-94-6427-015-0



9 789464 270150 >